

# Eigenelements of a General Aggregation-Fragmentation Model.

Marie Doumic <sup>\*</sup>      Pierre Gabriel <sup>†</sup>

June 12, 2009

## Abstract

We consider a linear integro-differential equation which arises to describe both aggregation-fragmentation processes and cell division. We prove the existence of a solution  $(\lambda, \mathcal{U}, \phi)$  to the related eigenproblem, where  $\lambda$  denotes its first eigenvalue. Our study concerns a non-constant transport term that can vanish at  $x = 0$ . Non lower-bounded transport terms bring difficulties to find *a priori* estimates. We use weighted-norms to solve this problem.

**Keywords** Aggregation-fragmentation equations, eigenproblem, size repartition, polymerization process, cell division, long-time asymptotic.

**AMS Class. No.** 35A05, 35B40, 45C05, 45K05, 82D60, 92D25

## 1 Introduction

Competition between growth and fragmentation is a common phenomenon for a structured population. It arises for instance in a context of cell division (see, among many others, [22, 1, 4, 5, 6, 14, 30, 18, 20]), polymerization (see [7, 13]), telecommunication (see [2]) or neurosciences (see [27]). It is also a mechanism which rules the proliferation of prion's proteins (see [10, 19, 21]). These proteins are responsible of spongiform encephalopathies and appear in the form of aggregates in infected cells. Such polymers grow attaching non infectious monomers and converting them into infectious ones. On the other hand they increase their number by splitting. To describe such phenomenons, we write the following integro-differential equation, used in [10, 19, 21] to study the proliferation of prion,

$$\left\{ \begin{array}{l} \frac{\partial}{\partial t} u(x, t) + \frac{\partial}{\partial x} (\tau(x) u(x, t)) + \beta(x) u(x, t) = 2 \int_x^\infty \beta(y) \kappa(x, y) u(y, t) dy, \quad x \geq 0, \\ u(x, 0) = u_0(x), \\ u(0, t) = 0. \end{array} \right. \quad (1)$$

---

<sup>\*</sup>INRIA Rocquencourt, projet BANG, Domaine de Voluceau, BP 105, F-78153 Rocquencourt, France.

<sup>†</sup>Université Pierre et Marie Curie-Paris 6, UMR 7598 LJLL, BC187, 4, place de Jussieu, F-75252 Paris cedex 5; corresponding author, email: gabriel@ann.jussieu.fr

The aggregates density function  $u(x, t)$  represents the quantity of polymers of size  $x$  at time  $t$ . They aggregate monomers with the rate  $\tau(x)$ . Equation (1) also takes into account the fragmentation of a polymer of size  $y$  into two smaller polymers of size  $x$  and  $y - x$ . This fragmentation occurs with a rate  $\beta(y)$  and produce an aggregate of size  $x$  with the rate  $\kappa(x, y)$ . Equation (1) is a particular case of the more general one

$$\frac{\partial}{\partial t}u(x, t) + \frac{\partial}{\partial x}(\tau(x)u(x, t)) + [\beta(x) + \mu(x)]u(x, t) = n \int_x^\infty \beta(y)\kappa(x, y) u(y, t) dy, \quad x \geq x_0, \quad (2)$$

with the bound condition  $u(x_0, t) = 0$  (see [3, 10, 21]). In this model, polymers are broken in an average of  $n > 1$  smaller ones by the fragmentation process, there is a death term  $\mu(x) \geq 0$  representing degradation, and a minimal size of polymers  $x_0$  which can be positive. Our results remain true for this generalization.

To study the asymptotic growth of the quantity of polymers, we are looking at the eigenelements  $(\lambda, \mathcal{U}, \phi)$  of (1), *i.e.*, the solution of the equation

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x}(\tau(x)\mathcal{U}(x)) + (\beta(x) + \lambda)\mathcal{U}(x) = 2 \int_x^\infty \beta(y)\kappa(x, y)\mathcal{U}(y)dy, \quad x \geq 0, \\ \tau\mathcal{U}(x=0) = 0, \quad \mathcal{U}(x) \geq 0, \quad \int_0^\infty \mathcal{U}(x)dx = 1, \\ -\tau(x)\frac{\partial}{\partial x}(\phi(x)) + (\beta(x) + \lambda)\phi(x) = 2\beta(x) \int_0^x \kappa(y, x)\phi(y)dy, \quad x \geq 0, \\ \phi(x) \geq 0, \quad \int_0^\infty \phi(x)\mathcal{U}(x)dx = 1. \end{array} \right. \quad (3)$$

Here  $\lambda$  is the first eigenvalue,  $\mathcal{U}$  the corresponding eigenfunction and  $\phi$  the dual eigenfunction. Our parameters  $\beta$ ,  $\tau$  and  $\kappa$  are not necessarily regular. For the first equation (equation on  $\mathcal{U}$ ) we are looking for  $\mathcal{D}'$  solutions defined as follows :  $\mathcal{U} \in L^1(\mathbb{R}^+)$  is a  $\mathcal{D}'$  solution if  $\forall \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^+)$ ,

$$-\int_0^\infty \tau(x)\mathcal{U}(x)\partial_x\varphi(x) dx + \lambda \int_0^\infty \mathcal{U}(x)\varphi(x) dx = \int_0^\infty \beta(x)\mathcal{U}(x) \left( 2 \int_0^\infty \varphi(y)\kappa(y, x) dy - \varphi(x) \right) dx. \quad (4)$$

Concerning the dual equation, we are looking for a solution  $\phi \in W_{loc}^{1,\infty}(0, \infty)$  such that the equality holds in  $L_{loc}^1(0, \infty)$ , *i.e.* almost everywhere.

Existence and uniqueness of such elements has already been proved for general fragmentation kernels  $\kappa(x, y)$  and fragmentation rates  $\beta(x)$ , but with very particular polymerization rates  $\tau(x)$ , namely constant ( $\tau \equiv 1$  in [28]), homogeneous ( $\tau(x) = x^\mu$  in [23]) or with a compact support ( $Supp \tau = [0, x_M]$  in [14]). It seems however of deep interest to consider more general  $\tau$  as [10, 32] suggest. The proof of [28] can be adapted for non constant rates but still positive and bounded ( $0 < m < \tau(x) < M$ ). The paper [23] gives results for  $\tau(0) = 0$ , but for a very restricted class of shape for  $\tau$ . The paper [14] gives results for  $\tau$  with general shape in the case where there is also an age variable (integration in age then allows to recover Problem (1)), but requires a compact support and regular parameters. Here we consider polymerization rates that can vanish at  $x = 0$ , positive for  $x$  positive, with general shape and few regularity for the all parameters ( $\tau$ ,  $\beta$  and  $\kappa$ ), and we have to reformulate the space for solutions.

**Theorem 1 (Existence and Uniqueness)** *Under assumptions (5)-(13), there exists a unique solution  $(\lambda, \mathcal{U}, \phi)$  (in the sense we have defined before) to the eigenproblem (3) and we have*

$$\begin{aligned} \lambda &> 0, \\ x^\alpha \tau \mathcal{U} &\in L^p(\mathbb{R}^+), \quad \forall \alpha \geq -\gamma, \quad \forall p \in [1, \infty], \\ x^\alpha \tau \mathcal{U} &\in W^{1,1}(\mathbb{R}^+), \quad \forall \alpha \geq 0 \\ \exists k > 0 \text{ s.t. } &\frac{\phi}{1+x^k} \in L^\infty(\mathbb{R}^+), \\ \tau \frac{\partial}{\partial x} \phi &\in L_{loc}^\infty(\mathbb{R}^+). \end{aligned}$$

The end of this paper is devoted to define precisely the assumptions and prove this theorem. It is organized as follows : in Section 2 we describe the assumptions and give some examples of interesting parameters. In Section 3 we prove Theorem 1 using *a priori* bounds on weighted norms and then we give some consequences and perspectives in Section 4. The proof of technical lemmas and theorem can be found in the Appendix.

## 2 Coefficients

### 2.1 Assumptions

For all  $y \geq 0$ ,  $\kappa(\cdot, y)$  is a nonnegative measure with a support included in  $[0, y]$ . We define  $\kappa$  on  $(\mathbb{R}_+)^2$  as follows :  $\kappa(x, y) = 0$  for  $x > y$ . We assume that for all continuous function  $\psi$ , the application  $f_\psi : y \mapsto \int \psi(x) \kappa(x, y) dx$  is Lebesgue measurable.

The natural assumptions on  $\kappa$  (see [19] for the motivations) are that polymers can split only in two pieces which is taken into account by

$$\int \kappa(x, y) dx = 1. \tag{5}$$

So  $\kappa(y, \cdot)$  is a probability measure and  $f_\psi \in L_{loc}^\infty(\mathbb{R}^+)$ . The conservation of mass imposes

$$\int x \kappa(x, y) dx = \frac{y}{2}, \tag{6}$$

a property that is automatically satisfied for a symmetric fragmentation (*i.e.*  $\kappa(x, y) = \kappa(y-x, y)$ ) thanks to (5). For the more general model (2), assumption (6) becomes  $\int x \kappa(x, y) dx = \frac{y}{n}$  to preserve the mass conservation.

We also assume that the second moment of  $\kappa$  is less than the first one

$$\int \frac{x^2}{y^2} \kappa(x, y) dx \leq c < 1/2 \tag{7}$$

(it becomes  $c < 1/n$  for model (2)). We refer to the Examples for an explanation of the physical meaning.

For the polymerization and fragmentation rates  $\tau$  and  $\beta$ , we introduce the set

$$\mathcal{P} := \{f \geq 0 : \exists \mu, \nu \geq 0, \limsup_{x \rightarrow \infty} x^{-\mu} f(x) < \infty \text{ and } \liminf_{x \rightarrow \infty} x^\nu f(x) > 0\}$$

and the space

$$L_0^1 := \{f, \exists a > 0, f \in L^1(0, a)\}.$$

We consider

$$\tau, \beta \in L_{loc}^1(\mathbb{R}^+) \cap \mathcal{P} \quad (8)$$

satisfying

$$\forall K \text{ compact of } (0, \infty), \exists m_K > 0 \text{ s.t. } \tau(x) \geq m_K \text{ for a.e. } x \in K \quad (9)$$

and

$$\exists b \geq 0, \text{ Supp}\beta = [b, \infty). \quad (10)$$

Assumption (10) is necessary to prove uniqueness and existence for the adjoint problem.

To avoid shattering (zero-size polymers formation, see [3, 21]), we assume

$$\exists C > 0, \gamma \geq 0 \text{ s.t. } \int_0^x \kappa(z, y) dz \leq \min\left(1, C\left(\frac{x}{y}\right)^\gamma\right) \quad \text{and} \quad \frac{x^\gamma}{\tau(x)} \in L_0^1 \quad (11)$$

$$\frac{\beta}{\tau} \in L_0^1. \quad (12)$$

On the other hand, to avoid forming infinitely long polymers (gelation phenomenon, see [17, 16]), we assume

$$\lim_{x \rightarrow +\infty} \frac{x\beta(x)}{\tau(x)} = +\infty. \quad (13)$$

**Remark 1** *In case when (11) is satisfied for  $\gamma > 0$ , then (7) is automatically fulfilled (see Lemma 3 in the Appendix).*

## 2.2 Examples

First we give some examples of coefficients which satisfy or not our previous assumptions.

For the fragmentation kernel, we first check the assumptions (5) and (6). They are satisfied for autosimilar measures, namely  $\kappa(x, y) = \frac{1}{y}\kappa_0(\frac{x}{y})$ , with  $\kappa_0$  a probability measure on  $[0, 1]$ , symmetric in  $1/2$ . Now we exhibit some  $\kappa_0$ .

**General mitosis :**

$$\kappa_0^r = \frac{1}{2}(\delta_r + \delta_{1-r}) \quad \text{for} \quad r \in [0, 1/2]. \quad (14)$$

Assumption (11) is satisfied for any  $\gamma > 0$  in the cases when  $r \in (0, 1/2]$ . So (7) is also fulfilled thanks to Remark 1. The particular value  $r = 1/2$  leads to equal mitosis ( $\kappa(x, y) = \delta_{x=\frac{y}{2}}$ ).

The case  $r = 0$  corresponds to the renewal equation ( $\kappa(x, y) = \frac{1}{2}(\delta_{x=0} + \delta_{x=y})$ ). In this case, the size of the polymers does not decrease during the fragmentation process. It is typically what we want to avoid with assumption (7). For such a fragmentation kernel, assumption (11) is satisfied only for  $\gamma = 0$ ,

and the moments  $\int z^k \kappa_0(z) dz$  are equal to  $1/2$  for all  $k > 0$ , so (7) does not hold true. However, if we consider a convex combination of  $\kappa_0^0$  with an other kernel such as  $\kappa_0^r$  with  $r \in (0, 1/2]$ , then (11) remains false for any  $\gamma > 0$  but (7) is fulfilled. Indeed we have for  $\rho \in (0, 1)$

$$\int z^2 (\rho \kappa_0^0(z) + (1 - \rho) \kappa_0^r(z)) dz = \frac{\rho}{2} + \frac{1 - \rho}{2} (r^2 + (1 - r)^2) = \frac{1}{2} (1 - 2r(1 - r)(1 - \rho)) < \frac{1}{2}.$$

**Homogeneous fragmentation :**

$$\kappa_0^\alpha(z) = \frac{\alpha + 1}{2} (z^\alpha + (1 - z)^\alpha) \quad \text{for} \quad \alpha > -1. \quad (15)$$

The parameter  $\gamma = 1 + \alpha > 0$  suits for (11) and so (7) is fulfilled. Uniform repartition ( $\kappa(x, y) = \frac{1}{y} \mathbb{1}_{0 \leq x \leq y}$ ) corresponds to  $\alpha = 0$  and is also included.

This last case of uniform repartition is useful because it provides us with explicit formulas for the eigenelements. For instance, we can consider the two following examples.

**First example :**  $\tau(x) = \tau_0$ ,  $\beta(x) = \beta_0 x$ .

In this case, widely used by [19], the eigenelements exist and we have

$$\begin{aligned} \lambda &= \sqrt{\beta_0 \tau_0}, \\ \mathcal{U}(x) &= 2\sqrt{\frac{\beta_0}{\tau_0}} \left( X + \frac{X^2}{2} \right) e^{-X - \frac{X^2}{2}}, \quad \text{with } X = \sqrt{\frac{\beta_0}{\tau_0}} x, \\ \phi(x) &= \frac{1}{2} (1 + X). \end{aligned}$$

**Second example :**  $\tau(x) = \tau_0 x$ .

For such  $\beta$  for which there exists eigenelements, we have

$$\lambda = \tau_0 \quad \text{and} \quad \phi(x) = \frac{x}{\int y \mathcal{U}(y)}.$$

For instance when  $\beta(x) = \beta_0 x^n$  with  $n \in \mathbb{N}^*$ , then the eigenelements exist and we can compute  $\mathcal{U}$  and  $\phi$  and we have the formulas in Table 1. In this table we can notice that  $\mathcal{U}(0) > 0$  but the boundary condition  $\tau \mathcal{U}(0) = 0$  is fulfilled.

$n = 1$	$\lambda = \tau_0$	$\mathcal{U}(x) = \frac{\beta_0}{\tau_0} e^{-\frac{\beta_0}{\tau_0} x}$	$\phi(x) = \frac{\beta_0}{\tau_0} x$
$n = 2$	$\lambda = \tau_0$	$\mathcal{U}(x) = \sqrt{\frac{2\beta_0}{\pi\tau_0}} e^{-\frac{1}{2} \frac{\beta_0}{\tau_0} x^2}$	$\phi(x) = \sqrt{\frac{\pi\beta_0}{2\tau_0}} x$
$n$	$\lambda = \tau_0$	$\mathcal{U}(x) = \left( \frac{n\beta_0}{\tau_0} \right)^{\frac{1}{n}} \frac{n}{\Gamma(\frac{1}{n})} e^{-\frac{1}{n} \frac{\beta_0}{\tau_0} x^n}$	$\phi(x) = \left( \frac{\tau_0}{n\beta_0} \right)^{\frac{1}{n}} \frac{\Gamma(\frac{1}{n})}{\Gamma(\frac{2}{n})} x$

Table 1: The example  $\tau(x) = \tau_0 x$ ,  $\beta(x) = \beta_0 x^n$  and uniform repartition  $\kappa(x, y) = \frac{1}{y} \mathbb{1}_{0 \leq x \leq y}$ . The table gives the eigenelements solution to (3).

Now we turn to non-existence cases. Let us consider constant fragmentation  $\beta(x) = \beta_0$  with an affine polymerization  $\tau(x) = \tau_0 + \tau_1 x$ , and any fragmentation kernel  $\kappa$  which satisfies to assumptions (5)-(6). We notice that (13) is not satisfied and look at two instructive cases.

**First case :**  $\tau_0 = 0$ .

In this case assumption (12) does not hold true. Assume that there exists  $\mathcal{U} \in L^1(\mathbb{R}^+)$  solution of (3) with the estimates of Theorem 1. Integrating the equation on  $\mathcal{U}$  we obtain that  $\lambda = \beta_0$ , but multiplying the equation by  $x$  before integration we have that  $\lambda = \tau_1$ . We conclude that eigenelements cannot exist if  $\tau_1 \neq \beta_0$ .

Moreover, if we take  $\kappa(x, y) = \frac{1}{y} \mathbb{1}_{0 \leq x \leq y}$ , then a formal computation shows that any solution to the first equation of (3) belongs to the plan  $Vect\{x^{-1}, x^{-\frac{2\beta_0}{\tau_1}}\}$ . So, even if  $\beta_0 = \tau_1$ , there does not exist an eigenvector in  $L^1$ .

**Second case :**  $\tau_0 > 0$ .

In this case (12) holds true but the same integrations than before lead to

$$\int x\mathcal{U}(x) dx = \frac{\tau_0}{\beta_0 - \tau_1}.$$

So there cannot exist any eigenvector  $\mathcal{U} \in L^1(x dx)$  for  $\tau_1 \geq \beta_0$ .

### 3 Proof of the main theorem

#### 3.1 A preliminary lemma

Before proving Theorem 1, we give a preliminary lemma, useful to prove uniqueness of the eigenfunctions.

**Lemma 1 (Positivity)** *Consider  $\mathcal{U}$  and  $\phi$  solutions to the eigenproblem (3).*

*We define  $m := \inf_{x,y} \{x : (x, y) \in \text{Supp } \beta(y)\kappa(x, y)\}$ . Then we have, under assumptions (5), (6), (9) and (10)*

$$\begin{aligned} \text{Supp } \mathcal{U} &= [m, \infty) \quad \text{and} \quad \tau\mathcal{U}(x) > 0 \quad \forall x > m, \\ \phi(x) &> 0 \quad \forall x > 0. \end{aligned}$$

*If additionally  $\frac{1}{\tau} \in L^1_0$ , then  $\phi(0) > 0$ .*

**Remark 2** *In case  $\text{Supp } \kappa = \{(x, y) / x \leq y\}$ , then  $m = 0$  and Lemma 1 and Theorem 1 can be proved without the connexity condition (10) on the support of  $\beta$ .*

**Proof.** Let  $x_0 > 0$ , we define  $F : x \mapsto \tau(x)\mathcal{U}(x)e^{\int_{x_0}^x \frac{\lambda+\beta(s)}{\tau(s)} ds}$ . We have that

$$F'(x) = 2e^{\int_{x_0}^x \frac{\lambda+\beta(s)}{\tau(s)} ds} \int \beta(y)\kappa(x, y)\mathcal{U}(y) dy \geq 0. \quad (16)$$

So, as soon as  $\tau\mathcal{U}(x)$  once becomes positive, it remains positive for larger  $x$ .

We define  $a := \inf\{x : \tau(x)\mathcal{U}(x) > 0\}$ . We first prove that  $a \leq \frac{b}{2}$ . For this we integrate the equation on  $[0, a]$  to obtain

$$\int_0^a \int_a^\infty \beta(y)\kappa(x, y)\mathcal{U}(y) dy dx = 0,$$

$$\int_a^\infty \beta(y)\mathcal{U}(y) \int_0^a \kappa(x,y) dx dy = 0.$$

Thus for almost every  $y \geq \max(a, b)$ ,  $\int_0^a \kappa(x, y) dx = 0$ . As a consequence we have

$$1 = \int \kappa(x, y) dx = \int_a^y \kappa(x, y) dx \leq \frac{1}{a} \int x\kappa(x, y) dx = \frac{y}{2a}$$

thanks to (5) and (6), and this is possible only if  $b \geq 2a$ .

Assume by contradiction that  $m < a$ , integrating (3) multiplied by  $\varphi$ , we have for all  $\varphi \in \mathcal{C}_c^\infty$  such that  $\text{Supp } \varphi \subset [0, a]$

$$\int \int \varphi(x)\beta(y)\kappa(x,y)\mathcal{U}(y) dy dx = 0. \quad (17)$$

By definition of  $m$  and using the fact that  $m < a$ , there exists  $(p, q) \in (m, a) \times (b, \infty)$  such that  $(p, q) \in \text{Supp } \beta(y)\kappa(x, y)$ . But we can choose  $\varphi$  positive such that  $\varphi(p)\mathcal{U}(q) > 0$  and this is a contradiction with (17). So we have  $m \geq a$ .

To conclude we notice that on  $[0, m]$ ,  $\mathcal{U}$  satisfies

$$\partial_x(\tau(x)\mathcal{U}(x)) + \lambda\mathcal{U}(x) = 0.$$

So, thanks to the condition  $\tau(0)\mathcal{U}(0) = 0$  and the assumption (9), we have  $\mathcal{U} \equiv 0$  on  $[0, m]$ , so  $m = a$  and the first statement is proved.

For  $\phi$ , we define  $G(x) := \phi(x)e^{-\int_{x_0}^x \frac{\lambda+\beta(s)}{\tau(s)} ds}$ . We have that

$$G'(x) = -2e^{-\int_{x_0}^x \frac{\lambda+\beta(s)}{\tau(s)} ds} \beta(x) \int_0^x \kappa(y, x)\phi(y) dy \leq 0, \quad (18)$$

so, as soon as  $\phi$  vanishes, it remains null. Therefore  $\phi$  is positive on an interval  $(0, x_1)$  with  $x_1 \in \mathbb{R}_+^* \cup \{+\infty\}$ . Assuming that  $x_1 < +\infty$  and using that  $x_1 > a = m$  because  $\int \phi(x)\mathcal{U}(x)dx = 1$ , we can find  $X \geq x_1$  such that

$$\int_{x_1}^X G'(x) dx = -2 \int_{x_1}^X \int_0^{x_1} e^{\int_{x_0}^x \frac{\lambda+\beta(s)}{\tau(s)} ds} \phi(y)\beta(x)\kappa(y, x) dy dx < 0.$$

This contradicts that  $\phi(x) = 0$  for  $x \geq x_1$ , and we have proved that  $\phi(x) > 0$  for  $x > 0$ .

If  $\frac{1}{\tau} \in L_0^1$ , we can take  $x_0 = 0$  in the definition of  $G$  and so  $\phi(0) > 0$  or  $\phi \equiv 0$ . The fact that  $\phi$  is positive ends the proof of the lemma.

□

### 3.2 Truncated problem

The proof of the theorem is based on uniform estimates on the solution to a truncated equation. Let  $\eta, \delta, R$  positive numbers and define

$$\tau_\eta(x) = \begin{cases} \eta & 0 \leq x \leq \eta \\ \tau(x) & x \geq \eta. \end{cases}$$

Then  $\tau_\eta$  is lower bounded on  $[0, R]$  thanks to (9) and we denote by  $\mu = \mu(\eta, R) := \inf_{[0, R]} \tau_\eta$ . The existence of eigenelements  $(\lambda_\eta^\delta, \mathcal{U}_\eta^\delta, \phi_\eta^\delta)$  for the following truncated problem when  $\delta R < \mu$  is standard (see Theorem 3 in the Appendix).

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x}(\tau_\eta(x)\mathcal{U}_\eta^\delta(x)) + (\beta(x) + \lambda_\eta^\delta)\mathcal{U}_\eta^\delta(x) = 2 \int_x^R \beta(y)\kappa(x, y)\mathcal{U}_\eta^\delta(y) dy, \quad 0 < x < R, \\ \tau_\eta\mathcal{U}_\eta^\delta(x=0) = \delta, \quad \mathcal{U}_\eta^\delta(x) > 0, \quad \int \mathcal{U}_\eta^\delta(x)dx = 1, \\ -\tau_\eta(x)\frac{\partial}{\partial x}\phi_\eta^\delta(x) + (\beta(x) + \lambda_\eta^\delta)\phi_\eta^\delta(x) - 2\beta(x) \int_0^x \kappa(y, x)\phi_\eta^\delta(y) dy = \delta\phi_\eta^\delta(0), \quad 0 < x < R, \\ \phi_\eta^\delta(R) = 0, \quad \phi_\eta^\delta(x) > 0, \quad \int \phi_\eta^\delta(x)\mathcal{U}_\eta^\delta(x)dx = 1. \end{array} \right. \quad (19)$$

The proof of the theorem 1 requires  $\lambda_\eta^\delta > 0$ . To enforce it, we take  $\delta R = \frac{\mu}{2}$  and we consider  $R$  large enough to satisfy the following lemma.

**Lemma 2** *Under assumptions (5), (8) and (13), there exists a  $R_0 > 0$  such that for all  $R > R_0$ , if we choose  $\delta = \frac{\mu}{2R}$ , then we have  $\lambda_\eta^\delta > 0$ .*

**Proof.** Assume by contradiction that  $R > 0$  and  $\lambda_\eta^\delta \leq 0$  with  $\delta = \frac{\mu}{2R}$ . Then, integrating between 0 and  $x > 0$ , we obtain

$$\begin{aligned} 0 &\geq \lambda \int_0^x \mathcal{U}(y) dy \\ &= \delta - \tau(x)\mathcal{U}(x) - \int_0^x \beta(y)\mathcal{U}(y) dy + 2 \int_0^x \int_z^R \beta(y)\kappa(z, y)\mathcal{U}(y) dy dz \\ &= \delta - \tau(x)\mathcal{U}(x) + \int_0^x \beta(y)\mathcal{U}(y) dy + 2 \int_x^R \left( \int_0^x \kappa(z, y) dz \right) \beta(y)\mathcal{U}(y) dy \\ &\geq \delta - \tau(x)\mathcal{U}(x) + \int_0^x \beta(y)\mathcal{U}(y) dy. \end{aligned}$$

Consequently

$$\tau(x)\mathcal{U}(x) \geq \delta + \int_0^x \frac{\beta(y)}{\tau(y)}\tau(y)\mathcal{U}(y) dy$$

and, thanks to Grönwall's lemma,

$$\tau(x)\mathcal{U}(x) \geq \delta e^{\int_0^x \frac{\beta(y)}{\tau(y)} dy}.$$

But assumption (13) ensures that for all  $n \geq 0$ , there is a  $A > 0$  such that

$$\frac{\beta(x)}{\tau(x)} \geq \frac{n}{x}, \quad \forall x \geq A$$

and thus we have

$$\tau(x)\mathcal{U}(x) \geq \delta x^n, \quad \forall x \geq A.$$

Thanks to (8) we can choose  $n$  and  $A$  such that  $x^{-n}\tau(x) \leq \frac{\mu}{4}$  for  $x \geq A$  and then we have

$$1 = \int_0^R \mathcal{U}(x) dx \geq \int_A^R \mathcal{U}(x) dx \geq \delta \int_A^R \frac{x^n}{\tau(x)} dx \geq \frac{2}{R}(R - A)$$

what is a contradiction as soon as  $R > 2A$ ; so Lemma 2 holds for  $R_0 = 2A$ .

□

### 3.3 Limit as $\delta \rightarrow 0$ for $\mathcal{U}_\eta^\delta$ and $\lambda_\eta^\delta$

Fix  $\eta$  and let  $\delta \rightarrow 0$  (then  $R \rightarrow \infty$  since  $\delta R = \frac{\mu}{2}$ ).

*First estimate:  $\lambda_\eta^\delta$  upper bound.* Integrating equation (19) between 0 and  $R$ , we find

$$\lambda_\eta^\delta \leq \delta + \int \beta(x)\mathcal{U}_\eta^\delta(x) dx,$$

then the idea is to prove a uniform estimate on  $\int \beta\mathcal{U}_\eta^\delta$ . For this we begin with bounding the higher moments  $\int x^\alpha \beta\mathcal{U}_\eta^\delta$  for  $\alpha \geq 2$ .

Let  $\alpha \geq 2$ , according to (7) we have

$$\int \frac{x^\alpha}{y^\alpha} \kappa(x, y) dx \leq \int \frac{x^2}{y^2} \kappa(x, y) dx \leq c < \frac{1}{2}.$$

Multiplying the equation on  $\mathcal{U}_\eta^\delta$  by  $x^\alpha$  and then integrating on  $[0, R]$ , we obtain for all  $A \geq \eta$

$$\begin{aligned} \int x^\alpha ((1 - 2c)\beta(x))\mathcal{U}_\eta^\delta(x) dx &\leq \alpha \int x^{\alpha-1} \tau_\eta(x)\mathcal{U}_\eta^\delta(x) dx \\ &= \alpha \int_{x \leq A} x^{\alpha-1} \tau_\eta(x)\mathcal{U}_\eta^\delta(x) dx + \alpha \int_{x \geq A} x^{\alpha-1} \tau(x)\mathcal{U}_\eta^\delta(x) dx \\ &\leq \alpha A^{\alpha-1} \sup_{(0, A)} \tau + \omega_{A, \alpha} \int x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx, \end{aligned}$$

where  $\omega_{A, \alpha}$  is a positive number chosen to have  $\alpha\tau(x) \leq \omega_{A, \alpha}x\beta(x)$ ,  $\forall x \geq A$ . Thanks to (7) and (13), we can choose  $A_\alpha$  large enough to have  $\omega_{A_\alpha, \alpha} < 1 - 2c$ . Thus we find

$$\forall \alpha \geq 2, \exists A_\alpha : \forall \eta, \delta > 0, \quad \int x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx \leq \frac{\alpha A_\alpha^{\alpha-1} \sup_{(0, A)} \tau}{1 - 2c - \omega_{A_\alpha, \alpha}} := B_\alpha. \quad (20)$$

The next step is to prove the same estimates for  $0 \leq \alpha < 2$  and for this we first give a bound on  $\tau_\eta\mathcal{U}_\eta^\delta$ . We fix  $\rho \in (0, 1/2)$  and define  $x_\eta > 0$  as the unique point such that  $\int_0^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} dy = \rho$ . It exists because  $\beta$  is nonnegative and locally integrable, and  $\tau_\eta$  is positive. Thanks to assumption (12), we know that  $x_\eta \xrightarrow{\eta \rightarrow 0} x_0$  where  $x_0 > 0$  satisfies  $\int_0^{x_0} \frac{\beta(y)}{\tau(y)} dy = \rho$ , so  $x_\eta$  is bounded by  $0 < \underline{x} \leq x_\eta \leq \bar{x}$ .

Then, integrating (19) between 0 and  $x \leq x_\eta$ , we find

$$\begin{aligned}
\tau_\eta(x)\mathcal{U}_\eta^\delta(x) &\leq \delta + 2 \int_0^x \int \beta(y)\mathcal{U}_\eta^\delta(y)\kappa(z,y) dy dz \\
&\leq \delta + 2 \int \beta(y)\mathcal{U}_\eta^\delta(y) dy \\
&= \delta + 2 \int_0^{x_\eta} \beta(y)\mathcal{U}_\eta^\delta(y) dy + 2 \int_{x_\eta}^\infty \beta(y)\mathcal{U}_\eta^\delta(y) dy \\
&\leq \delta + 2 \sup_{(0,x_\eta)} \{\tau_\eta\mathcal{U}_\eta^\delta\} \int_0^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} dy + \frac{2}{x_\eta^2} \int_0^\infty y^2 \beta(y)\mathcal{U}_\eta^\delta(y) dy \\
&\leq \delta + 2\rho \sup_{(0,x_\eta)} \{\tau_\eta\mathcal{U}_\eta^\delta\} + \frac{2}{x_\eta^2} B_2.
\end{aligned}$$

Consequently, if we consider  $\delta \leq 1$  for instance, we obtain

$$\sup_{x \in (0,x)} \tau_\eta(x)\mathcal{U}_\eta^\delta(x) \leq \frac{1 + 2B_2/x^2}{1 - 2\rho} := C \quad (21)$$

so  $\tau_\eta\mathcal{U}_\eta^\delta$  is uniformly bounded in a neighborhood of zero.

Now we can prove a bound  $B_\alpha$  for  $x^\alpha\beta\mathcal{U}_\eta^\delta$  in the case  $0 \leq \alpha < 2$ . Thanks to the estimates (20) and (21) we have

$$\begin{aligned}
\int x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx &= \int_0^{\bar{x}} x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx + \int_{\bar{x}}^R x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx \\
&\leq \bar{x}^\alpha \sup_{(0,\bar{x})} \{\tau_\eta\mathcal{U}_\eta^\delta\} \int_0^{\bar{x}} \frac{\beta(y)}{\tau_\eta(y)} dy + \bar{x}^{\alpha-2} \int_{\bar{x}}^R x^2 \beta(x)\mathcal{U}_\eta^\delta(x) dx \\
&\leq C\rho\bar{x}^\alpha + B_2\bar{x}^{\alpha-2} := B_\alpha.
\end{aligned} \quad (22)$$

Combining (20) and (22) we obtain

$$\forall \alpha \geq 0, \exists B_\alpha : \forall \eta, \delta > 0, \quad \int x^\alpha \beta(x)\mathcal{U}_\eta^\delta(x) dx \leq B_\alpha, \quad (23)$$

and finally we bound  $\lambda_\eta^\delta$

$$\lambda_\eta^\delta = \delta + \int \beta\mathcal{U}_\eta^\delta \leq \delta + B_0. \quad (24)$$

So the family  $\{\lambda_\eta^\delta\}_\delta$  belong to a compact interval and we can extract a converging subsequence  $\lambda_\eta^\delta \xrightarrow{\delta \rightarrow 0} \lambda_\eta$ .

*Second estimate :  $W^{1,1}$  bound for  $x^\alpha\tau_\eta\mathcal{U}_\eta^\delta$ ,  $\alpha \geq 0$ .* We use the estimate (23). First we give a  $L^\infty$  bound for  $\tau_\eta\mathcal{U}_\eta^\delta$  by integrating (19) between 0 and  $x$

$$\tau_\eta(x)\mathcal{U}_\eta^\delta(x) \leq \delta + 2 \int_0^R \beta(y)\mathcal{U}_\eta^\delta(y) dy \leq \delta + 2B_0 := D_0. \quad (25)$$

Then we bound  $x^\alpha \tau_\eta \mathcal{U}_\eta^\delta$  in  $L^1$  for  $\alpha > -1$ . Assumption (13) ensures that there exists  $X > 0$  such that  $\tau(x) \leq x\beta(x)$ ,  $\forall x \geq X$ , so we have for  $R > X$

$$\begin{aligned} \int x^\alpha \tau_\eta(x) \mathcal{U}_\eta^\delta(x) dx &\leq \sup_{(0,X)} \{\tau_\eta \mathcal{U}_\eta^\delta\} \int_0^X x^\alpha dx + \int_X^R x^{\alpha+1} \beta(x) \mathcal{U}_\eta^\delta(x) dx \\ &\leq \sup_{(0,X)} \{\tau_\eta \mathcal{U}_\eta^\delta\} \frac{X^{\alpha+1}}{\alpha+1} + B_{\alpha+1} := C_\alpha. \end{aligned}$$

Finally

$$\forall \alpha > -1, \exists C_\alpha : \forall \eta, \delta > 0, \quad \int x^\alpha \tau_\eta(x) \mathcal{U}_\eta^\delta(x) dx \leq C_\alpha \quad (26)$$

and we also have that  $x^\alpha \mathcal{U}_\eta^\delta$  is bounded in  $L^1$  because  $\tau \in \mathcal{P}$  (see assumption (8)).

A consequence of (23) and (26) is that  $x^\alpha \tau_\eta \mathcal{U}_\eta^\delta$  is bound in  $L^\infty$  for all  $\alpha \geq 0$ . We already have (25) and for  $\alpha > 0$ , we multiply (19) by  $x^\alpha$ , integrate on  $[0, x]$  and obtain

$$x^\alpha \tau_\eta(x) \mathcal{U}_\eta^\delta(x) \leq \alpha \int_0^R y^{\alpha-1} \tau_\eta(y) \mathcal{U}_\eta^\delta(y) dy + 2 \int_0^R y^\alpha \beta(y) \mathcal{U}_\eta^\delta(y) dy \leq \alpha C_\alpha + 2B_\alpha := D_\alpha,$$

that give immediately

$$\forall \alpha \geq 0, \exists D_\alpha : \forall \eta, \delta > 0, \quad \sup_{x>0} x^\alpha \tau_\eta(x) \mathcal{U}_\eta^\delta(x) \leq D_\alpha. \quad (27)$$

To conclude we use the fact that neither the parameters nor  $\mathcal{U}_\eta^\delta$  are negative and we find by the chain rule, for  $\alpha \geq 0$

$$\begin{aligned} \int \left| \frac{\partial}{\partial x} (x^\alpha \tau_\eta(x) \mathcal{U}_\eta^\delta(x)) \right| dx &\leq \alpha \int x^{\alpha-1} \tau_\eta(x) \mathcal{U}_\eta^\delta(x) dx + \int x^\alpha |\partial_x (\tau_\eta(x) \mathcal{U}_\eta^\delta(x))| dx \\ &\leq \alpha \int x^{\alpha-1} \tau_\eta(x) \mathcal{U}_\eta^\delta(x) dx + \lambda_\eta^\delta \int x^\alpha \mathcal{U}_\eta^\delta(x) dx + 3 \int x^\alpha \beta(x) \mathcal{U}_\eta^\delta(x) dx \end{aligned} \quad (28)$$

and all the terms in the right hand side are uniformly bounded thanks to the previous estimates.

Since we have proved that the family  $\{x^\alpha \tau_\eta \mathcal{U}_\eta^\delta\}_\delta$  is bounded in  $W^{1,1}(\mathbb{R}^+)$  for all  $\alpha \geq 0$ , then, because  $\tau_\eta$  is positive and belongs to  $\mathcal{P}$ , we can extract from  $\{\mathcal{U}_\eta^\delta\}_\delta$  a subsequence which converges in  $L^1(\mathbb{R}^+)$  when  $\delta \rightarrow 0$ . Passing to the limit in equation (19) we find that

$$\begin{cases} \frac{\partial}{\partial x} (\tau_\eta(x) \mathcal{U}_\eta(x)) + (\beta(x) + \lambda_\eta) \mathcal{U}_\eta(x) = 2 \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_\eta(y) dy, \\ \mathcal{U}_\eta(0) = 0, \quad \mathcal{U}_\eta(x) \geq 0, \quad \int \mathcal{U}_\eta = 1, \end{cases} \quad (29)$$

with  $\lambda_\eta \geq 0$ .

### 3.4 Limit as $\eta \rightarrow 0$ for $\mathcal{U}_\eta$ and $\lambda_\eta$

All the estimates (20)-(28) remain true for  $\delta = 0$ . So we still know that the family  $\{x^\alpha \tau_\eta \mathcal{U}_\eta\}_\eta$  belongs to a compact set of  $L^1$ , but not necessarily  $\{\mathcal{U}_\eta\}_\eta$  because in the limit  $\tau$  can vanish at zero. We need one more estimate to study the limit  $\eta \rightarrow 0$ .

*Third estimate:  $L^\infty$  bound for  $x^\alpha \tau_\eta \mathcal{U}_\eta$ ,  $\alpha \geq -\gamma$ .* We already know that  $x^\alpha \tau_\eta \mathcal{U}_\eta$  is bounded for  $\alpha \geq 0$ . So, to prove the bound, it only remains to prove that  $x^{-\gamma} \tau_\eta \mathcal{U}_\eta$  is bounded in a neighborhood of zero. Let define  $f_\eta : x \mapsto \sup_{(0,x)} \tau_\eta \mathcal{U}_\eta$ . If we integrate (29) between 0 and  $x' < x$ , we find

$$\tau_\eta(x') \mathcal{U}_\eta(x') \leq 2 \int_0^{x'} \int \beta(y) \mathcal{U}_\eta(y) \kappa(z, y) dy dz \leq 2 \int_0^x \int \beta(y) \mathcal{U}_\eta(y) \kappa(z, y) dy dz$$

and so for all  $x$

$$f_\eta(x) \leq 2 \int_0^x \int \beta(y) \mathcal{U}_\eta(y) \kappa(z, y) dy dz.$$

We consider  $x_\eta$  and  $\underline{x}$  defined in the first estimate and, using (11) and (12), we have for all  $x < x_\eta$

$$\begin{aligned} f_\eta(x) &\leq 2 \int_0^x \int \beta(y) \mathcal{U}_\eta(y) \kappa(z, y) dy dz \\ &= 2 \int \beta(y) \mathcal{U}_\eta(y) \int_0^x \kappa(z, y) dz dy \\ &\leq 2 \int_0^\infty \beta(y) \mathcal{U}_\eta(y) \min\left(1, C \left(\frac{x}{y}\right)^\gamma\right) dy \\ &= 2 \int_0^x \beta(y) \mathcal{U}_\eta(y) dy + 2C \int_x^{x_\eta} \beta(y) \mathcal{U}_\eta(y) \left(\frac{x}{y}\right)^\gamma dy + 2C \int_{x_\eta}^\infty \beta(y) \mathcal{U}_\eta(y) \left(\frac{x}{y}\right)^\gamma dy \\ &= 2 \int_0^x \frac{\beta(y)}{\tau_\eta(y)} \tau_\eta(y) \mathcal{U}_\eta(y) dy + 2Cx^\gamma \int_x^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} \frac{\tau_\eta(y) \mathcal{U}_\eta(y)}{y^\gamma} dy + 2C \int_{x_\eta}^\infty \beta(y) \mathcal{U}_\eta(y) \left(\frac{x}{y}\right)^\gamma dy \\ &\leq 2f_\eta(x) \int_0^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} dy + 2Cx^\gamma \int_x^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} \frac{f_\eta(y)}{y^\gamma} dy + 2C \|\beta \mathcal{U}_\eta\|_{L^1} x^\gamma. \end{aligned}$$

We set  $\mathcal{V}_\eta(x) = x^{-\gamma} f_\eta(x)$  and we obtain

$$(1 - 2\rho) \mathcal{V}_\eta(x) \leq K + 2C \int_x^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} \mathcal{V}_\eta(y) dy.$$

Hence, using Grönwall's lemma, we find that  $\mathcal{V}_\eta(x) \leq \frac{K e^{\frac{2C\rho}{1-2\rho}}}{1-2\rho}$  and consequently

$$x^{-\gamma} \tau_\eta(x) \mathcal{U}_\eta(x) \leq \frac{K e^{\frac{2C\rho}{1-2\rho}}}{1-2\rho} := \tilde{C}, \quad \forall x \in [0, \underline{x}]. \quad (30)$$

This last estimate allows us to bound  $\mathcal{U}_\eta$  by  $\frac{x^\gamma}{\tau}$  which is in  $L^1_0$  by the assumption (11). Thanks to the second estimate, we also have that  $\int x^\alpha \mathcal{U}_\eta$  is bounded in  $L^1$  and so, thanks to the Dunford-Pettis theorem (see [8] for instance),  $\{\mathcal{U}_\eta\}_\eta$  belong to a  $L^1$ -weak compact set. Thus we can extract a subsequence which converges  $L^1$ -weak toward  $\mathcal{U}$ . But for all  $\varepsilon > 0$ ,  $\{x^\alpha \mathcal{U}_\eta\}_\eta$  is bounded in  $W^{1,1}([\varepsilon, \infty))$  for all  $\alpha \geq 1$  thanks to (28) and so the convergence is strong on  $[\varepsilon, \infty)$ . Then we write

$$\begin{aligned} \int |\mathcal{U}_\eta - \mathcal{U}| &= \int_0^\varepsilon |\mathcal{U}_\eta - \mathcal{U}| + \int_\varepsilon^\infty |\mathcal{U}_\eta - \mathcal{U}| \\ &\leq 2\tilde{C} \int_0^\varepsilon \frac{x^\gamma}{\tau(x)} + \int_\varepsilon^\infty |\mathcal{U}_\eta - \mathcal{U}|. \end{aligned}$$

The first term on the right hand side is small for  $\varepsilon$  small because  $\frac{x^\gamma}{\tau} \in L_0^1$  and then the second term is small for  $\eta$  small because of the strong convergence. Finally  $\mathcal{U}_\eta \xrightarrow{\eta \rightarrow 0} \mathcal{U}$  strongly in  $L^1(\mathbb{R}^+)$  and  $\mathcal{U}$  solution of the eigenproblem (3).

### 3.5 Limit as $\delta, \eta \rightarrow 0$ for $\phi_\eta^\delta$

We prove uniform estimates on  $\phi_\eta^\delta$  which are enough to pass to the limit and prove the result.

*Fourth estimate : uniform  $\phi_\eta^\delta$ -bound on  $[0, A]$ .* Let  $A > 0$ , our first goal is to prove the existence of a constant  $C_0(A)$  such that

$$\forall \eta, \delta, \quad \sup_{(0, A)} \phi_\eta^\delta \leq C_0(A).$$

We divide the equation on  $\phi_\eta^\delta$  by  $\tau_\eta$  and we integrate between  $x$  and  $x_\eta$  with  $0 < x < x_\eta$ , where  $x_\eta$ , bounded by  $\underline{x}$  and  $\bar{x}$ , is defined in the first estimate. Considering  $\delta < \frac{\mu(1-2\rho)}{\bar{x}}$  (fulfilled for  $R > \frac{\bar{x}}{2(1-2\rho)}$  since  $\delta = \frac{\mu}{2R}$ ), we find

$$\begin{aligned} \phi_\eta^\delta(x) &\leq \phi_\eta^\delta(x_\eta) + 2 \int_x^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} \int_0^y \kappa(z, y) \phi_\eta^\delta(z) dz + x_\eta \frac{\delta}{\mu} \phi_\eta^\delta(0) \\ &\leq \phi_\eta^\delta(x_\eta) + \sup_{(0, x_\eta)} \{\phi_\eta^\delta\} \left( 2 \int_0^{x_\eta} \frac{\beta(y)}{\tau_\eta(y)} \int_0^y \kappa(z, y) dz + x_\eta \frac{\delta}{\mu} \right) \end{aligned}$$

and we obtain

$$\sup_{x \in (0, \underline{x})} \phi_\eta^\delta(x) \leq \frac{1}{1 - 2\rho - \delta \bar{x} / \mu} \phi_\eta^\delta(x_\eta).$$

Using the decay of  $\phi_\eta^\delta(x) e^{-\int_x^{x_\eta} \frac{\beta + \lambda_\eta^\delta}{\tau_\eta}}$ , there exists  $C(A)$  such that

$$\sup_{x \in (0, A)} \phi_\eta^\delta(x) \leq C(A) \phi_\eta^\delta(x_\eta).$$

Noticing that  $\int \phi_\eta^\delta(x) \mathcal{U}_\eta^\delta(x) dx = 1$ , we conclude

$$1 \geq \int_0^{x_\eta} \phi_\eta^\delta(x) \mathcal{U}_\eta^\delta(x) dx \geq \phi_\eta^\delta(x_\eta) \int_0^{x_\eta} e^{-\int_x^{x_\eta} \frac{\beta + \lambda_\eta^\delta}{\tau_\eta}} \mathcal{U}_\eta^\delta(x) dx,$$

so, as  $x_\eta \rightarrow x_0$  and  $\int_0^{x_0} \mathcal{U}(x) dx > 0$  (thanks to Lemma 1 and because  $x_0 > b \geq a$ ), we have

$$\sup_{(0, A)} \phi_\eta^\delta \leq C_0(A). \tag{31}$$

*Fifth estimate : uniform  $\phi_\eta^\delta$ -bound on  $[A, \infty)$ .* Following an idea introduced in [29] we notice that the equation in (19) satisfied by  $\phi_\eta^\delta$  is a transport equation and therefore satisfies the maximum principle (see Lemma 4 in the Appendix). Therefore it remains to build a supersolution  $\bar{\phi}$  that is positive at  $x = R$ , to conclude  $\phi_\eta^\delta(x) \leq \bar{\phi}(x)$  on  $[0, R]$ .

This we cannot do on  $[0, R]$ , but on a subinterval  $[A_0, R]$  only. So we begin with an auxiliary function  $\bar{\varphi}(x) = x^k + \theta$  with  $k$  and  $\theta$  positive numbers to be determined. We have to check that on  $[A_0, R]$

$$-\tau(x) \frac{\partial}{\partial x} \bar{\varphi}(x) + (\lambda_\eta^\delta + \beta(x)) \bar{\varphi}(x) \geq 2\beta(x) \int \kappa(y, x) \bar{\varphi}(y) dy + \delta \phi_\eta^\delta(0),$$

*i.e.*

$$-k\tau(x)x^{k-1} + (\lambda_\eta^\delta + \beta(x)) \bar{\varphi}(x) \geq \left(2\theta + 2 \int \kappa(y, x) y^k dy\right) \beta(x) + \delta \phi_\eta^\delta(0).$$

For  $k \geq 2$ , we know that  $\int \kappa(y, x) \frac{y^k}{x^k} dy \leq c < 1/2$  so it is sufficient to prove that there exists  $A_0 > 0$  such that we have

$$-k\tau(x)x^{k-1} + (\lambda_\eta^\delta + \beta(x))(x^k + \theta) \geq (2\theta + 2cx^k)\beta(x) + \delta C_0(1) \quad (32)$$

for all  $x > A_0$ , where  $C_0$  is defined in (31). For this, dividing (32) by  $x^{k-1}\tau(x)$ , we say that if we have

$$(1 - 2c) \frac{x\beta(x)}{\tau(x)} \geq k + \frac{2\theta\beta(x) + \delta C_0(1)}{x^{k-1}\tau(x)}, \quad (33)$$

then (32) holds true. Thanks to assumptions (8) and (13) we know that there exists  $k > 0$  such that for any  $\theta > 0$ , there exists  $A_0 > 0$  for which (33) is true on  $[A_0, +\infty)$ .

Then we conclude by choosing the supersolution  $\bar{\phi}(x) = \frac{C_0(A_0)}{\theta} \bar{\varphi}(x)$  so that

$$\bar{\phi}(x) \geq \phi_\eta^\delta(x) \quad \text{on } [0, A_0],$$

and on  $[A_0, R]$ , we have

$$\begin{cases} -\tau(x) \frac{\partial}{\partial x} \bar{\phi}(x) + (\lambda_\eta^\delta + \beta(x)) \bar{\phi}(x) \geq 2\beta(x) \int_0^x \kappa(y, x) \bar{\phi}(y) dy + \delta \phi_\eta^\delta(0), \\ \bar{\phi}(R) > 0, \end{cases} \quad (34)$$

which is a supersolution to the equation satisfied by  $\phi_\eta^\delta$ . Therefore  $\phi_\eta^\delta \leq \bar{\phi}$  uniformly in  $\eta$  and  $\delta$  and we get

$$\exists k, \theta, C \text{ s.t. } \forall \eta, \delta, \quad \phi_\eta^\delta(x) \leq (Cx^k + \theta). \quad (35)$$

Equation (19) and the fact that  $\phi_\eta^\delta$  is uniformly bounded in  $L_{loc}^\infty(\mathbb{R}^+)$  give immediately that  $\partial_x \phi_\eta^\delta$  is uniformly bounded in  $L_{loc}^\infty(\mathbb{R}^+, \tau(x)dx)$ , so in  $L_{loc}^\infty(0, \infty)$  thanks to (9).

Then we can extract a subsequence of  $\{\phi_\eta^\delta\}$  which converges  $C^0(0, \infty)$  toward  $\phi$ . Now we check that  $\phi$  satisfied the adjoint equation of (3). We consider the terms of (19) one after another.

First  $(\lambda_\eta^\delta + \beta(x))\phi_\eta^\delta(x)$  converges to  $(\lambda + \beta(x))\phi(x)$  in  $L_{loc}^\infty$ .

For  $\partial_x \phi_\eta^\delta$ , we have an  $L^\infty$  bound on each compact of  $(0, \infty)$ . So it converges  $L^\infty$  - *weak* toward  $\partial_x \phi$ .

It remains the last term which we write, for all  $x > 0$ ,

$$\int_0^x \kappa(y, x) (\phi_\eta^\delta(y) - \phi(y)) dy \leq \|\phi_\eta^\delta - \phi\|_{L^\infty(0, x)} \xrightarrow{\eta, \delta \rightarrow 0} 0.$$

The fact that  $\int \phi \mathcal{U} = 1$  comes from the convergence  $L^\infty - L^1$  when written as

$$1 = \int \phi_\eta^\delta(x) \mathcal{U}_\eta^\delta(x) dx = \int \frac{\phi_\eta^\delta(x)}{1+x^k} (1+x^k) \mathcal{U}_\eta^\delta(x) dx \longrightarrow \int \frac{\phi(x)}{1+x^k} (1+x^k) \mathcal{U}(x) dx = \int \phi \mathcal{U}.$$

At this stage we have found  $(\lambda, \mathcal{U}, \phi) \in \mathbb{R}^+ \times L^1(\mathbb{R}^+) \times \mathcal{C}(\mathbb{R}^+)$  solution of (3). The estimates announced in Theorem 1 also follow from those uniform estimates. It remains to prove that  $\lambda > 0$  and the uniqueness.

### 3.6 Proof of $\lambda > 0$

We prove a little bit more, namely that

$$\lambda \geq \frac{1}{2} \sup_{x \geq 0} \{\tau(x) \mathcal{U}(x)\}. \quad (36)$$

We integrate the first equation of (3) between 0 and  $x$  and find

$$\begin{aligned} 0 \leq \lambda \int_0^x \mathcal{U}(y) dy &= -\tau(x) \mathcal{U}(x) - \int_0^x \beta(y) \mathcal{U}(y) dy + 2 \int_0^x \int_z^\infty \beta(y) \kappa(z, y) \mathcal{U}(y) dy dz \\ &\leq -\tau(x) \mathcal{U}(x) + 2 \int_0^\infty \int_z^\infty \beta(y) \kappa(z, y) \mathcal{U}(y) dy dz \\ &= -\tau(x) \mathcal{U}(x) + 2 \int_0^\infty \beta(y) \mathcal{U}(y) dy \\ &= -\tau(x) \mathcal{U}(x) + 2\lambda, \end{aligned}$$

Hence  $2\lambda \geq \tau(x) \mathcal{U}(x)$  and (36) is proved.

### 3.7 Uniqueness

We follow the idea of [23]. Let  $(\lambda_1, \mathcal{U}_1, \phi_1)$  and  $(\lambda_2, \mathcal{U}_2, \phi_2)$  two solutions to the eigenproblem (3). First we have

$$\begin{aligned} \lambda_1 \int \mathcal{U}_1(x) \phi_2(x) dx &= \int \left( -\partial_x(\tau(x) \mathcal{U}_1(x)) - \beta(x) \mathcal{U}_1(x) + 2 \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_1(y) dy \right) \phi_2(x) dx \\ &= \int \left( \tau(x) \partial_x \phi_2(x) - \beta(x) \phi_2(x) + 2\beta(x) \int_0^x \kappa(y, x) \phi_2(y) dy \right) \mathcal{U}_1(x) dx \\ &= \lambda_2 \int \mathcal{U}_1(x) \phi_2(x) dx \end{aligned}$$

and then  $\lambda_1 = \lambda_2 = \lambda$  because  $\int \mathcal{U}_1 \phi_2 > 0$  thanks to Lemma 1.

For the eigenvectors we use the General Relative Entropy method introduced in [25, 26]. For  $C > 0$ , we test the equation on  $\mathcal{U}_1$  against  $\text{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2} - C\right) \phi_1$ ,

$$0 = \int \left[ \partial_x(\tau(x) \mathcal{U}_1(x)) + (\lambda + \beta(x)) \mathcal{U}_1(x) - 2 \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_1(y) dy \right] \text{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx.$$

Deriving  $\left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \tau(x) \mathcal{U}_2(x) \phi_1(x)$  we find

$$\begin{aligned} \int \partial_x(\tau(x) \mathcal{U}_1(x)) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx &= \int \partial_x\left(\left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \tau(x) \mathcal{U}_2(x) \phi_1(x)\right) dx \\ &+ \int \partial_x(\tau(x) \mathcal{U}_2(x)) \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx - \int \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \partial_x(\tau(x) \mathcal{U}_2(x) \phi_1(x)) dx \end{aligned}$$

and then

$$\begin{aligned} \int \partial_x(\tau(x) \mathcal{U}_1(x)) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx &= \\ 2 \int \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \left[ \int_0^x \beta(y) \kappa(y, x) \mathcal{U}_2(y) \phi_1(y) dy - \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_2(y) \phi_1(y) dy \right] dx \\ &+ 2 \int \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_2(y) dy \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx \\ &- \int (\lambda + \beta(x)) \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \mathcal{U}_2(x) \phi_1(x) dx, \end{aligned}$$

$$\begin{aligned} \int \partial_x(\tau(x) \mathcal{U}_1(x)) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx &= \\ 2 \int \int \beta(y) \kappa(x, y) \left[ \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - C \right| - \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \right] \mathcal{U}_2(y) \phi_1(x) dx dy \\ &+ 2 \int \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_2(y) dy \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx \\ &- \int (\lambda + \beta(x)) \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \mathcal{U}_2(x) \phi_1(x) dx. \end{aligned}$$

So

$$\begin{aligned} 0 &= 2 \int \int \beta(y) \kappa(x, y) \left[ \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - C \right| - \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C \right| \right] \mathcal{U}_2(y) \phi_1(x) dx dy \\ &+ 2 \int \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_2(y) dy \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx \\ &- 2 \int \int_x^\infty \beta(y) \kappa(x, y) \mathcal{U}_1(y) dy \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \phi_1(x) dx \end{aligned}$$

$$0 = \int \int \beta(y) \kappa(x, y) \mathcal{U}_2(y) \left| \frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - C \right| \left[ 1 - \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - C\right) \right] \phi_1(x) dx dy.$$

Hence  $\left[1 - \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) - C\right) \operatorname{sgn}\left(\frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - C\right)\right] = 0$  on the support of  $\kappa(x, y)$  for all  $C$  thus  $\frac{\mathcal{U}_1}{\mathcal{U}_2}(x) = \frac{\mathcal{U}_1}{\mathcal{U}_2}(y)$  on the support of  $\kappa(x, y)$  and

$$\partial_x \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) = \int \beta(y) \kappa(x, y) \left( \frac{\mathcal{U}_1}{\mathcal{U}_2}(y) - \frac{\mathcal{U}_1}{\mathcal{U}_2}(x) \right) \frac{\mathcal{U}_2(y)}{\mathcal{U}_2(x)} dy = 0 \quad (37)$$

so  $\frac{\mathcal{U}_1}{\mathcal{U}_2} \equiv cst = 1$ .

We can prove in the same way that  $\phi_1 = \phi_2$  even if we can have  $\mathcal{U} \equiv 0$  on  $[0, m]$  with  $m > 0$ . Indeed in this case we know that  $\beta \equiv 0$  on  $[0, m]$  and so

$$\phi_i(x) = \phi_i(0) e^{\int_0^x \frac{\lambda}{\tau(s)} ds} \quad \forall x \in [0, m], \quad i \in \{1, 2\}.$$

## 4 Consequences, Perspectives

Existence of eigenelements is the basic stone for asymptotic studies when  $t \rightarrow \infty$ . The so-called "General Relative Entropy" method, introduced in [28, 25] and widely used (see [10, 9, 5] for instance) takes advantage of the eigenvalue problem and its adjoint to build entropy functionals. We recall for instance the following theorem (see [25, 26, 29]) for the linear case of Equation (1).

**Theorem 2** *Let  $u_0$  a function defined on  $\mathbb{R}^+$  satisfying*

$$\exists C_0 > 0, \quad s.t. \quad \forall x \geq 0 \quad |u_0(x)| \leq C_0 \mathcal{U}(x).$$

*Then there exists a unique solution  $u(x, t)$  to the equation (1) and we have for all  $t > 0$  the a priori bounds*

$$|u(x, t)| e^{-\lambda t} \leq C_0 \mathcal{U}(x),$$

$$\int u(y, t) e^{-\lambda t} \phi(y) dy = \int u_0(y) \phi(y) dy := \langle u_0, \phi \rangle,$$

*and the asymptotic behaviour*

$$\int_0^\infty |u(y, t) e^{-\lambda t} - \langle u_0, \phi \rangle \mathcal{U}(y)| \phi(y) dy \xrightarrow[t \rightarrow \infty]{} 0.$$

The eigenvalue problem can also be used in nonlinear cases, such as prion proliferation equations, where there is a quadratic coupling of Equation (1) with a differential equation. In [10, 9] for instance, the stability of steady states is investigated. The use of entropy methods in the case of nonlinear problems remains however a challenging and widely open field (see [30] for a recent review).

A following work is to study the dependency of the eigenvalue  $\lambda$  on parameters  $\tau$  and  $\beta$  (see [24]). For instance, our assumptions allow  $\tau$  to vanish at zero, what is a necessary condition to ensure that  $\lambda$  tends to zero when the fragmentation tends to infinity. Such results give precious information on the qualitative behaviour of the solution.

Another possible extension of the present work is to prove existence of eigenlements in the case of time-periodic parameters, using the Floquet's theory, and then compare the new  $\lambda_F$  with the time-independent one  $\lambda$  (see [11]). Such studies can help to choose a right strategy in order to optimize, for instance, the total mass  $\int xu(t, x)dx$  in the case of prion proliferation (see [10]) or on the contrary minimize the total population  $\int u(t, x)dx$  in the case of cancer therapy (see [12, 11]).

Finally, this eigenvalue problem could be used to recover some of the equation parameters like  $\tau$  and  $\beta$  from the knowledge of the asymptotic profile of the solution, as introduced in [31, 15] in the case of symmetric division ( $\tau = 1$  and  $\kappa = \delta_{x=\frac{y}{2}}$ ), by the use of inverse problems techniques. The method of [31] has to be adapted to our general case, in order to model prion proliferation for instance, or yet to recover the aggregation rate  $\tau$  ; this is another direction of future research.

### **Aknowledgment**

The authors thank a lot Benoît Perthame for his precious help and his corrections.

# Appendix

## A Assumption on $\kappa$ .

**Lemma 3** *Assumptions (5),(6) and (11) with  $\gamma > 0$  imply that*

$$\inf_y \lim_{\eta \rightarrow 0} \int_{\eta y}^{(1-\eta)y} \kappa(x, y) dx > 0,$$

*which means that polymers undergo a decrease in the size during fragmentation process. As a consequence, assumption (7) holds true.*

**Proof.** With the first assumption (5) we have

$$1 = \int_0^y \kappa(x, y) dx = \int_0^{\eta y} \kappa(x, y) dx + \int_{\eta y}^{(1-\eta)y} \kappa(x, y) dx + \int_{(1-\eta)y}^y \kappa(x, y) dx.$$

The two other assumptions (6) and (11) allow to control the mass of  $\kappa$  at the ends :

$$\begin{aligned} \int_{\eta y}^{(1-\eta)y} \kappa(x, y) dx &= 1 - \int_0^{\eta y} \kappa(x, y) dx - \int_{(1-\eta)y}^y \kappa(x, y) dx \\ &\geq 1 - C\eta^\gamma - \frac{1}{1-\eta} \int_{(1-\eta)y}^y \frac{x}{y} \kappa(x, y) dx \\ &\geq 1 - C\eta^\gamma - \frac{1}{2(1-\eta)} \xrightarrow{\eta \rightarrow 0} \frac{1}{2}, \end{aligned}$$

which gives the first assertion of the lemma.

Now we can prove (7) :

$$\begin{aligned} \int_0^y \frac{x^2}{y^2} \kappa(x, y) dx &\leq \left[ \int_0^{\eta y} \frac{x}{y} \kappa(x, y) dx + \int_{(1-\eta)y}^y \frac{x}{y} \kappa(x, y) dx \right] + \int_{\eta y}^{(1-\eta)y} \frac{x^2}{y^2} \kappa(x, y) dx \\ &\leq \left[ \frac{1}{2} - \int_{\eta y}^{(1-\eta)y} \frac{x}{y} \kappa(x, y) dx \right] + (1-\eta) \int_{\eta y}^{(1-\eta)y} \frac{x}{y} \kappa(x, y) dx \\ &= \frac{1}{2} - \eta \int_{\eta y}^{(1-\eta)y} \frac{x}{y} \kappa(x, y) dx \\ &\leq \frac{1}{2} - \eta^2 \int_{\eta y}^{(1-\eta)y} \kappa(x, y) dx. \end{aligned}$$

We use the first part of the proof to conclude. Taking  $\eta = \min\left(\frac{1}{4}, \frac{1}{(4C)^{1/\gamma}}\right)$  for instance, we obtain

$$\int_{\eta y}^{(1-\eta)y} \kappa(x, y) dx \geq \frac{1}{3},$$

and the lemma is proved for  $c = \frac{1}{2} - \frac{1}{48}$ .  $\square$

## B Krein-Rutman

We prove existence of solution for the truncated equation (19). In this part  $\eta$  and  $\delta$  are fixed (with  $\delta R < \mu$ ), so we will omit these indices for  $\tau$ ,  $\lambda$ ,  $\mathcal{U}$  and  $\phi$  but we keep in mind that  $\tau(x) \geq \mu > 0$ .

We use the Krein-Rutman theorem which requires working in the space of continuous functions (see [28] for instance). First we define regularized parameters as follows :

$$\tau_\varepsilon = \rho_\varepsilon * \tau, \quad \beta_\varepsilon = \rho_\varepsilon * \beta, \quad \text{and} \quad \forall y \geq 0, \quad \kappa_\varepsilon(\cdot, y) = \rho_\varepsilon * \kappa(\cdot, y),$$

where  $\rho_\varepsilon(x) = \frac{1}{\varepsilon} \rho(\frac{x}{\varepsilon})$  with  $\rho \in \mathcal{C}_c^\infty((0, \infty))$ , positive and such that  $\int_0^\infty \rho = 1$ . Then we have the theorem

**Theorem 3** *Under assumptions (5)-(13) on the parameters and for all  $\varepsilon > 0$ , there is a unique solution  $\lambda_\varepsilon \in \mathbb{R}$  and  $\mathcal{U}_\varepsilon, \phi_\varepsilon \in \mathcal{C}^1([0, R])$  to the regularized eigenproblem*

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x}(\tau_\varepsilon(x)\mathcal{U}_\varepsilon(x)) + (\beta_\varepsilon(x) + \lambda_\varepsilon)\mathcal{U}_\varepsilon(x) = 2 \int_0^R \beta_\varepsilon(y)\kappa_\varepsilon(x, y)\mathcal{U}_\varepsilon(y) dy, \quad 0 < x < R, \\ \tau_\varepsilon\mathcal{U}_\varepsilon(x=0) = \delta \int_0^R \mathcal{U}_\varepsilon(y) dy, \quad \mathcal{U}_\varepsilon(x) > 0, \quad \int_0^R \mathcal{U}_\varepsilon(x) dx = 1, \\ -\tau_\varepsilon(x)\frac{\partial}{\partial x}\phi_\varepsilon(x) + (\beta_\varepsilon(x) + \lambda_\varepsilon)\phi_\varepsilon(x) - 2\beta_\varepsilon(x) \int_0^R \kappa_\varepsilon(y, x)\phi_\varepsilon(y) dy = \tau_\varepsilon(0)\delta\phi_\varepsilon(0), \quad 0 < x < R, \\ \phi_\varepsilon(R) = 0, \quad \phi_\varepsilon(x) > 0, \quad \int_0^R \phi_\varepsilon(x)\mathcal{U}_\varepsilon(x) dx = 1. \end{array} \right. \quad (38)$$

**Proof.** We follow the proof of [28]. We define linear operators on  $E := \mathcal{C}^0([0, R])$  to apply the Krein-Rutman theorem.

**Direct equation.** For  $\nu > 0$  we consider the following equation on  $E$

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x}(n(x)) + \frac{\nu + \beta_\varepsilon(x)}{\tau_\varepsilon(x)}n(x) - 2 \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x, y)n(y) dy = \frac{f(x)}{\tau_\varepsilon(x)}, \quad 0 \leq x \leq R, \\ n(x=0) = \delta \int_0^R \frac{n(y)}{\tau_\varepsilon(y)} dy, \end{array} \right. \quad (39)$$

and we prove that the linear operator  $A : f \mapsto n$  (solution of (39)) satisfies to the assumptions of the Krein-Rutman theorem.

*First step: construction of A.* Fix  $f \in E$  and for  $m \in E$ , we define  $n = T(m) \in E$  as the (explicit) solution to

$$\left\{ \begin{array}{l} \frac{\partial}{\partial x}(n(x)) + \frac{\nu + \beta_\varepsilon(x)}{\tau_\varepsilon(x)}n(x) = 2 \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x, y)m(y) dy + \frac{f(x)}{\tau_\varepsilon(x)}, \quad 0 \leq x \leq R, \\ n(x=0) = \delta \int_0^R \frac{m(y)}{\tau_\varepsilon(y)} dy, \end{array} \right.$$

We prove that  $T$  is a strict contraction. Therefore it has a unique fixed point thanks to the Banach-Picard theorem. This fixed point is a solution to (39).

In order to prove that  $T$  is a strict contraction, we consider  $m_1$  and  $m_2$  two functions in  $E$ , we compute for  $n = n_1 - n_2$ ,  $m = m_1 - m_2$ ,

$$\begin{cases} \frac{\partial}{\partial x}(n(x)) + \frac{\nu + \beta_\varepsilon(x)}{\tau_\varepsilon(x)}n(x) = 2 \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x, y)m(y) dy, & 0 \leq x \leq R, \\ n(x=0) = \delta \int_0^R \frac{m(y)}{\tau_\varepsilon(y)} dy, \end{cases}$$

therefore

$$\begin{cases} \frac{\partial}{\partial x}|n(x)| + \frac{\nu + \beta_\varepsilon(x)}{\tau_\varepsilon(x)}|n(x)| \leq 2 \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x, y)|m(y)| dy, & 0 \leq x \leq R, \\ n(x=0) \leq \delta \int_0^R \frac{|m(y)|}{\tau_\varepsilon(y)} dy. \end{cases}$$

After integration, we obtain

$$|n(x)|e^{\int_0^x \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \leq \delta \int_0^R \frac{|m(y)|}{\tau_\varepsilon(y)} dy + \int_0^x e^{\int_0^{x'} \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x', y)|m(y)| dy dx'$$

and thus

$$\begin{aligned} |n(x)| &\leq \delta \int_0^R \frac{|m(y)|}{\tau_\varepsilon(y)} dy + \int_0^x e^{-\int_{x'}^x \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x', y)|m(y)| dy dx' \\ &\leq \|m\|_E \frac{1}{\mu} \left[ \delta R + \int_0^x e^{-\int_{x'}^x \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \int_0^R \beta_\varepsilon(y)\kappa_\varepsilon(x', y) dy dx' \right] \\ &\leq \|m\|_E \frac{1}{\mu} \left[ \delta R + \left\| \int_0^R \beta_\varepsilon(y)\kappa_\varepsilon(\cdot, y) dy \right\|_{L^\infty} \int_0^x e^{-\frac{\nu}{\|\tau_\varepsilon\|_{L^\infty}}(x-x')} dx' \right] \\ &\leq \|m\|_E \underbrace{\frac{1}{\mu} \left[ \delta R + \nu^{-1} \|\tau_\varepsilon\|_{L^\infty} \left\| \int_0^R \beta_\varepsilon(y)\kappa_\varepsilon(\cdot, y) dy \right\|_{L^\infty} \right]}_{:=k}. \end{aligned}$$

Because  $\delta R < \mu$  by assumption, we can choose  $\nu$  large so that  $k < 1$  and we obtain

$$\|n\|_E \leq k \|m\|_E.$$

Thus  $T$  is a strict contraction and we have proved the existence of a solution to (39).

*Second step: A is continuous.* This relies on a general argument which in fact shows that the linear mapping  $A$  is Lipschitz continuous. Indeed, arguing as above

$$|n(x)|e^{\int_0^x \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \leq \delta \int_0^R \frac{|n(y)|}{\tau_\varepsilon(y)} dy + \int_0^x e^{\int_0^{x'} \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \int_0^R \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)}\kappa_\varepsilon(x', y)|n(y)| dy dx' + \int_0^x e^{\int_0^{x'} \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \frac{|f(x')|}{\tau_\varepsilon(x')} dx',$$

and thus

$$|n(x)| \leq k \|n\|_E + \int_0^R \frac{|f(x')|}{\tau_\varepsilon(x')} dx' \leq k \|n\|_E + \frac{R}{\mu} \|f\|_E.$$

This indeed proves that

$$\|n\|_E \leq \frac{R}{\mu(1-k)} \|f\|_E.$$

*Third step: A is strongly positive.* For  $f \geq 0$ , the operator  $T$  of the first step maps  $m \geq 0$  to  $n \geq 0$ . Therefore the fixed point  $n$  is nonnegative. In other words  $n = A(f) \geq 0$ . If additionally  $f$  does not vanish, then  $n$  does not vanish either. Therefore  $n(0) = \delta \int_0^R \frac{n(y)}{\tau_\varepsilon(y)} dy > 0$  and thus

$$n(x) \geq n(0) + e^{-\int_0^x \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \int_0^x e^{-\int_0^{x'} \frac{\nu + \beta_\varepsilon}{\tau_\varepsilon}} \frac{f(x')}{\tau_\varepsilon(x')} dx' > 0.$$

*Fourth step: A is compact.* For  $\|f\|_E \leq 1$ , the third step proves that  $n$  is bounded in  $E$  and thus

$$\frac{\partial}{\partial x} n = -\frac{\nu + \beta_\varepsilon}{\tau_\varepsilon} n + \int \frac{\beta_\varepsilon(y)}{\tau_\varepsilon(y)} \kappa(x, y) n(y) dy + \frac{f}{\tau_\varepsilon}$$

is also bounded in  $E$ . Therefore by the Ascoli-Arzelà theorem the family  $n$  is relatively compact in  $E$ .

**Adjoint equation.** A function  $\phi$  is a solution to the adjoint equation of (38) if and only if  $\tilde{\phi}(x) := \phi(R - x)$  satisfies

$$\begin{cases} \tilde{\tau}_\varepsilon(x) \frac{\partial}{\partial x} \tilde{\phi}(x) + (\tilde{\beta}_\varepsilon(x) + \lambda_\varepsilon) \tilde{\phi}(x) - 2\tilde{\beta}_\varepsilon(x) \int_0^R \kappa_\varepsilon(y, R - x) \tilde{\phi}_\varepsilon(y) dy = \delta \tilde{\phi}_\varepsilon(R), & 0 < x < R, \\ \tilde{\phi}_\varepsilon(0) = 0, \end{cases} \quad (40)$$

where  $\tilde{\tau}_\varepsilon(x) = \tau_\varepsilon(R - x)$  and  $\tilde{\beta}_\varepsilon(x) = \beta_\varepsilon(R - x)$ . Then the same method than for the direct equation give the result, namely the existence of  $\lambda$  and  $\tilde{\phi}$  solution to (40).

Finally we have proved existence of  $(\lambda_{\mathcal{U}}, \mathcal{U}_\varepsilon)$  and  $(\lambda_\phi, \phi_\varepsilon)$  solution to the direct and adjoint equations of (38). It remains to prove that  $\lambda_{\mathcal{U}} = \lambda_\phi$  but it is nothing but integrating the direct equation against the adjoint eigenvector, what gives

$$\lambda_{\mathcal{U}} \int \mathcal{U}_\varepsilon \phi_\varepsilon = \lambda_\phi \int \mathcal{U}_\varepsilon \phi_\varepsilon.$$

□

To have existence of solution for (19), it remains to do  $\varepsilon \rightarrow 0$ . For this we can prove uniform bounds in  $L^\infty$  for  $\mathcal{U}_\varepsilon$  and  $\phi_\varepsilon$  because we are on the fixed compact  $[0, R]$ . Then we can extract subsequences which converge  $L^\infty$  \*-weak toward  $\mathcal{U}$  and  $\phi$ , solutions to (19) because  $\tau_\varepsilon$  and  $\beta_\varepsilon$  converge in  $L^1$  toward  $\tau$  and  $\beta$ . Concerning  $\kappa_\varepsilon$ , we have that for all  $\varphi \in C_c^\infty$ ,  $\int \varphi(x) \kappa_\varepsilon(x, y) dx \rightarrow \int \varphi(x) \kappa(x, y) dx \quad \forall y$ , and it is sufficient to pass to the limit in the equations.

## C Maximum principle

**Lemma 4** *If there exists  $A_0 > 0$  such that  $\bar{\phi} \geq \phi$  on  $[0, A_0]$  and  $\bar{\phi}$  a supersolution of (3) on  $[A_0, R]$  with  $\bar{\phi}(R) \geq \phi(R)$ , then  $\bar{\phi} \geq \phi$  on  $[0, R]$ .*

**Proof.** The proof is based on the same tools than to prove uniqueness (see above) or to establish GRE principles (see [25, 26] for instance).

We know that  $\bar{\phi} \geq \phi$  on  $[0, A_0]$  and that  $\bar{\phi}$  is a supersolution to the equation satisfied by  $\phi$  on  $[A_0, R]$ , *i.e.* there exists a function  $f \geq \delta\phi(0)$  such that

$$-\tau(x) \frac{\partial}{\partial x} \bar{\phi}(x) + (\lambda + \beta(x)) \bar{\phi}(x) = 2\beta(x) \int_0^x \kappa(y, x) \bar{\phi}(y) dy + f(x), \quad \forall x \in [A_0, R].$$

So we have for all  $x \in [A_0, R]$

$$-\tau(x) \frac{\partial}{\partial x} (\phi(x) - \bar{\phi}(x)) + (\lambda + \beta(x)) (\phi(x) - \bar{\phi}(x)) = 2\beta(x) \int_0^x \kappa(y, x) (\phi(y) - \bar{\phi}(y)) dy - f(x).$$

Then, multiplying by  $\mathbb{1}_{\phi \geq \bar{\phi}}$ , we obtain (see [28] for a justification)

$$-\tau(x) \frac{\partial}{\partial x} (\phi - \bar{\phi})_+(x) + (\lambda + \beta(x)) (\phi - \bar{\phi})_+(x) \leq 2\beta(x) \int_0^x \kappa(y, x) (\phi - \bar{\phi})_+(y) dy - f(x) \mathbb{1}_{\phi \geq \bar{\phi}}(x),$$

and this inequality is satisfied on  $[0, R]$  since  $(\phi - \bar{\phi})_+ \equiv 0$  on  $[0, A_0]$ .

If we test against  $\mathcal{U}$  we have, using the fact that  $\phi(R) = 0 < \bar{\phi}(R)$ ,

$$\begin{aligned} & \int_0^R (\phi - \bar{\phi})_+(x) \frac{\partial}{\partial x} (\tau(x) \mathcal{U}(x)) dx + \int_0^R (\lambda + \beta(x)) (\phi - \bar{\phi})_+(x) \mathcal{U}(x) dx \\ & \leq 2 \int_0^R (\phi - \bar{\phi})_+(y) \int_y^R \beta(x) \kappa(y, x) \mathcal{U}(x) dx dy - \int_0^R f(x) \mathbb{1}_{\phi \geq \bar{\phi}}(x) \mathcal{U}(x) dx. \end{aligned}$$

But if we test the equation (3) satisfied by  $\mathcal{U}$  against  $(\phi - \bar{\phi})_+$ , we find

$$\begin{aligned} & \int_0^R (\phi - \bar{\phi})_+(x) \frac{\partial}{\partial x} (\tau(x) \mathcal{U}(x)) dx + \int_0^R (\lambda + \beta(x)) (\phi - \bar{\phi})_+(x) \mathcal{U}(x) dx \\ & = 2 \int_0^R (\phi - \bar{\phi})_+(y) \int_y^R \beta(x) \kappa(y, x) \mathcal{U}(x) dx dy, \end{aligned}$$

and finally, subtracting,

$$0 \leq - \int_0^R f(x) \mathbb{1}_{\phi \geq \bar{\phi}}(x) \mathcal{U}(x) dx,$$

so

$$\delta\phi(0) \int_0^R \mathbb{1}_{\phi \geq \bar{\phi}}(x) \mathcal{U}(x) dx \leq 0$$

and this can hold only if  $\mathbb{1}_{\phi \geq \bar{\phi}} \equiv 0$  or  $\phi(0) = 0$ . But we deal with the truncated problem with  $\tau(x) \geq \eta > 0$ , so  $\frac{1}{\tau} \in L_0^1$  and  $\phi(0) > 0$  thanks to the lemma 1. Thus  $\mathbb{1}_{\phi \geq \bar{\phi}} \equiv 0$  and the lemma 4 is proved.  $\square$

## References

- [1] M. Adimy and L. Pujo-Menjouet. Asymptotic behaviour of a singular transport equation modelling cell division. *Dis. Cont. Dyn. Sys. Ser. B*, 3(3):439–456, 2003.
- [2] F. Baccelli, D. R. McDonald, and J. Reynier. A mean-field model for multiple tcp connections through a buffer implementing red. *Performance Evaluation*, 49(1-4):77 – 97, 2002.
- [3] J. Banasiak and W. Lamb. On a coagulation and fragmentation equation with mass loss. *Proc. Roy. Soc. Edinburgh Sect. A*, 136(6):1157–1173, 2006.
- [4] B. Basse, B.C. Baguley, E.S. Marshall, W.R. Joseph, B. van Brunt, G. Wake, and D. J. N. Wall. A mathematical model for analysis of the cell cycle in cell lines derived from human tumors. *J. Math. Biol.*, 47(4):295–312, 2003.
- [5] F. Bekkal Brikci, J. Clairambault, and B. Perthame. Analysis of a molecular structured population model with possible polynomial growth for the cell division cycle. *Math. Comput. Modelling*, 47(7-8):699–713, 2008.
- [6] F. Bekkal Brikci, J. Clairambault, B. Ribba, and B. Perthame. An age-and-cyclin-structured cell population model for healthy and tumoral tissues. *J. Math. Biol.*, 57(1):91–110, 2008.
- [7] T. Biben, J.-C. Geminard, and F. Melo. Dynamics of Bio-Polymeric Brushes Growing from a Cellular Membrane: Tentative Modelling of the Actin Turnover within an Adhesion Unit; the Podosome. *J. Biol. Phys.*, 31:87–120, 2005.
- [8] H. Brezis. *Analyse fonctionnelle*. Collection Mathématiques Appliquées pour la Maîtrise. [Collection of Applied Mathematics for the Master’s Degree]. Masson, Paris, 1983. Théorie et applications. [Theory and applications].
- [9] V. Calvez, N. Lenuzza, M. Doumic, J.-P. Deslys, F. Mouthon, and B. Perthame. Prion dynamic with size dependency - strain phenomena. *J. of Biol. Dyn.*, in press, 2008.
- [10] V. Calvez, V. Lenuzza, D. Oelz, J.-P. Deslys, P. Laurent, F. Mouthon, and B. Perthame. Size distribution dependence of prion aggregates infectivity. *Math. Biosci.*, 1:88–99, 2009.
- [11] J. Clairambault, S. Gaubert, and T. Lepoutre. Comparison of perron and floquet eigenvalues in age structured cell division cycle models. *Math. Model. Nat. Phenom.*, page in press, 2009.
- [12] J. Clairambault, S. Gaubert, and B. Perthame. An inequality for the Perron and Floquet eigenvalues of monotone differential systems and age structured equations. *C. R. Math. Acad. Sci. Paris*, 345(10):549–554, 2007.
- [13] O. Destaing, F. Saltel, J.-C. Geminard, P. Jurdic, and F. Bard. Podosomes Display Actin Turnover and Dynamic Self- Organization in Osteoclasts Expressing Actin-Green Fluorescent Protein. *Mol. Biol. of the Cell*, 14, 2003.
- [14] M. Doumic. Analysis of a population model structured by the cells molecular content. *Math. Model. Nat. Phenom.*, 2(3):121–152, 2007.
- [15] Marie Doumic, Benoît Perthame, and Jorge P. Zubelli. Numerical solution of an inverse problem in size-structured population dynamics. *Inverse Problems*, 25, 2009.

- [16] M. Escobedo, Ph. Laurençot, S. Mischler, and B. Perthame. Gelation and mass conservation in coagulation-fragmentation models. *J. Differential Equations*, 195(1):143–174, 2003.
- [17] M. Escobedo, S. Mischler, and B. Perthame. Gelation in coagulation and fragmentation models. *Comm. Math. Phys.*, 231(1):157–188, 2002.
- [18] J.Z. Farkas. Stability conditions for a nonlinear size-structured model. *Nonlin. Anal. Real World App.*, 6:962–969, 2005.
- [19] M.L. Greer, L. Pujó-Menjouet, and G.F. Webb. A mathematical analysis of the dynamics of prion proliferation. *J. Theoret. Biol.*, 242(3):598–606, 2006.
- [20] M. Gyllenberg and G.F. Webb. A Nonlinear Structured Population Model of Tumor Growth With Quiescence. *J. Math. Biol.*, 28:671–694, 1990.
- [21] P. Laurençot and C. Walker. Well-posedness for a model of prion proliferation dynamics. *J. Evol. Equ.*, 7(2):241–264, 2007.
- [22] J. A. J. Metz and O. Diekmann, editors. *The dynamics of physiologically structured populations*, volume 68 of *Lecture Notes in Biomathematics*. Springer-Verlag, Berlin, 1986. Papers from the colloquium held in Amsterdam, 1983.
- [23] P. Michel. Existence of a solution to the cell division eigenproblem. *Math. Models Methods Appl. Sci.*, 16(7, suppl.):1125–1153, 2006.
- [24] P. Michel. Optimal proliferation rate in a cell division model. *Math. Model. Nat. Phenom.*, 1(2):23–44, 2006.
- [25] P. Michel, S. Mischler, and B. Perthame. General entropy equations for structured population models and scattering. *C. R. Math. Acad. Sci. Paris*, 338(9):697–702, 2004.
- [26] P. Michel, S. Mischler, and B. Perthame. General relative entropy inequality: an illustration on growth models. *J. Math. Pures Appl. (9)*, 84(9):1235–1260, 2005.
- [27] K. Pakdaman, B. Perthame, and D. Salort. Dynamics of a structured neuron population. (submitted), 2009.
- [28] B. Perthame. *Transport equations in biology*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [29] B. Perthame and L. Ryzhik. Exponential decay for the fragmentation or cell-division equation. *J. Differential Equations*, 210(1):155–177, 2005.
- [30] B. Perthame and T.M. Touaoula. Analysis of a cell system with finite divisions. *Boletín SEMA*, (44), 2008.
- [31] Benoît Perthame and Jorge P. Zubelli. On the inverse problem for a size-structured population model. *Inverse Problems*, 23(3):1037–1052, 2007.
- [32] J.R. Silveira, G.J. Raymond, A.G. Hughson, R.E. Race, V.L. Sim, S.F. Hayes, and B. Caughey. The most infectious prion protein particles. *Nature*, 437(7056):257–261, September 2005.