

# FINITE-DIMENSIONAL APPROXIMATION OF THE SELF-DIFFUSION COEFFICIENT FOR THE EXCLUSION PROCESS

C. LANDIM, S. OLLA AND S. R. S. VARADHAN

IMPA and CNRS UPRES-A 6085  
Université de Cergy CNRS UPRES-A 8088  
Courant Institute, New York University

ABSTRACT. We show that for the symmetric simple exclusion process on  $\mathbb{Z}^d$  the selfdiffusion coefficient of a tagged particle is stable when approximated by simple exclusion processes on large periodic lattices. The proof depends on a similar stability property of the asymptotic variance of additive functionals of mean zero. This requires establishing a property for the Dirichlet space known as Liouville-D property.

## 1. INTRODUCTION

In the early 80's Kipnis and Varadhan proved an invariance principle for the position of a tagged particle in a symmetric simple exclusion process in equilibrium (cf. [KV]). Their proof relies on a general central limit theorem for additive functionals of a reversible Markov process that they establish. Time reversibility and translation invariance of the system are the basic ingredients of this method, and in principle it can be applied to any system with these two symmetries. Later it has been extended to non-reversible processes that satisfy a strong sector condition. (cf. [V1], [OS]).

The (effective) diffusion matrix of the limiting Brownian Motion is a function  $D(\alpha)$  of the density  $\alpha$  of the particles and in the reversible case it is given by a variational formula.

This method works directly in infinite systems and it raises the natural question of the stability of the effective diffusion matrix under finite dimensional approximations. More precisely, consider a finite dimensional version of the same process on

---

1991 *Mathematics Subject Classification.* 60K35.

*Key words and phrases.* selfdiffusion, tagged particle, exclusion process, Liouville property.

C. L. was partially supported by CNPq grant 300358/93-8, FAPERJ grant E26/150.940/99 and PRONEX grant 41.96.0923.00. S. R. S. Varadhan was supported by the National Science Foundation grant DMS-9803140.

the torus  $\{-N, \dots, 0, \dots, N\}^d$  (i.e. with periodic boundary conditions, preserving in this manner the translation symmetry). Since we want to work with an ergodic process, we also fix the total number  $K$  of particles. Consider now a tagged particle in this finite system. If  $N$  is much larger than the size of a single jump, the motion of the tagged particle has a unique canonical lifting to  $\mathbb{Z}^d$ . We get in this manner a process  $X_N(t)$  with values in  $\mathbb{Z}^d$ . Let us denote by  $D_{[N,K]}$  the variance of the Brownian motion which is the limit of the scaled process  $\varepsilon X_N(\varepsilon^{-2}t)$  as  $\varepsilon \rightarrow 0$ . We expect that

$$\lim_{\substack{N \rightarrow \infty \\ K/(2N)^d \rightarrow \alpha}} D_{[N,K]} = D(\alpha). \quad (1.1)$$

Related questions arise in other contexts, for example, the effective conductivity in percolation regime (cf. [K], [DFGW]) and fluctuations in massless interface models (cf. [GOS]).

In general it is easy to prove that

$$\limsup_{\substack{N \rightarrow \infty \\ K/(2N)^d \rightarrow \alpha}} D_{[N,K]} \leq D(\alpha).$$

In this paper we prove the equality (1.1) for symmetric simple exclusion processes on  $\mathbb{Z}^d$ .

A related problem that is somewhat easier to handle is to study the scaling behavior of an additive functional  $A(t) = \int_0^t V(\eta(s)) ds$  where  $V$  is a suitable local function that depends on the configuration at a few sites. There is again a limiting Brownian motion for  $\varepsilon A(\varepsilon^{-2}t)$  with variance  $\sigma_{N,K}^2$ . The problem is to show that

$$\lim_{\substack{N \rightarrow \infty \\ K/(2N)^d \rightarrow \alpha}} \sigma_{N,K}^2 = \sigma^2(\alpha), \quad (1.2)$$

where  $\sigma^2(\alpha)$  is the corresponding variance in the infinite volume situation where the central limit theorem can be established directly. In this context the nontrivial part is the inequality

$$\limsup_{\substack{N \rightarrow \infty \\ K/(2N)^d \rightarrow \alpha}} \sigma_{N,K}^2 \leq \sigma^2(\alpha)$$

The proofs depend on *approximating* the *solutions* of the infinite dimensional variational problem for the limiting variance by solutions of the corresponding finite dimensional problem.

A basic step in this approximation is to prove a certain regularity property of the Dirichlet form associated to the simple exclusion process. Let us describe this regularity property in a more general context:

Suppose we have a collection of invertible measure preserving transformations  $\{T_z\}$  on some probability space  $(\Omega, \mathcal{F}, \mu)$ , then we can define a Dirichlet form

$$D(u) = \frac{1}{2} \int_{\Omega} \sum_z [u(T_z \omega) - u(\omega)]^2 d\mu$$

corresponding to the generator

$$Lu = \sum_z [u(T_z \omega) - u(\omega)]$$

of a process consisting of random stirrings. If we define  $v_z(\omega) = (\nabla_z u)(\omega) = u(T_z \omega) - u(\omega)$ , there is natural map

$$u \rightarrow \delta u = \{v_z(\omega)\} \in \oplus_z L_2(\mu) .$$

The problem is to characterize the closure of the range of this map by relations that remain stable under suitable approximations done with finitely supported functions.

In section 3 we prove this regularity property for the symmetric simple exclusion processes and in section 4 for the version with a tagged particle. The main tool in the proof is duality, that allows us to reduce the problem to a finite dimensional situation (cf. [LY], [SVY]).

A similar question arises in the context of hydrodynamic limit for non-gradient models (cf. [V2], [KL] appendix 3). But there the approximation is needed in a different norm and the proof relies usually on a suitable estimate of the spectral gap for the corresponding dynamics in a finite volume.

## 2. NOTATION AND RESULTS

Let us fix a symmetric, finite-range, probability measure  $p(\cdot)$  on  $\mathbb{Z}^d$ :  $p(0) = 0$ ,  $p(z) = p(-z)$  and  $p(z) = 0$  for  $|z|$  large. Let us assume that the random walk with transition probability  $p(\cdot)$  is irreducible, i.e. the finite set  $\{z : p(z) > 0\}$  generates the group  $\mathbb{Z}^d$ . Consider the symmetric simple exclusion process associated to  $p(\cdot)$ . This is the Markov process on  $\mathcal{X} = \{0, 1\}^{\mathbb{Z}^d}$  whose generator  $L$  acting on cylinder functions  $f$  is given by

$$\begin{aligned} (Lf)(\eta) &= \sum_{x, y \in \mathbb{Z}^d} p(y-x) \eta(x) [1 - \eta(y)] [f(\sigma^{x, y} \eta) - f(\eta)] \\ &= \frac{1}{2} \sum_{x, y \in \mathbb{Z}^d} p(y-x) [f(\sigma^{x, y} \eta) - f(\eta)] . \end{aligned} \tag{2.1}$$

Here and below, a configuration of particles in  $\mathbb{Z}^d$ , i.e a point in  $\mathcal{X}$ , will be denoted by  $\eta$ . In particular, for  $x$  in  $\mathbb{Z}^d$ ,  $\eta(x)$  is equal to 1 if the site  $x$  is occupied and 0 otherwise. Moreover, for a configuration  $\eta$  and  $x, y$  in  $\mathbb{Z}^d$ ,  $\sigma^{x, y} \eta$  is the configuration obtained from  $\eta$  by exchanging the occupation variables  $\eta(x)$  and  $\eta(y)$  :

$$(\sigma^{x, y} \eta)(z) = \begin{cases} \eta(y) & \text{if } z = x, \\ \eta(x) & \text{if } z = y, \\ \eta(z) & \text{otherwise .} \end{cases} \tag{2.2}$$

Note that  $\sigma^{x,y} = \sigma^{y,x}$  and we can think of either one as being  $\sigma^e$  where  $e$  is an edge or a bond connecting  $x$  and  $y$ . Fix  $0 \leq \alpha \leq 1$  and denote by  $\nu_\alpha$  the Bernoulli product measure on  $\mathcal{X}$ . This is the probability measure on  $\mathcal{X}$  obtained by placing a particle with probability  $\alpha$  at each site  $x$ , and independently for different sites. It is easy to check that the one-parameter family of probability measures  $\{\nu_\alpha, 0 \leq \alpha \leq 1\}$  are reversible invariant measures for the process with generator  $L$  on  $\mathcal{X}$ . The Dirichlet form is given by

$$D(f) = \frac{1}{4} \int_{\mathcal{X}} \sum_{x,y \in \mathbb{Z}^d} p(y-x) [f(\sigma^{x,y}\eta) - f(\eta)]^2 d\nu_\alpha .$$

We will think of  $\mathbb{Z}^d$  as a connected graph with an edge  $e$  connecting  $x, y$  being present whenever  $p(y-x) > 0$ . Since  $p(\cdot)$  is symmetric, one can view  $p(y-x)$  as defined on edges with  $p(e) = p(y-x)$ . Denote by  $\mathbb{E}$  the set of such edges. For each local function  $f$ , denote by  $\delta f$  the gradient obtained from  $f$ . This is the collection  $\{(\nabla_e f)(\eta)\}$  of functions on  $\mathcal{X}$  defined for  $e = (x, y) \in \mathbb{E}$ , by

$$(\nabla_e f)(\eta) = f(\sigma^{x,y}\eta) - f(\eta) .$$

Since  $\sigma^{x,y} = \sigma^{y,x}$  we can define  $\sigma^e$  for  $e \in \mathbb{E}$ . If  $e_1, \dots, e_n$  are  $n$  edges in  $\mathbb{E}$  such that  $\sigma^{e_1} \dots \sigma^{e_n}$  is the identity permutation then

$$\begin{aligned} 0 &= f(\sigma^{e_1} \dots \sigma^{e_n} \eta) - f(\eta) \\ &= \sum_{j=1}^n [f(\sigma^{e_j} \dots \sigma^{e_n} \eta) - f(\sigma^{e_{j+1}} \dots \sigma^{e_n} \eta)] \\ &= \sum_{j=1}^n (\nabla_{e_j} f)(\sigma^{e_{j+1}} \dots \sigma^{e_n} \eta) \\ &= \sum_{j=1}^n \sigma^{e_{j+1}} \dots \sigma^{e_n} (\nabla_{e_j} f)(\eta) . \end{aligned}$$

It is understood here that for  $j = n$  the expression is just  $(\nabla_{e_n} f)(\eta)$ . The above conditions are called the compatibility conditions for a gradient and they permit us to recognize when an arbitrary collection  $\Phi = \{\Phi_e, e \in \mathbb{E}\}$  of  $L^2(\nu_\alpha)$  functions could be a gradient. We say that such a collection is a formal gradient if each  $\Phi_e$  is square integrable and whenever  $e_1, \dots, e_n$  is a finite set of edges in  $\mathbb{E}$  such that  $\sigma^{e_1} \dots \sigma^{e_n}$  is equal to the identity permutation, we have

$$\sum_{j=1}^n \sigma^{e_{j+1}} \dots \sigma^{e_n} \Phi_{e_j} = 0 . \quad (2.3)$$

As before, for  $j = n$  this expression is  $\Phi_{e_n}$ . We associate a Hilbert Space with norm defined by

$$\|\Phi\|_\alpha^2 = \sum_{e \in \mathbb{E}} p(e) \langle \Phi_e^2 \rangle_{\nu_\alpha} . \quad (2.4)$$

Any  $\Phi$  for which the norm is finite will be called a square integrable formal gradient. Although we have defined  $\nabla_e f$  only for edges  $e = (x, y)$  with  $p(y - x) > 0$ , if the random walk with transition probability  $p(\cdot)$  is irreducible then knowing  $\nabla_e$  for edges in  $\mathbb{E}$  determines it uniquely for all edges  $(x, y)$ . The Dirichlet forms corresponding to different irreducible finitely supported probabilities  $p(\cdot)$  are equivalent. For qualitative statements we could take as Dirichlet form simply

$$\frac{1}{4} \int \sum_{\substack{x, y \\ y-x \in F}} |f(\sigma^{x, y} \eta) - f(\eta)|^2 d\nu_\alpha ,$$

where  $F$  is any symmetric set that generates  $\mathbb{Z}^d$  as a group. The natural question in this context is whether any square integrable formal gradient can be approximated by actual gradients in the  $L^2$  sense, i.e. for each  $\Phi = \{\Phi_e\}$ , does there exist a sequence of local functions  $f_n$  such that

$$\lim_{n \rightarrow \infty} \|\Phi - \delta f_n\|_\alpha = 0 . \quad (2.5)$$

The next result states that this is indeed possible.

**Theorem 2.1 .** *For the symmetric exclusion process on  $\mathbb{Z}^d$ , any square integrable formal gradient can be approximated by gradients of local functions in the sense of (2.5).*

**Corollary 2.2 .** *For a symmetric simple exclusion process on  $\mathbb{Z}^d$  the limit (1.2) is valid for any local function  $V$  which has mean 0 with respect to every  $\nu_\alpha$ .*

In order to study the evolution of a tagged particle we must consider simple exclusion on the lattice  $\mathbb{Z}_*^d = \mathbb{Z}^d - \{0\}$ . We have analogous results in that case. Of course in dealing with simple exclusion on  $\mathbb{Z}_*^d$ ,  $\mathbb{E}$  contains only edges  $(x, y)$  such that  $y - x \in F$  and  $x, y \in \mathbb{Z}_*^d$  and the Dirichlet form is now given by

$$D(f) = \frac{1}{4} \int_{\mathcal{X}} \sum_{\substack{x, y \in \mathbb{Z}_*^d \\ y-x \in F}} p(y-x) [f(\sigma^{x, y} \eta) - f(\eta)]^2 d\nu_\alpha .$$

**Theorem 2.3 .** *For the symmetric exclusion process (other than the 1-dimensional nearest neighbor walk) on  $\mathbb{Z}_*^d$ , any square integrable formal gradient can be approximated by gradients of local functions in the sense of (2.5).*

**Corollary 2.4 .** *For a symmetric simple exclusion process (other than the 1-dimensional nearest neighbor walk) on  $\mathbb{Z}_*^d$  the limit (1.2) is valid for any local function  $V$  which has mean 0 with respect to every  $\nu_\alpha$ .*

We will apply Theorem 2.3 in order to study the evolution of a tagged particle in the symmetric simple exclusion process. Let  $\eta$  be an initial configuration with a particle at the origin:  $\eta(0) = 1$ . Tag this particle and denote by  $\eta_t$  and  $X_t$  the state of the process and the position of the tagged particle respectively, at time  $t$ .

We shall refer to  $\eta_t$  as the environment. Let  $\xi_t$  be the state of the environment as seen from the tagged particle:  $\xi_t = \tau_{X_t}\eta_t$ . Here, for  $x$  in  $\mathbb{Z}^d$  and a configuration  $\eta$ ,  $\tau_x$  stands for the translation of  $\eta$  by  $x$  units :  $(\tau_x\eta)(y) = \eta(x+y)$ . Notice that the origin is always occupied (by the tagged particle) for the environment as seen from the tagged particle. For this reason, we shall consider the process  $\xi_t$  as taking values in  $\{0, 1\}^{\mathbb{Z}_*^d}$ .

While  $X_t$  is not a Markov process due to the presence of the environment,  $(X_t, \eta_t)$  and  $\xi_t$  are Markov processes. A simple computation shows that the generator  $\mathcal{L}$  of the Markov process  $\xi_t$  is given by  $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_\tau$ , where

$$\begin{aligned} (\mathcal{L}_0 f)(\xi) &= \sum_{x, y \in \mathbb{Z}_*^d} p(y-x) \xi(x) [1 - \xi(y)] [f(\sigma^{x,y}\xi) - f(\xi)], \\ &= \frac{1}{2} \sum_{x, y \in \mathbb{Z}_*^d} p(y-x) [f(\sigma^{x,y}\xi) - f(\xi)], \\ (\mathcal{L}_\tau f)(\xi) &= \sum_{z \in \mathbb{Z}_*^d} p(z) [1 - \xi(z)] [f(\tau_z \xi) - f(\xi)]. \end{aligned}$$

The first part of the generator takes into account the jumps of the environment, while the second one corresponds to jumps of the tagged particle. In the above formula,  $\tau_z \xi$  stands for the configuration where the tagged particle, sitting at the origin, is first transferred to site  $z$  and then the entire configuration is translated by  $-z$ : i.e. for  $x \neq 0$ ,

$$(\tau_z \xi)(x) = \begin{cases} 0 & \text{if } x = -z, \\ \xi(x+z) & \text{for } x \neq -z. \end{cases}$$

For  $0 \leq \alpha \leq 1$ , denote by  $\mu_\alpha$  the Bernoulli product measure on  $\mathcal{X}_* = \{0, 1\}^{\mathbb{Z}_*^d}$ . A simple computation shows that  $\mu_\alpha$  is reversible for the Markov process  $\xi_t$ .

Kipnis and Varadhan [KV] proved a central limit theorem for the position of the tagged particle when the environment process is in equilibrium with marginal distribution  $\mu_\alpha$ . They showed that  $\varepsilon X_{t\varepsilon^{-2}}$  converges, as  $\varepsilon \downarrow 0$ , to a Brownian motion with self-diffusion coefficient  $D(\alpha)$  that can be described as follows. Fix  $\lambda$  in  $\mathbb{R}^d$ . Then the quadratic form  $\langle \lambda, D(\alpha)\lambda \rangle$  is given by the variational formula :

$$\begin{aligned} \langle \lambda, D(\alpha)\lambda \rangle &= \\ \inf_f \left\{ \sum_{z \in \mathbb{Z}_*^d} p(z) \langle [1 - \xi(z)] \{ \lambda \cdot z - T_z f \}^2 \rangle_\alpha + \sum_{x, y \in \mathbb{Z}_*^d} p(y-x) \langle \{ T_{x,y} f \}^2 \rangle_\alpha \right\}. \end{aligned} \tag{2.6}$$

In this formula the infimum is taken over all cylinder functions  $f$ ,  $\langle \cdot \rangle_\alpha$  stands for the expectation with respect to the measure  $\mu_\alpha$  and  $T_z, T_{x,y}$ ,  $x, y, z$  in  $\mathbb{Z}_*^d$ , are the operators defined by

$$\begin{aligned} (T_z f)(\xi) &= [1 - \xi(z)] \{ f(\tau_z \xi) - f(\xi) \}, \\ (T_{x,y} f)(\xi) &= \{ f(\sigma^{x,y}\xi) - f(\xi) \} \end{aligned}$$

and the second summation is performed over all bonds of  $\mathbb{Z}_*^d$ .

The diffusion coefficient  $D(\alpha)$  can also be written as

$$\langle \lambda \cdot D(\alpha) \lambda \rangle = (1 - \alpha) \sum_{z \in \mathbb{Z}_*^d} (z \cdot \lambda)^2 p(z) - 2 \|\psi_\lambda\|_{-1, \alpha}^2, \quad (2.7)$$

where  $\psi_\lambda$  is the cylinder function given by

$$\psi_\lambda(\xi) = \sum_{z \in \mathbb{Z}_*^d} p(z) (z \cdot \lambda) [1 - \xi(z)] \quad (2.8)$$

and  $\|\psi_\lambda\|_{-1, \alpha}^2$  is the  $H_{-1}$  norm of  $\psi_\lambda$ , which is given by

$$\|\psi\|_{-1, \alpha}^2 = \sup_f \left\{ 2 \langle f, \psi_\lambda \rangle_\alpha - \langle f, (-\mathcal{L})f \rangle_\alpha \right\}. \quad (2.9)$$

In this formula the supremum is carried over all cylinder functions  $f$  in  $\mathcal{X}_*$  and  $\langle \cdot, \cdot \rangle_\alpha$  stands for the inner product in  $L^2(\mu_\alpha)$ .

Fix a positive integer  $N$  and denote by  $\mathbb{T}_N^d = \{-N, \dots, N\}^d$  the discrete  $d$ -dimensional torus with  $N$  and  $-N$  identified, consisting of  $(2N)^d$  points. For the same transition probability  $p(\cdot)$ , one can consider the symmetric simple exclusion process evolving on  $\mathbb{T}_N^d$ . The state space is now  $\mathcal{X}_N = \{0, 1\}^{\mathbb{T}_N^d}$  and the generator  $L_N$  acting on any function  $f$  is given by

$$(L_N f)(\eta) = \sum_{x, y \in \mathbb{T}_N^d} p(y - x) \eta(x) [1 - \eta(y)] [f(\sigma^{x, y} \eta) - f(\eta)].$$

In a similar way, we can examine the evolution of a tagged particle on the torus  $\mathbb{T}_N^d$ . The environment as seen from the tagged particle is a Markov process with generator  $\mathcal{L}_N = \mathcal{L}_{0, N} + \mathcal{L}_{\tau, N}$ , where

$$\begin{aligned} (\mathcal{L}_{0, N} f)(\xi) &= \sum_{x, y \in \mathbb{T}_{N, *}^d} p(y - x) \xi(x) [1 - \xi(y)] [f(\sigma^{x, y} \xi) - f(\xi)], \\ (\mathcal{L}_{\tau, N} f)(\xi) &= \sum_{z \in \mathbb{T}_{N, *}^d} p(z) [1 - \xi(z)] [f(\tau_z \xi) - f(\xi)]. \end{aligned} \quad (2.10)$$

Let the total number  $K$  of particles be fixed and of course  $1 \leq K \leq (2N)^d$ . A simple computation shows that the measure  $\mu_{N, K}$ , which is uniform over the configurations of  $\mathcal{X}_{N, *} = \{0, 1\}^{\mathbb{T}_{N, *}^d}$  with  $K - 1$  particles is reversible with respect to  $\mathcal{L}_N$ . Since we are in a finite state space it is not hard to prove an invariance principle for the motion  $X_t^N$  of the tagged particle lifted to  $\mathbb{Z}^d$  in this context:  $\varepsilon X_{t\varepsilon^{-2}}^N$  converges as  $\varepsilon \downarrow 0$  to a Brownian motion with diffusion coefficient  $D_{N, K}$  given by

$$\langle \lambda, D_{N, K} \lambda \rangle = \left( 1 - \frac{K - 1}{(2N)^d - 1} \right) \sum_{z \in \mathbb{Z}_*^d} (z \cdot \lambda)^2 p(z) - 2 \|\psi_\lambda\|_{-1, N, K}^2, \quad (2.11)$$

where  $\|\psi_\lambda\|_{-1,N,K}^2$  stands for the  $H_{-1}$  norm of  $\psi_\lambda$  in  $L^2(\mu_{N,K})$ , given by the variational formula

$$\|\psi_\lambda\|_{-1,N,K}^2 = \sup_f \left\{ 2 \langle \psi_\lambda, f \rangle_{N,K} - \langle f, (-\mathcal{L}_N)f \rangle_{N,K} \right\}. \quad (2.12)$$

Here the supremum is carried over all functions in  $L^2(\mu_{N,K})$ .

The main result of this article concerns the behavior of the finite volume diffusion coefficient  $D_{N,K}$  as  $N \uparrow \infty$ :

**Theorem 2.5 .** *Fix  $\alpha$  in  $(0, 1)$  and a sequence  $K_N$  such that  $K_N/(2N)^d$  converges to  $\alpha$  as  $N \uparrow \infty$ . Then,*

$$\lim_{N \rightarrow \infty} D_{N,K_N} = D(\alpha).$$

We conclude this section with a sketch of the proof of Theorem 2.5. In view of formulae (2.7), (2.11), Theorem 2.5 follows from the convergence of the  $H_{-1}$  norms of  $\psi_\lambda$ . It is enough to show that

$$\lim_{\substack{N \uparrow \infty \\ K_N/(2N)^d \rightarrow \alpha}} \langle \psi_\lambda, (-\mathcal{L}_N)^{-1} \psi_\lambda \rangle_{N,K_N} = \langle \psi_\lambda, (-\mathcal{L})^{-1} \psi_\lambda \rangle_\alpha. \quad (2.13)$$

Here, to emphasize the dependence of the  $H_{-1}$  norm on the generators, we have denoted the  $H_{-1}$  norms  $\|\cdot\|_{-1,N,K}^2$  and  $\|\cdot\|_{-1,\alpha}^2$  respectively by  $\langle \cdot, (-\mathcal{L}_N)^{-1} \cdot \rangle_{N,K}$  and  $\langle \cdot, (-\mathcal{L})^{-1} \cdot \rangle_\alpha$ .

The generator  $\mathcal{L}$  has two pieces. The first one,  $\mathcal{L}_0$ , corresponds to jumps of the environment and the second one,  $\mathcal{L}_\tau$ , corresponds to jumps of the tagged particle. From Corollary 2.4 we conclude that

$$\lim_{\substack{N \uparrow \infty \\ K_N/(2N)^d \rightarrow \alpha}} \langle \psi_\lambda, (-\mathcal{L}_{0,N})^{-1} \psi_\lambda \rangle_{N,K_N} = \langle \psi_\lambda, (-\mathcal{L}_0)^{-1} \psi_\lambda \rangle_\alpha. \quad (2.14)$$

In section 4 we prove, in addition that, for each  $\varepsilon > 0$ , we may express the local function  $\psi_\lambda$  as the sum of  $\mathcal{L}g_\varepsilon + u_\varepsilon$ , where  $g_\varepsilon, u_\varepsilon$  are local functions,  $u_\varepsilon$  has mean zero with respect to all invariant measures and  $\|u_\varepsilon\|_{0,-1}^2 = \langle u_\varepsilon, (-\mathcal{L}_0)^{-1} u_\varepsilon \rangle_\alpha \leq \varepsilon^2$ . Since  $\|\cdot\|_{-1} \leq \|\cdot\|_{0,-1}$  and (2.13) is obvious if we replace  $\psi_\lambda$  by a local function of the form  $\mathcal{L}f$ , (2.13) follows from the previous decomposition and (2.14). Details are given in section 4.

### 3. FORMAL GRADIENTS OF THE SYMMETRIC EXCLUSION PROCESS

We prove in this section Theorems 2.1, 2.3 and Corollaries 2.2 and 2.4. The proof relies on the duality properties of the symmetric exclusion process that we now explain. Let  $\mathbb{X}$  be a countable set. For us it will be either  $\mathbb{Z}^d$  or  $\mathbb{Z}^d - \{0\}$ . We denote by  $\mathcal{X} = \{0, 1\}^{\mathbb{X}}$  the set of configurations  $\eta$  on  $\mathbb{X}$ . For  $0 \leq \alpha \leq 1$ ,  $\nu_\alpha$  is the Bernoulli product measure on  $\mathcal{X}$  with  $P[\eta(x) = 1] = \alpha$  for each  $x \in \mathbb{X}$ . Averages or norms with respect to  $\nu_\alpha$  will be denoted by the subscript  $\alpha$ . Let  $\mathbb{E}$  be a collection

of edges  $e = (x, y)$  connecting some pairs of points in  $\mathbb{X}$ . Assume that the resulting graph is connected and that every vertex in it has a finite degree. There will also be given transition rates  $\{p(x, y) : \mathbb{X} \times \mathbb{X} \rightarrow [0, \infty)\}$  that are symmetric and satisfy  $\sum_{y \in \mathbb{X}} p(x, y) \leq 1$  for every  $x \in \mathbb{X}$ . Moreover  $\mathbb{E} = \{e = (x, y) : p(x, y) > 0\}$ . In our context  $p(x, y) = p(y - x)$  for  $x, y \in \mathbb{Z}^d$  or  $x, y \in \mathbb{Z}_*^d$  as the case may be. We have a simple exclusion process on  $\mathcal{X}$  with generator

$$(\mathcal{L}f)(\eta) = \sum_{x, y \in \mathbb{X}} p(x, y) \eta(x) (1 - \eta(y)) [f(\sigma^{x, y} \eta) - f(\eta)]$$

and Dirichlet form

$$D(f) = \frac{1}{4} \sum_{x, y \in \mathbb{X}} p(x, y) \|f(\sigma^{x, y} \eta) - f(\eta)\|_\alpha^2.$$

For each  $n \geq 0$ , denote by  $\mathcal{E}_n$  the subsets of  $\mathbb{X}$  with  $n$  points. For each  $A$  in  $\mathcal{E}_n$ , let  $\Psi_A$  be the local function

$$\Psi_A = \prod_{x \in A} \frac{\eta(x) - \alpha}{\sqrt{\alpha(1 - \alpha)}}. \quad (3.1)$$

By convention,  $\Psi_\emptyset = 1$ . It is easy to check that  $\{\Psi_A, A \subset \mathbb{X}\}$  is an orthonormal basis of  $L^2(\nu_\alpha)$ . For each  $n \geq 0$ , denote by  $\mathcal{H}_n$  the subspace of  $L^2(\nu_\alpha)$  generated by  $\{\Psi_A, A \in \mathcal{E}_n\}$ , so that  $L^2(\nu_\alpha) = \bigoplus_{n \geq 0} \mathcal{H}_n$ . Functions of  $\mathcal{H}_n$  are said to have degree  $n$ .

For  $n \geq 0$ , denote by  $\pi_n$  the projection of  $L^2(\nu_\alpha)$  on  $\mathcal{H}_n$ . It follows from definition (2.3) of formal gradients that  $\{\pi_n \Phi_e, e \in \mathbb{E}\}$  is a formal gradient if  $\{\Phi_e, e \in \mathbb{E}\}$  is one:

**Lemma 3.1** . *Let  $\{\Phi_e, e \in \mathbb{E}\}$  be a formal gradient. Then, for every  $n \geq 0$ ,  $\{\pi_n \Phi_e, e \in \mathbb{E}\}$  is also a formal gradient.*

*Proof.* The proof is elementary. We verify that the projection  $\pi_n$  commutes with the exchanges  $\sigma^e$ , which follows from a direct computation.  $\square$

This lemma allows us to decompose a formal gradient into components of different degrees. We shall say that a formal gradient  $\{\Phi_e, e \in \mathbb{E}\}$  has degree  $n$  if each function  $\Phi_e$  has degree  $n$ . The proof of Theorems 2.1 and 2.3 is thus reduced to proving that a square integrable formal gradient of degree  $n$  can be approximated by gradients of local functions of degree  $n$ .

Let us compute  $\sigma^e \Psi_A$ . If  $e = (x, y)$ ,  $\sigma^e \Psi_A = 0$  unless  $e$  connects  $A$  and  $A^c$ . If  $x \in A$  and  $y \notin A$ , then  $\sigma^e \Psi_A = \Psi_{(A \setminus \{x\}) \cup \{y\}}$  and if  $y \in A$  and  $x \notin A$ , then  $\sigma^e \Psi_A = \Psi_{(A \setminus \{y\}) \cup \{x\}}$ . In other words  $\sigma^e \Psi_A = \Psi_{\sigma^e A}$ , with the obvious definition of  $\sigma^e A = (A \setminus \{x\}) \cup \{y\}$  or  $(A \setminus \{y\}) \cup \{x\}$ . This suggests that we should turn the space  $\mathbb{X}_n$  of subsets  $A$  of cardinality  $n$  in  $\mathbb{X}$  into a graph with edges  $\mathbb{E}_n$  by connecting two vertices  $A$  and  $B$  (i.e. subsets of  $\mathbb{X}$  of cardinality  $n$ ) in  $\mathbb{X}_n$  with

an edge if  $B = (A \setminus \{x\}) \cup \{y\}$  for some  $x \in A$  and  $y \notin A$  that are connected by an edge in  $\mathbb{X}$ . The transition probabilities  $p_n(A, B)$  on  $\mathbb{X}_n$  are given by  $p(x, y)$  if  $B = (A \setminus \{x\}) \cup \{y\}$  for some  $x \in A$  and  $y \notin A$  and 0 otherwise.

Any local function  $f$  of degree  $n$  has a representation as

$$f(\eta) = \sum_A f(A) \Psi_A(\eta)$$

and

$$(\sigma^e f)(\eta) = f(\sigma^e \eta) = \sum_A f(A) \sigma^e \Psi_A = \sum_A f(\sigma^e A) \Psi_A .$$

The Dirichlet form  $D(f)$  for a function of degree  $n$  reduces to

$$\frac{1}{2} \sum_{\substack{e \in \mathbb{E} \\ A \in \mathbb{X}_n}} p(e) |f(\sigma^e A) - f(A)|^2 .$$

The factor 1/2 is explained by the sum over the unoriented edges. The total Dirichlet form of  $f$  can be written as the sum

$$\frac{1}{2} \sum_{n \geq 1} \sum_{\substack{e \in \mathbb{E} \\ A \in \mathbb{X}_n}} p(e) |\pi_n f(\sigma^e A) - \pi_n f(A)|^2 ,$$

where

$$\pi_n f = \sum_{A \in \mathbb{X}_n} f(A) \Psi_A .$$

The approximation of any square integrable formal gradient in  $\mathbb{X}$  by gradients of local functions reduces to a similar property for each of the graphs  $\mathbb{X}_n$ . Note that for the purpose of the approximation result we can replace the Dirichlet form by any equivalent one.

Let  $\mathcal{G}$  be a connected graph with vertices  $V = \{x\}$  and edges  $\mathbb{E} = \{e\}$ . We assume that the graph is countable and that there is an upper bound  $D$  on the degree of all of its vertices. Let us pick for each edge an arbitrary orientation as the positive direction but keep the edges in both directions so we have the set  $\widehat{\mathbb{E}}$  of oriented edges. Given function  $f : V \rightarrow \mathbb{R}$  we can define  $g = \delta f : \widehat{\mathbb{E}} \rightarrow \mathbb{R}$  by  $(\delta f)(e) = f(y) - f(x)$  where  $(x, y) = e$  is an oriented edge. It is clear that  $g : \widehat{\mathbb{E}} \rightarrow \mathbb{R}$  satisfies

$$\sum_{e \in \ell} g(e) = 0 \tag{3.2}$$

for any loop  $\ell$  of oriented edges. We can turn the set of  $g$ 's satisfying (3.2) into a Hilbert space  $\mathcal{H}_1$  by defining

$$\|g\|^2 = \sum_{e \in \mathbb{E}} |g(e)|^2 = \frac{1}{2} \sum_{e \in \widehat{\mathbb{E}}} |g(e)|^2 . \tag{3.3}$$

Given any  $g$  satisfying (3.2) we can construct an  $f$ , unique up to an additive constant, such that  $g = \delta f$ . However, in general we have no control on the growth of  $f$  at  $\infty$ . We say that the graph has property **LD**, if for any given  $g \in \mathcal{H}_1$  we can find a sequence of functions  $f_n$  each of which vanishes outside a finite set of vertices, such that

$$\lim_{n \rightarrow \infty} \|g - \delta f_n\| = 0$$

(cf. [G]). Not every graph has this property: a counterexample is given by the homogeneous tree of degree 3 (cf. [S], example (3.19), pag. 37).

The following lemma will allow us deduce property **LD** for a graph from a closely related one.

**Lemma 3.2 .** *Let  $\mathcal{G}$  be a graph with the degree of its vertices bounded by  $D$ . Let  $\mathcal{G}_1$  be obtained from  $\mathcal{G}$  by deleting a set of vertices and retaining all the edges of  $\mathcal{G}$  that are contained in  $\mathcal{G}_1$ . Assume that there are finite constants  $c(k)$ , depending only on  $(k, \mathcal{G}_1)$  such that any two vertices in  $\mathcal{G}_1$  that are connected by a path of  $k$  edges in  $\mathcal{G}$  are connected by a path of at most  $c(k)$  edges contained in  $\mathcal{G}_1$ . Assume also that there is a bound  $\ell$  such that any deleted vertex of  $\mathcal{G}$  is connected to an undeleted vertex in  $\mathcal{G}_1$  by a path of edges in  $\mathcal{G}$  of length at most  $\ell$ . Then if  $\mathcal{G}$  has property **LD** so does  $\mathcal{G}_1$ .*

*Proof.* The proof proceeds along the following lines. Given a  $g$  defined on the oriented edges of  $\mathcal{G}_1$  that satisfies (3.2) we can find  $f$  on the vertices of  $\mathcal{G}_1$  such that  $g(e) = (\delta f)(e)$ . Assume that we can extend  $f$  as a function  $h$  defined on the vertices of  $\mathcal{G}$  in such a way that  $\|\delta h\|_{\mathcal{G}}^2 \leq C\|g\|_{\mathcal{G}_1}^2$ . Because  $\mathcal{G}$  has property **LD** we can find a sequence of functions  $h_n$  defined on the vertices of  $\mathcal{G}$ , each  $h_n$  having finite support such that  $\|\delta h_n - \delta h\|_{\mathcal{G}}^2 \rightarrow 0$  as  $n \rightarrow \infty$ . It follows that the restriction  $f_n$  of  $h_n$  to  $\mathcal{G}_1$  satisfies  $\|\delta f_n - g\|_{\mathcal{G}_1}^2 \rightarrow 0$  proving that  $\mathcal{G}_1$  has the property **LD** as well.

This shows that all we need to do is to construct such extension  $h$  of  $f$ . In order to do this let us proceed as follows. For any  $x$  that is a deleted vertex, there is at least one vertex  $x'$  of  $\mathcal{G}_1$  within a distance  $\ell$ . Pick any one. Define  $h(x) = f(x')$ . Then if  $e = (x, y)$  is a deleted edge, then  $(\delta h)(e) = f(x') - f(y')$  where  $x', y'$  are vertices in  $\mathcal{G}_1$ . The graph distance between  $x'$  and  $y'$  can be at most  $2\ell + 1$  in  $\mathcal{G}$  and therefore at most  $c(2\ell + 1)$  in  $\mathcal{G}_1$ . It is clear that there is a constant  $K = K(\ell, c(\cdot))$  such that all the edges involved in the path from  $x'$  to  $y'$  through the vertices of  $\mathcal{G}_1$  are within a distance  $K$  from  $x$  and  $y$ . For any edge  $e$  let us define

$$S_K(e) = \sum_{\substack{e': d(e, e') \leq K \\ e' \in \mathcal{G}_1}} |(\delta f)(e')|^2,$$

where the sum extends only over the edges of  $\mathcal{G}_1$  within a distance  $K$  of  $e$ . We can define the distance between two edges as the largest of the four possible distances between the two pairs of vertices. They differ from one another by at most 2. By an application of Schwarz's inequality and the crudest upper bound

$$|(\delta h)(e)|^2 \leq c(2\ell + 1)S_K(e)$$

and

$$\|\delta h\|_{\mathcal{G}}^2 \leq c(2\ell + 1) \sum_e S_K(e) = C c(2\ell + 1) \|\delta f\|_{\mathcal{G}_1}^2,$$

where

$$C = \sup_e \{ |e'| : d(e, e') \leq K \}.$$

Clearly  $C \leq C(K, D) = (D + D^2 + \dots + D^K)$ , where  $D$  is a bound on the degree of any vertex.  $\square$

**Lemma 3.3 .** *Any symmetric random walk on  $\mathbb{Z}^d$  has property **LD**.*

*Proof.* We can assume without loss of generality that the walk is the nearest neighbor one. On any graph, if  $\delta f$  with  $f$  having finite support is not dense in the Hilbert space  $\mathcal{H}_1$ , there will be a  $g \in \mathcal{H}_1$  that is orthogonal to  $\delta f$  for all finitely supported  $f$ . In other words, if we represent  $g = \delta h$ , after a summation by parts

$$\sum_e (\delta h)(e)(\delta f)(e) = \sum_e (\delta^* \delta h)(e) f(e) = 0 \text{ for all } f.$$

For the operator

$$L = \delta^* \delta$$

we have a solution of  $Lh \equiv 0$  with  $\delta h \in \mathcal{H}_1$ . We will show that this is impossible for the simple random walk on any  $\mathbb{Z}^d$  (except for the trivial solution  $h = 0$ ). Define  $h_i(x) = h(x + z_i) - h(x)$  where  $z_i : 1 \leq i \leq d$  are the unit vectors in the coordinate directions. Then (3.2) implies  $(e^{i\theta_j} - 1)\hat{h}_k(\theta) = (e^{i\theta_k} - 1)\hat{h}_j(\theta)$  a.e. for  $1 \leq j < k \leq d$ , where  $\hat{h}_j$  is the Fourier transform of the  $\ell_2$  function  $h_j(x)$ . Therefore there exists a function  $u(\theta)$  on the  $d$ -torus such that for  $1 \leq j \leq d$ ,  $\hat{h}_j(\theta) = (e^{i\theta_j} - 1)u(\theta)$ . It is easy to see that  $Lh = 0$  now implies  $k(\theta)u(\theta) = 0$ , where  $k(\theta) = \sum_j (1 - \cos \theta_j) > 0$  for  $\theta \neq 0$ . Thus  $\hat{h}_j \equiv 0$ .  $\square$

Clearly, the proof only depends on the property

$$\sum_x p(x) e^{i\theta \cdot x} = \sum_x p(x) \cos(\theta \cdot x) < 1$$

for  $\theta \neq 0$  which is the same as irreducibility.

**Lemma 3.4 .** *Any symmetric exclusion process involving a finite number  $k$  of particles on  $\mathbb{Z}_*^d$  has property **LD**.*

*Proof.* With out exclusion and allowing 0, we have essentially a symmetric random walk on  $\mathbb{Z}^{kd}$  which has property **LD**. (We only have to deal with symmetric functions of the  $k$  positions  $x_1, \dots, x_k$ ). We now delete the set of vertices  $A = \{(x_1, \dots, x_k)\}$ , where either some  $x_i = 0$  or  $x_i = x_j$  for some pair  $i, j$ . It is easy to verify that  $A$  satisfies the conditions of Lemma 3.2 except when  $d = 1$  and the walk is the nearest neighbor one.  $\square$

**Remark 3.5 .** *Tom Liggett pointed out to us that this result can be also obtained by using coupling techniques due to Spitzer and Liggett, as presented in [Li], chapter 8, section 1.*

We have thus proved Theorems 2.1 and 2.3. We shall now prove Corollaries 2.2 and 2.4. Since the proofs are identical we will deal with only Corollary 2.2.

*Proof of Corollary 2.2.* Consider a local function  $h$  such that  $\langle h \rangle_\alpha \equiv 0$  for any  $\alpha \in [0, 1]$ . This is equivalent to  $\langle h \rangle_{N,K} \equiv 0$  for any  $K$  and  $N$  sufficiently large. We define for such a function

$$\begin{aligned} \|h\|_{-1,\alpha}^2 &= \sup_f \left\{ 2 \langle h, f \rangle_\alpha - \langle f, (-\mathcal{L})f \rangle_\alpha \right\} \\ \|h\|_{-1,N,K}^2 &= \sup_f \left\{ 2 \langle h, f \rangle_{N,K} - \langle f, (-\mathcal{L}_N)f \rangle_{N,K} \right\}. \end{aligned}$$

In the first formula one uses only local functions  $f$ . The second formula is in a finite dimensional space and we are free to use arbitrary functions  $f$ . We need to prove

$$\lim_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K_N}^2 = \|h\|_{-1,\alpha}^2. \quad (3.4)$$

It is easy to see that

$$\|h\|_{-1,\alpha}^2 \leq \liminf_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K_N}^2. \quad (3.5)$$

Indeed, by definition, for every  $\varepsilon > 0$ , there exists a local function  $f_\varepsilon$  such that

$$\|h\|_{-1,\alpha}^2 \leq 2 \langle h, f_\varepsilon \rangle_\alpha - \langle f_\varepsilon, (-\mathcal{L})f_\varepsilon \rangle_\alpha + \varepsilon.$$

Since  $h$  and  $f_\varepsilon$  are local functions, the right hand side is equal to

$$\begin{aligned} & \lim_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \{ 2 \langle h, f_\varepsilon \rangle_{N,K} - \langle f_\varepsilon, (-\mathcal{L})f_\varepsilon \rangle_{N,K} \} + \varepsilon \\ &= \lim_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \{ 2 \langle h, f_\varepsilon \rangle_{N,K} - \langle f_\varepsilon, (-\mathcal{L}_N)f_\varepsilon \rangle_{N,K} \} + \varepsilon \\ &\leq \liminf_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K}^2 + \varepsilon. \end{aligned}$$

The reverse inequality is more demanding. Our goal is to prove that

$$\limsup_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K_N} \leq \|h\|_{-1,\alpha}. \quad (3.6)$$

It is more convenient to express the norms  $\|h\|_{-1,N,K}$  and  $\|h\|_{-1,\alpha}$  as

$$\|h\|_{-1,\alpha} = \sup \left\{ \langle h, f \rangle_\alpha : \langle f, -\mathcal{L}f \rangle_\alpha \leq 1 \right\}, \quad (3.7)$$

$$\|h\|_{-1,N,K} = \sup \left\{ \langle h, f \rangle_\alpha : \langle f, -\mathcal{L}_N f \rangle_{N,K} \leq 1 \right\}, \quad (3.8)$$

where the suprema are taken over  $f$  from the unit balls in the corresponding Dirichlet spaces.

It is easy to see that the following lemma is equivalent to (3.6). In fact we can choose  $f_N$  as the solution of the variational problem (3.8) and applying the Lemma 3.6 we obtain (3.6).  $\square$

**Lemma 3.6 .** *Let  $K_N/(2N)^d \rightarrow \alpha$ . Assume that for some  $A \geq 0$  we have a sequence of functions  $f_N$  in  $L_2(\mu_{N,K_N})$  that satisfy*

$$\begin{aligned} \sup_N \langle f_N, -\mathcal{L}_N f_N \rangle_{N,K_N} &\leq 1, \\ \lim_{N \rightarrow \infty} \langle h, f_N \rangle_{N,K_N} &= A. \end{aligned} \quad (3.9)$$

Then, for any  $\varepsilon > 0$ , there is a local function  $f_\varepsilon$  such that

$$\langle f_\varepsilon, h \rangle_\alpha \geq A - \varepsilon \text{ and } \langle f_\varepsilon, -\mathcal{L} f_\varepsilon \rangle_\alpha \leq 1. \quad (3.10)$$

*Proof of Lemma 3.6.* Fix a positive integer  $\ell$  large enough so that  $\Lambda_\ell = \{-\ell, \dots, \ell\}^d$  contains the support of  $h$ . Since  $h$  has mean zero with respect to all the canonical measures  $\mu_{\ell,K}$ , from linear algebra,  $H = \mathcal{L}_{\Lambda_\ell}^{-1} h$  is a well defined cylinder function. Here  $\mathcal{L}_{\Lambda_\ell}$  is the generator of the exclusion process restricted to  $\Lambda_\ell$ . Notice the difference between  $\mathcal{L}_{\Lambda_\ell}$  and  $\mathcal{L}_\ell$ . While the first generator stands for the exclusion process on the cube  $\Lambda_\ell$ , the second generator stands for the exclusion process on  $\Lambda_\ell$  with periodic boundary conditions, that is, on  $\mathbb{T}_\ell^d$ . Since  $h = \mathcal{L}_{\Lambda_\ell} H$ , for any  $N > \ell$

$$\langle h, f_N \rangle_{N,K_N} = -\frac{1}{2} \sum_{x,y \in \Lambda_\ell} p(y-x) \langle T_{x,y} f_N, T_{x,y} H \rangle_{N,K_N}. \quad (3.11)$$

The summation in the above formula is carried over all sites in the cube  $\Lambda_\ell$  (without periodic boundary condition). Fix a cube  $\Lambda_L$  with  $L \geq \ell$ . Denote by  $f_{N,L}$  the conditional expectation of  $f_N$  given  $\mathcal{F}_{\Lambda_L} = \sigma(\eta(x), x \in \Lambda_L)$ . A simple computation shows that  $E[T_{x,y} f_N | \mathcal{F}_{\Lambda_L}] = T_{x,y} E[f_N | \mathcal{F}_{\Lambda_L}] = T_{x,y} f_{N,L}$  for all bonds  $\{x, y\}$  in  $\Lambda_\ell$ . Therefore, since  $H$  is  $\mathcal{F}_{\Lambda_L}$ -measurable, taking conditional expectations with respect to  $\mathcal{F}_{\Lambda_L}$ , we obtain that the right hand side of (3.11) is equal to

$$-\frac{1}{2} \sum_{x,y \in \Lambda_\ell} p(y-x) \langle T_{x,y} f_{N,L}, T_{x,y} H \rangle_{N,K_N}. \quad (3.12)$$

On the other hand, by Jensen's inequality and (3.9)

$$\begin{aligned} &\frac{1}{2} \sum_{x,y \in \Lambda_L} p(x-y) \langle (T_{x,y} f_{N,L})^2 \rangle_{N,K_N} \\ &\leq \frac{1}{2} \sum_{x,y \in \Lambda_L} p(x-y) \langle (T_{x,y} f_N)^2 \rangle_{N,K_N} \\ &= \langle f_N, -\mathcal{L}_N f_N \rangle_{N,K_N} \\ &\leq 1. \end{aligned} \quad (3.13)$$

Since  $f_{N,L}$  is a cylinder function, the state space  $\{0, 1\}^{\Lambda_L}$  is finite and we have a uniform bound on the  $L^2(\mu_{N,K_N})$  norms of the gradients of  $f_{N,L}$ , there exists a subsequence  $N'$  and a cylinder function  $g_L$  such that the gradients of  $f_{N',L}$  converge to the gradients of  $g_L$  pointwise. On the other hand, by the equivalence of ensembles, as  $N \uparrow \infty$  and  $K_N/(2N)^d \rightarrow \alpha$ , the finite dimensional marginals of the canonical measure  $\mu_{N,K_N}$  converge to the corresponding marginals of  $\mu_\alpha$ . Therefore, as  $N' \uparrow \infty$  and as  $K_{N'}/(2N')^d \rightarrow \alpha$ , the expression in (3.12) converges to

$$-\frac{1}{2} \sum_{x,y \in \Lambda_\ell} p(y-x) \langle T_{x,y} H, T_{x,y} g_L \rangle_\alpha \cdot \quad (3.14)$$

Moreover, since  $T_{x,y} f_{N,L}$  converges to  $T_{x,y} g_L$ , by (3.13),

$$\begin{aligned} & \frac{1}{2} \sum_{x,y \in \Lambda_L} p(x-y) \langle \{T_{x,y} g_L\}^2 \rangle_\alpha \\ &= \frac{1}{4} \sum_{x,y \in \Lambda_L} p(x-y) \langle \{\nabla_{x,y} g_L\}^2 \rangle_\alpha \\ &\leq 1 \end{aligned} \quad (3.15)$$

uniformly in  $L$ .

For each  $x, y$  in  $\mathbb{Z}^d$  such that  $y-x \in F$  where  $F = \{z : p(z) > 0\}$ ,  $T_{x,y} g_L$  is a bounded sequence in  $L^2(\mu_\alpha)$ . There exists, therefore, a subsequence  $L'$  for which  $T_{x,y} g_{L'}$  converges weakly for each  $x, y$  in  $\mathbb{Z}^d$  such that  $y-x \in F$ . Denote by  $\Phi_{x,y}$  the weak  $L^2(\mu_\alpha)$  limit. By construction, as  $L' \uparrow \infty$ , (3.14) converges to

$$-\frac{1}{2} \sum_{x,y \in \Lambda_\ell} p(x-y) \langle T_{x,y} H, \Phi_{x,y} \rangle_\alpha \cdot$$

Moreover, the collection of  $L^2(\mu_\alpha)$ -functions  $\Phi_{x,y}$  satisfy conditions (2.3) of a formal gradient.

At this point we have proved that

$$-\frac{1}{2} \sum_{x,y \in \Lambda_\ell} p(x-y) \langle T_{x,y} H, \Phi_{x,y} \rangle_\alpha \geq A \quad (3.16)$$

and from (3.15) it follows that

$$\frac{1}{4} \sum_{x,y \in \mathbb{Z}^d} p(x-y) \langle \Phi_{x,y}^2 \rangle_\alpha \leq 1 \cdot \quad (3.17)$$

We can now apply Theorem 2.1 and approximate  $\{\Phi_{x,y}\}$  by  $\delta f_\varepsilon$  thereby establishing (3.10).  $\square$

## 4. APPROXIMATIONS FOR THE TAGGED PARTICLE

Given,  $1 \leq K \leq (2N)^d$ , denote by  $\mathcal{X}_{N,K}$  the space of configurations of  $\mathcal{X}_N$  with  $K - 1$  particles:  $\mathcal{X}_{N,K} = \{\xi \in \mathcal{X}_N, \sum_{x \in \mathbb{T}_{N,*}^d} \xi(x) = K - 1\}$ . Recall from section 2 that  $\mu_{N,K}$  stands for the uniform measure on  $\mathcal{X}_{N,K}$  and that  $\langle \cdot \rangle_{N,K}$  represents the expectation with respect to  $\mu_{N,K}$ .

Denote by  $\mathcal{C}$  (resp.  $\mathcal{C}_0$ ) the space of local functions (resp. local functions that have mean zero with respect to all canonical measures). Thus, a local function  $h$  belongs to  $\mathcal{C}_0$  provided  $\langle h \rangle_{N,K} = 0$  for all large enough  $N$  and all  $1 \leq K \leq (2N)^d$ . Examples of subspaces of  $\mathcal{C}_0$  are  $\mathcal{L}_0\mathcal{C} = \{\mathcal{L}_0h, h \in \mathcal{C}\}$ ,  $\mathcal{L}_\tau\mathcal{C} = \{\mathcal{L}_\tau h, h \in \mathcal{C}\}$ . Also  $\{\psi_\lambda, \lambda \in \mathbb{R}^d\}$  defined by (2.8) are functions in  $\mathcal{C}_0$ .

On the space of local functions  $\mathcal{C}$ , consider the inner product  $\langle \cdot, \cdot \rangle_{0,\alpha,1}$  defined by

$$\langle f, g \rangle_{0,1,\alpha} = \langle -\mathcal{L}_0f, g \rangle_\alpha$$

Here the index 0 signifies that only the jump part of the generator enters in the definition. Denote by  $\mathcal{H}_{1,\alpha}^0$  the Hilbert space obtained by completing the space of local functions endowed with the above inner product and by  $D_{0,\alpha}$  the Dirichlet form associated to the generator  $\mathcal{L}_0$  and the probability measure  $\nu_\alpha$ : i.e.  $D_{0,\alpha}(f) = \langle -\mathcal{L}_0f, f \rangle_\alpha$ . For a local function  $f$ , let

$$\|f\|_{0,-1,\alpha}^2 = \sup_{g \in \mathcal{C}} \left\{ 2 \langle f, g \rangle_\alpha - D_{0,\alpha}(g) \right\}$$

be the dual  $H_{-1}$  norm.

**Remark 4.1** . *We could have defined the  $H_{-1}$  norm as*

$$\|f\|_{0,-1,\alpha} = \sup \left\{ \langle f, g \rangle_\alpha : D_{0,\alpha}(g) \leq 1 \right\},$$

where the supremum is taken over either local functions or functions in  $L_2(\nu_\alpha)$  with Dirichlet norm bounded by 1.

$\|f\|_{0,-1,\alpha}$  is called the  $\mathcal{H}_{-1,\alpha}^0$ -norm of  $f$ . It is known that every local function in  $\mathcal{C}_0$  has finite  $\mathcal{H}_{-1,\alpha}^0$ -norm (cf. [KL], section 7.2). On the other hand, constants have infinite  $\mathcal{H}_{-1,\alpha}^0$ -norm. For two local functions  $f, g$  with finite  $\mathcal{H}_{-1,\alpha}^0$ -norm, define by polarization the  $\mathcal{H}_{-1,\alpha}^0$  inner product of  $f, g$ , denoted by  $\langle f, g \rangle_{0,-1,\alpha}$ . Let  $\mathcal{H}_{-1,\alpha}^0$  be the Hilbert space induced by the local functions endowed with the inner product  $\langle f, g \rangle_{0,-1,\alpha}$ .

In a similar way, for each  $N \geq 1$  and  $1 \leq K \leq (2N)^d$ , we define the spaces  $\mathcal{H}_{1,N,K}^0, \mathcal{H}_{-1,N,K}^0$  and the norms  $\|\cdot\|_{0,1,N,K}$  and  $\|\cdot\|_{0,-1,N,K}$ . According to Corollary 2.4, for any local function in  $h \in \mathcal{C}_0$ ,

$$\lim_{\substack{N \rightarrow \infty \\ K_N / (2N)^d \rightarrow \alpha}} \|h\|_{0,-1,N,K_N}^2 = \|h\|_{0,-1,\alpha}^2. \quad (4.1)$$

Since the density  $\alpha$  is fixed throughout this section, to keep notation simple, for  $n = -1, 1$ , we denote the norms  $\|\cdot\|_{0,n,\alpha}$  and  $\|\cdot\|_{n,\alpha}$  by  $\|\cdot\|_{0,n}$  and  $\|\cdot\|_n$  and we denote the inner product  $\langle \cdot, \cdot \rangle_\alpha$  simply by  $\langle \cdot, \cdot \rangle$ . The main result of this section states that any  $h \in \mathcal{C}_0$  can be approximated in  $\|\cdot\|_{0,-1}$  norm by  $\mathcal{L}f$  with a local function  $f$ . Notice that while the  $H_{-1}$  norm corresponds to the generator  $\mathcal{L}_0$ , the generator that appears in front of  $f$  is the full generator  $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_\tau$ .

**Theorem 4.2 .** *Let  $h$  be a local function in  $\mathcal{C}_0$ . For every  $\varepsilon > 0$ , there exist local functions  $u_\varepsilon, f_\varepsilon$  such that  $h = \mathcal{L}f_\varepsilon + u_\varepsilon$  and  $\|u_\varepsilon\|_{0,-1} \leq \varepsilon$ .*

Notice that  $u_\varepsilon$  belongs to  $\mathcal{C}_0$  because  $\mathcal{L}f_\varepsilon$  belongs automatically to  $\mathcal{C}_0$ . Theorem 4.2 is based on the following two results. Fix a local function  $h$  in  $\mathcal{C}_0$ . For each  $\lambda > 0$ , let  $u_\lambda$  be the solution of the resolvent equation

$$\lambda u_\lambda - \mathcal{L}u_\lambda = -h. \quad (4.2)$$

**Proposition 4.3 .** *For every  $\varepsilon > 0$ , there exists  $\lambda_0 > 0$  such that  $\|\mathcal{L}u_\lambda - h\|_{0,-1} \leq \varepsilon$  for all  $\lambda < \lambda_0$ . In other words*

$$\lim_{\lambda \rightarrow 0} \|\mathcal{L}u_\lambda - h\|_{0,-1} = 0. \quad (4.3)$$

**Proposition 4.4 .** *Fix a local function  $h$  in  $\mathcal{C}_0$  and let  $u_\lambda$  be the solution of the resolvent equation (4.2). For every  $\varepsilon > 0$  and every  $\lambda > 0$ , there exists a local function  $g_{\lambda,\varepsilon}$  such that  $\|\mathcal{L}(u_\lambda - g_{\lambda,\varepsilon})\|_{0,-1} \leq \varepsilon$ .*

*Proof of Theorem 4.2.* Since

$$h = \mathcal{L}g_{\lambda,\varepsilon} + \mathcal{L}(u_\lambda - g_{\lambda,\varepsilon}) + (h - \mathcal{L}u_\lambda)$$

and we can make

$$\|\mathcal{L}(u_\lambda - g_{\lambda,\varepsilon})\|_{0,-1} + \|(h - \mathcal{L}u_\lambda)\|_{0,-1} < \varepsilon$$

by first picking a small  $\lambda$  according to Proposition 4.3 to make the second term small and then by picking  $\varepsilon$  small according to Proposition 4.4 so that the first term small, Theorem 4.2 follows from the two Propositions 4.3 and 4.4.  $\square$

We now turn to the proofs of Propositions 4.3 and 4.4, which are based on the duality properties of the symmetric exclusion process explained in section 3. Denote by  $D_0(\cdot)$  the Dirichlet form associated to the jump part of the generator, by  $D_\tau(\cdot)$  the Dirichlet form associated to the translation part of the generator and by  $D(\cdot)$  the full Dirichlet form:

$$\begin{aligned} D_0(f) &= \langle f, (-\mathcal{L}_0)f \rangle \\ D_\tau(f) &= \langle f, (-\mathcal{L}_\tau)f \rangle \\ D(f) &= D_0(f) + D_\tau(f). \end{aligned}$$

Recall the definition of the set  $A_{x,y}$  given in section 3 and notice that  $\sigma^{x,y}\Psi_A = \Psi_{A_{x,y}}$ . In particular, if  $f = \sum_{n \geq 1} \sum_{A \in \mathcal{E}_n} f(A)\Psi_A$ ,

$$\begin{aligned} D_0(f) &= \frac{1}{4} \sum_{x,y \in \mathbb{Z}_*^d} p(y-x) \int [f(\sigma^{x,y}\xi) - f(\xi)]^2 d\nu_\alpha \\ &= \frac{1}{4} \sum_{n \geq 1} \sum_{x,y \in \mathbb{Z}_*^d} p(y-x) \sum_{A \in \mathcal{E}_n} \left\{ f(A_{x,y}) - f(A) \right\}^2. \end{aligned}$$

Actually if  $F$  is any symmetric finite set that generates  $\mathbb{Z}^d$ ,  $D_0(f)$  is seen to be equivalent to the form

$$\begin{aligned} \widehat{D}_0(f) &= \frac{1}{2} \sum_{\substack{x,y \in \mathbb{Z}_*^d \\ y-x \in F}} \int [f(\sigma^{x,y}\xi) - f(\xi)]^2 d\nu_\alpha \\ &= \frac{1}{2} \sum_{n \geq 1} \sum_{\substack{x,z \in \mathbb{Z}_*^d \\ y-x \in F}} \sum_{A \in \mathcal{E}_n} \left\{ f(A_{x,y}) - f(A) \right\}^2 \quad (4.4) \\ &= \sum_{n \geq 1} \sum_{e \in \mathbb{E}_n} \sum_{A \in \mathcal{E}_n} \left\{ f(\sigma^e A) - f(A) \right\}^2, \end{aligned}$$

where  $\mathbb{E}_n$ , is the set of edges in  $\mathcal{E}_n$  with an edge  $e$  connecting  $A$  and  $B$  present if  $A = \sigma^{x,y}B$ , for some  $x,y \in \mathbb{Z}_*^d$  such that  $y-x \in F$ . A consequence of the equivalence of the two Dirichlet forms is their equivalence on all of the spaces  $\mathcal{H}_n$  with constants that are independent of  $n$ .

On the other hand, for a set  $A$  of  $\mathcal{E}_n$ , denote by  $\tau_z A$  the set defined by

$$(\tau_z A) = \begin{cases} A + z & \text{if } -z \notin A, \\ (A + z) \setminus \{0\} \cup \{z\} & \text{if } -z \in A, \end{cases}$$

where  $B + z$  is the set  $\{x + z; x \in B\}$ . Therefore, in the case where  $-z$  belongs to  $A$ , we first translate  $A$  by  $z$  (obtaining a new set that contains the origin) and then we remove the origin and we add the site  $z$ . Of course  $\tau_z: \mathcal{E}_n \rightarrow \mathcal{E}_n$  is a one-to-one function. We denote its inverse by  $\tau_z^{-1}$ , which is seen to be  $\tau_{-z}$ . A straightforward computation shows that  $\tau_z \Psi_A = \Psi_{\tau_z A}$ .

We now use the results in the Appendix to prove Proposition 4.3 along the lines of the proofs of Lemma 2.5 of [LY] or Theorem 5.1 of [SVY].

*Proof of Proposition 4.3.* From the results in [KV] one can show that

$$\lim_{\lambda \rightarrow 0} \|\mathcal{L}u_\lambda - h\|_{-1} = 0 \quad (4.5).$$

Our goal is to prove that

$$\lim_{\lambda \rightarrow 0} \|\mathcal{L}u_\lambda - h\|_{0,-1} = 0.$$

To this end we write  $u_\lambda = \sum_n u_{n,\lambda}$  where  $u_{n,\lambda} = \pi_n u_\lambda$  is the projection of  $u_\lambda$  into  $\mathcal{H}_n$ , the space functions of degree  $n$  and a similar decomposition for  $\mathcal{L}u_\lambda = v_\lambda = \sum_n v_{n,\lambda}$  as well as  $h = \sum h_n$ . Since  $h$  is local, the decomposition for  $h$  is only a finite sum, and since the decomposition is orthogonal in  $\mathcal{H}_{-1,\alpha}^0$

$$\|\mathcal{L}u_\lambda - h\|_{0,-1}^2 = \sum_{n \leq n_0} \|v_{n,\lambda} - h_n\|_{0,-1}^2 + \sum_{n > n_0} \|v_{n,\lambda}\|_{0,-1}^2.$$

From (4.5) and Lemma 5.1

$$\|f\|_1^2 \leq C n \|f\|_{0,1}^2$$

valid for  $f \in \mathcal{H}_n$ , it follows that for each fixed  $n$  and any function  $u \in L^2$ ,

$$\begin{aligned} \|\pi_n u\|_{0,-1} &= \sup_{\substack{f \in \mathcal{H}_n \\ \|f\|_{0,1} \leq 1}} |\langle u, f \rangle| \leq \sup_{\|f\|_1 \leq \sqrt{Cn}} |\langle u, f \rangle| \\ &\leq \sqrt{Cn} \|u\|_{-1}. \end{aligned} \tag{4.6}$$

Then we have

$$\begin{aligned} \|v_{n,\lambda} - h_n\|_{0,-1} &\leq \sqrt{Cn} \|\mathcal{L}u_\lambda - h\|_{-1} \\ &\rightarrow 0 \quad \text{as } \lambda \rightarrow 0. \end{aligned}$$

It is therefore sufficient to control

$$\sum_n \sup_\lambda \|v_{n,\lambda}\|_{0,-1} < \infty.$$

On the other hand by the tridiagonal property of  $\mathcal{L}$  and using Lemma 5.1 and (4.6), we have

$$\begin{aligned} \|v_{n,\lambda}\|_{0,-1} &= \|\pi_n \mathcal{L}(u_{n-1,\lambda} + u_{n,\lambda} + u_{n+1,\lambda})\|_{0,-1} \\ &\leq \sqrt{Cn} \|\mathcal{L}(u_{n-1,\lambda} + u_{n,\lambda} + u_{n+1,\lambda})\|_{-1} \\ &= \sqrt{Cn} \|u_{n-1,\lambda} + u_{n,\lambda} + u_{n+1,\lambda}\|_1 \\ &\leq C'n \left\{ \|u_{n-1,\lambda}\|_{0,1} + \|u_{n,\lambda}\|_{0,1} + \|u_{n+1,\lambda}\|_{0,1} \right\}. \end{aligned}$$

In view of estimate (5.5) the proof of the proposition is complete.  $\square$

*Proof of Proposition 4.4.* Let  $u_\lambda$  be the solution of the resolvent equation. During the proof of Proposition 4.3 we saw that

$$\mathcal{L}u_\lambda = \mathcal{L} \left[ \sum_{n \geq 1} u_{\lambda,n} \right] = \sum_{n \geq 1} \mathcal{L}u_{\lambda,n}$$

can be evaluated term by term and the second sum converges rapidly in  $\mathcal{H}_{0,-1}$ . It is therefore enough to show that if  $g = \sum_{j=1}^n u_{\lambda,j}$ , we can find for any given  $\varepsilon$  a

local function  $g_\epsilon$  such that  $\|\mathcal{L}(g - g_\epsilon)\|_{0,-1} \leq \epsilon$ . If  $u \in \oplus_{j=1}^n \mathcal{H}_j$ , with components  $\{u_j\}$ ,

$$\|\mathcal{L}u\|_{0,-1} \leq \sum_{j=1}^n \|\mathcal{L}u_j\|_{0,-1} \leq C \sum_{j=1}^n j \|u_j\|_1 \leq C_n \|u\|_{0,1}.$$

All we need to do now is to find a local function  $g_\epsilon \in \oplus_{j=1}^n \mathcal{H}_j$ , such that

$$\|g - g_\epsilon\|_{0,1} < \epsilon.$$

Note that  $g$  is in  $L_2(\nu_\alpha)$  and the approximation can be done in each  $\mathcal{H}_j$ .  $\square$

We now prove the main result of the article.

*Proof of Theorem 2.5.* In view of (3.5), which holds with  $\|\cdot\|_{-1,\alpha}$  in place of  $\|\cdot\|_{0,-1,\alpha}$ , to prove Theorem 2.5, it is enough to show that

$$\limsup_{\substack{N \rightarrow \infty \\ K_N/(2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K_N}^2 \leq \|h\|_{-1,\alpha}^2 \quad (4.7)$$

for every local function  $h$  in  $\mathcal{C}_0$ . Notice first that this statement is elementary if  $h = \mathcal{L}h_0$  for some local function  $h_0$ .

Fix  $\epsilon > 0$ . By Theorem 4.2, there exist local functions  $f_\epsilon$  and  $u_\epsilon$  such that  $h = \mathcal{L}f_\epsilon + u_\epsilon$  and  $\|u_\epsilon\|_{0,-1,\alpha} \leq \epsilon$ . Let  $N$  be larger that the supports of  $\mathcal{L}f_\epsilon$  and  $u_\epsilon$ . In this case,

$$\|h\|_{-1,N,K} = \|\mathcal{L}f_\epsilon + u_\epsilon\|_{-1,N,K} \leq \|\mathcal{L}f_\epsilon\|_{-1,N,K} + \|u_\epsilon\|_{-1,N,K}.$$

Since (4.7) is elementary for functions of type  $\mathcal{L}g$ , as  $N \uparrow \infty$  and  $K_N/(2N)^d \rightarrow \alpha$ , the first term on the right hand side converges to  $\|\mathcal{L}f_\epsilon\|_{-1,\alpha}^2$ . By Corollary 2.4, since  $u_\epsilon$  belongs to  $\mathcal{C}_0$ , as  $N \uparrow \infty$  and  $K_N/(2N)^d \rightarrow \alpha$ , the second term converges to  $\|u_\epsilon\|_{0,-1,\alpha}$ , which is at most  $\epsilon$ . Therefore,

$$\limsup_{\substack{N \rightarrow \infty \\ K_N/(2N)^d \rightarrow \alpha}} \|h\|_{-1,N,K_N} \leq \|\mathcal{L}f_\epsilon\|_{-1,\alpha} + \epsilon \leq \|h\|_{-1,\alpha} + 2\epsilon.$$

Since  $\epsilon > 0$  is arbitrary we are done.  $\square$

## 5. APPENDIX

We begin with a lemma.

**Lemma 5.1 .** *There exists a finite constant depending only on the transition probability  $p$  such that*

$$D_\tau(f) \leq C n D_0(f)$$

for all functions  $f$  in  $\mathcal{H}_n$ .

*Proof.* We note that on  $\mathcal{H}_n$ ,  $D_0(f)$  can be replaced by the expression

$$\widehat{D}_0(f) = \frac{1}{2} \sum_{e \in \mathbb{E}_n} \sum_{A \in \mathcal{E}_n} \left\{ f(\sigma^e A) - f(A) \right\}^2$$

because of its equivalence to  $D_0(f)$  which is in fact uniform in  $n$ . Here  $\mathbb{E}_n$ , is the set of edges in  $\mathcal{E}_n$  connecting  $A$  and  $B$  if  $A = \sigma^{x,y}B$ , for some  $x, y \in \mathbb{Z}_*^d$  such that  $y - x \in F$ . Recall that  $F$  can be taken to be the set of points in  $\mathbb{Z}^d$  where every coordinate is bounded by 2 in absolute value.

An easy calculation of

$$\int (1 - \xi(z))[f(\tau_z \xi) - f(\xi)]^2 d\nu_\alpha$$

for  $f \in \mathcal{H}_n$  of the form  $f = \sum_{|A|=n} f(A)\Psi_A$  leads to

$$D_\tau(f) = \frac{1}{2} \sum_z \sum_{|A|=n} p(z)r_z(A)[f(\tau_{-z}A) - f(A)]^2 \quad (5.1)$$

with  $r_z(A) = \alpha$  if  $z \in A$  and  $1 - \alpha$  otherwise. Since  $r_z(A) \leq 1$  and (5.1) is really a finite sum over  $\{z : p(z) > 0\}$ , it is basically the question of estimating

$$\sum_{A \in \mathcal{E}_n} \left( f(\tau_{-z}A) - f(A) \right)^2$$

for each  $z$ . We can assume with out loss of generality that  $z = (1, 0, \dots)$ , the unit vector in the direction of the first coordinate axis.

The problem is now reduced to the following: we are given a function  $g : \mathcal{E}_n \rightarrow \mathbb{R}$ . We think of  $\mathcal{E}_n$  as a graph with edges  $\mathbb{E}_n$ . The Dirichlet form is

$$\widehat{\mathcal{D}}_0(g) = \frac{1}{2} \sum_{e \in \mathbb{E}_n} |(\delta g)(e)|^2$$

and we are claiming an estimate of the form

$$\frac{1}{2} \sum_{A \in \mathcal{E}_n} |g(\tau_z A) - g(A)|^2 \leq C n \widehat{\mathcal{D}}_0(g)$$

with a constant independent of  $n$ . It is clear that for any  $A$  and  $B = \tau_z A$  one can move from  $A$  to  $B$  along the edges of the graph  $\mathcal{E}_n$ , using only the edges in  $\mathbb{E}_n$ . We will verify that we can assign a set of edges  $\mathbb{E}_A \subset \mathbb{E}_n$  such that, (i) for every  $A$  one can use the edges of  $\mathbb{E}_A$  to go from  $A$  to  $\tau_z A$ , (ii) for any  $A$  there are at most  $n$  edges in  $\mathbb{E}_A$  and (iii) the subsets  $\{\mathbb{E}_A\}$  are mutually disjoint as  $A$  varies over  $\mathcal{E}_n$ . Then it is easy to see that

$$|g(\tau_z A) - g(A)|^2 = \left| \sum_{e \in \mathbb{E}_A} (\delta g)(e) \right|^2 \leq n \sum_{e \in \mathbb{E}_A} |(\delta g)(e)|^2$$

and summing over  $A \in \mathcal{E}_n$ , because  $\mathbb{E}_A$  are disjoint, we can establish the Lemma.

To construct the paths from  $A$  to  $\tau_z A$  we totally order the points of  $\mathbb{Z}_*^d$  by lexicographic ordering. We say that  $z = (z_1, \dots, z_d) \in \mathbb{Z}_*^d$  is *positive* if, either  $z_1 > 0$ ; or  $z_1 = 0, z_2 = 0, \dots, z_{j-1} = 0$  and  $z_j > 0$  for some  $j, 2 \leq j \leq d$ . The total ordering declares  $y > x$  if  $y - x$  is *positive*. Let the set  $A \in \mathcal{E}_n$  consist of the  $n$  points  $(x_1, x_2, \dots, x_n)$  of  $\mathbb{Z}_*^d$ . We can assume that they are ordered so that  $x_1 > x_2 > \dots > x_n$ . Then  $\tau_z A = (x_1^*, \dots, x_n^*)$  where  $x_j^* = x_j + z$  unless  $x_j = -z$  in which case  $x_j^* = x_j + 2z = z$ . We use the edges in  $\mathbb{E}_n$  to shift successively each  $x_i$  to  $x_i^*$  starting from  $x_1$  and proceeding in order and ending with shifting  $x_n$ . Any edge that is used goes from some  $A_1$  to  $A_2 = \sigma^{x, x^*} A_1$ . Since the shifts were made in lexicographic order we can determine without ambiguity which points of  $A_1$  have already been shifted and which have not been. In other words the paths from any two different  $A$  to the corresponding  $\tau_y A$  do not share a common edge. It is also clear that exactly  $n$  edges are used. Any edge corresponds to  $x^* - x = z$  or  $2z$  both of which are in  $F$ .  $\square$

We want to look at the resolvent equation

$$\lambda u_\lambda - \mathcal{L}u_\lambda = -h \quad (5.2)$$

and obtain some decay estimates on the contributions to the Dirichlet form from terms of large degree. We follow essentially the method carried out in [LY] or [SVY]. The lattice is  $\mathbb{Z}_*^d$  and we have the Bernoulli measure  $\nu_\alpha$  on  $\mathcal{X} = \{0, 1\}^{\mathbb{Z}_*^d}$ . We consider the orthonormal basis  $\{\Psi_A\}$  introduced in (3.1) and the decomposition of  $L_2(\nu_\alpha)$  as  $\oplus_j \mathcal{H}_j$ . We will consider operators  $\mathbb{T}$  acting on  $L_2(\nu_\alpha)$  as scalar multiplication by  $t(j)$  on  $\mathcal{H}_j$ . Assume that  $t(j)$  is constant for  $j \leq n_1$ . These or the operators that commute with the projection operators  $\pi_j$  onto the subspaces  $\mathcal{H}_j$ . The operator  $\mathcal{L}$  can be written as  $\mathcal{L}_0 + \mathcal{L}_\tau$  the sum of the parts corresponding to the motions of untagged particles and the tagged particle. The operator  $\mathcal{L}_0$  commutes with  $\mathbb{T}$ . Corresponding to the decomposition  $L_2(\nu_\alpha) = \oplus \mathcal{H}_j$  we can write a block decomposition for  $\mathcal{L}_\tau$  involving a tridiagonal block matrix  $\mathfrak{L}_{i,j}$ . If we write  $u = \sum_j u_j$  where  $u_j = \pi_j u$  then we can compute the commutator

$$\begin{aligned} [\mathbb{T}, \mathcal{L}] &= \mathbb{T}\mathcal{L} - \mathcal{L}\mathbb{T} \\ &= \sum_j t(j) [\mathfrak{L}_{j-1,j} u_{j-1} + \mathfrak{L}_{j,j} u_j + \mathfrak{L}_{j+1,j} u_{j+1}] \\ &\quad - \sum_j [t(j-1)\mathfrak{L}_{j-1,j} u_{j-1} + t(j)\mathfrak{L}_{j,j} u_j + t(j+1)\mathfrak{L}_{j+1,j} u_{j+1}] \\ &= \sum_j [t(j) - t(j-1)] \mathfrak{L}_{j-1,j} u_{j-1} + [t(j) - t(j+1)] \mathfrak{L}_{j+1,j} u_{j+1} \end{aligned}$$

We let  $t(j)u_j = v_j$  and  $t(j) - t(j-1) = s(j)$ . From the reversibility of  $\mathcal{L}_\tau$  we can conclude that

$$\mathfrak{L}_{j-1,j}^* = \mathfrak{L}_{j,j-1}$$

We then use it to estimate the following quadratic form

$$\begin{aligned}
& \langle [\mathbb{T}, \mathcal{L}]\mathbb{T}^{-1}\mathbb{T}u, \mathbb{T}u \rangle \\
& \langle [\mathbb{T}, \mathcal{L}]\mathbb{T}^{-1}\mathbb{T}u, \mathbb{T}u \rangle \\
& = \sum_j \langle [t(j) - t(j-1)]\mathfrak{L}_{j-1,j} u_{j-1} + [t(j) - t(j+1)]\mathfrak{L}_{j+1,j} u_{j+1}, t(j)u_j \rangle \\
& = \sum_j \frac{s(j)}{t(j-1)} \langle \mathfrak{L}_{j-1,j} v_{j-1}, v_j \rangle - \sum_j \frac{s(j+1)}{t(j+1)} \langle \mathfrak{L}_{j+1,j} v_{j+1}, v_j \rangle \\
& = \sum_j \frac{s(j)}{t(j-1)} \langle \mathfrak{L}_{j-1,j} v_{j-1}, v_j \rangle - \sum_j \frac{s(j+1)}{t(j+1)} \langle \mathfrak{L}_{j,j+1} v_j, v_{j+1} \rangle \\
& = \sum_j \frac{s(j)^2}{t(j-1)t(j)} \langle \mathfrak{L}_{j-1,j} v_{j-1}, v_j \rangle \\
& = \sum_j \frac{s(j)^2}{t(j-1)t(j)} \langle \mathcal{L}_\tau v_{j-1}, v_j \rangle \\
& \leq \sum_j \frac{s(j)^2}{t(j-1)t(j)} \sqrt{\langle -\mathcal{L}_\tau v_{j-1}, v_{j-1} \rangle} \sqrt{\langle -\mathcal{L}_\tau v_j, v_j \rangle} \\
& \leq \frac{1}{2} \sum_j \frac{s(j)^2}{t(j-1)t(j)} [\langle -\mathcal{L}_\tau v_{j-1}, v_{j-1} \rangle + \langle -\mathcal{L}_\tau v_j, v_j \rangle] .
\end{aligned} \tag{5.3}$$

Since  $-\mathcal{L}_\tau$  is symmetric and positive definite the penultimate inequality is valid. We can now combine the last inequality of our calculation in (5.3) with Lemma 5.1 to arrive at

$$\langle [\mathbb{T}, \mathcal{L}]\mathbb{T}^{-1}\mathbb{T}u, \mathbb{T}u \rangle \leq \sum_j \frac{C}{2} \frac{js(j)^2}{t(j-1)t(j)} [\langle -\mathcal{L}_0 v_{j-1}, v_{j-1} \rangle + \langle -\mathcal{L}_0 v_j, v_j \rangle] .$$

Let us suppose that

$$\sup_j \frac{Cjs(j)^2}{t(j-1)t(j)} \leq \delta .$$

Then

$$\langle [\mathbb{T}, \mathcal{L}]\mathbb{T}^{-1}\mathbb{T}u, \mathbb{T}u \rangle \leq \delta \langle -\mathcal{L}_0 v, v \rangle \leq \delta \langle -\mathcal{L}v, v \rangle = \delta \langle -\mathcal{L}\mathbb{T}u, \mathbb{T}u \rangle . \tag{5.4}$$

We start with the resolvent equation

$$\lambda u_\lambda - \mathcal{L}u_\lambda = -h$$

operate by  $\mathbb{T}$  and take inner product with  $v = \mathbb{T}u_\lambda$ . We get

$$\lambda \langle v, v \rangle - \langle \mathbb{T}\mathcal{L}\mathbb{T}^{-1}v, v \rangle = - \langle h, v \rangle ,$$

which can be rewritten using (5.4) as

$$\begin{aligned} \lambda \langle v, v \rangle - \langle \mathcal{L}v, v \rangle &= \langle [\mathbb{T}\mathcal{L} - \mathcal{L}\mathbb{T}]\mathbb{T}^{-1}v, v \rangle - \langle h, v \rangle \\ &\leq \delta \langle -\mathcal{L}v, v \rangle - \langle h, v \rangle . \end{aligned}$$

This implies the estimate

$$\|v\|_1^2 = \langle -\mathcal{L}v, v \rangle \leq (1 - \delta)^{-1} | \langle h, v \rangle | \leq \|\mathbb{T}h\|_{-1} \|v\|_1 \leq \|\mathbb{T}h\|_{0,-1} \|v\|_1 .$$

In particular

$$\sup_{\lambda > 0} \|\mathbb{T}u_\lambda\|_{0,1}^2 \leq (1 - \delta)^{-1} \|\mathbb{T}h\|_{0,-1} .$$

It is not hard to construct sequences  $t(j)$  that satisfy

$$\sup_j \frac{Cjs(j)^2}{t(j-1)t(j)} \leq \delta$$

and increase rapidly to infinity. We can get up to  $e^{c\sqrt{n}}$  for some small  $c > 0$  that depends on  $C$  and  $\delta$ . This ‘a priori’ estimate can easily be formally established to show that for any local  $h \in \mathcal{H}_{0,-1}$  there is a constant  $C(k, h)$  such that

$$D_0(\pi_n u_\lambda) \leq C(k, h) n^{-k} \tag{5.5}$$

for all  $\lambda > 0$  and  $n \geq 1$ . Note that the mean zero property implies that  $\pi_0(u_\lambda) = 0$ .

#### REFERENCES

- [DFGW] De Masi A., Ferrari P.A., Goldstein S., Wick W.D.; An Invariance principle for reversible Markov processes. Applications to random motions in random environments. *J. Stat. Phys.* **55**, nos.3/4, 787–855 (1989).
- [GOS] Giacomin G., Olla S., Spohn H., Equilibrium fluctuations for  $\nabla\phi$  interface models. Preprint Université de Cergy-Pontoise, CNRS-UPRES-A 8088, n.12/99, (1999). To appear in *Ann.Prob.*
- [G] Grigor’yan A.A.; On Liouville theorems for harmonic functions with finite Dirichlet integral, *Math. USSR Sbornik* **60**, 485-504 (1988).
- [K] Kesten H.; *Percolation Theory for Mathematicians*, Birkhauser (1982).
- [KL] Kipnis C., Landim C.; *Scaling Limit of Interacting Particle Systems*, Grundlehren der mathematischen Wissenschaften **320**, Springer-Verlag, Berlin, New York, (1999).
- [KV] Kipnis C., Varadhan S. R. S.; Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusion. *Commun. Math. Phys.* **106** 1–19 (1986).

- [L] Landim C.; Decay to equilibrium in  $L^\infty$  of asymmetric simple exclusion processes in infinite volume. *Markov Proc. Rel. Fields* **4**, 517–534, (1998).
- [LY] Landim C., Yau H. T.; Fluctuation–dissipation equation of asymmetric simple exclusion processes, *Probab. Th. Rel. Fields* **108**, 321–356, (1997)
- [Li] Liggett T.; *Interacting Particles Systems*, Grundlehren der mathematischen Wissenschaften **276**, Springer-Verlag, Berlin, New York, (1985).
- [OS] Osada H., Saitoh T.; An invariance principle for non-symmetric Markov processes and reflecting diffusions in random domains, *Prob. Th. Relat. Fields* **101**, 45–63 (1995).
- [SVY] Sethuraman S., Varadhan S. R. S., Yau H. T.; Diffusive limit of a tagged particle in asymmetric exclusion process. *Comm.Pure Appl.Math.* **53**, 972–1006, (2000).
- [S] Soardi P.M.; *Potential Theory on Infinite Networks*, Lect. Notes in Math. **1590**, Springer (1994).
- [V1] Varadhan S.R.S.; Self diffusion of a tagged particle in equilibrium for asymmetric mean zero random walks with simple exclusion, *Ann. Inst. H. Poincaré (Probabilités)*, **31**, 273–285.
- [V2] Varadhan S.R.S.; Non-linear diffusion limit for a system with nearest neighbor interactions II. In *Asymptotic Problems in Probability Theory: Stochastic Models and diffusion on Fractals*, Elworthy K.D. and Ikeda N. editors. Pitman Research Notes in Mathematics vol. 283, 75–128, John Wiley & Sons, New York (1994).

IMPA, ESTRADA DONA CASTORINA 110, CEP 22460 RIO DE JANEIRO, BRASIL AND  
CNRS UPRES-A 6085, UNIVERSITÉ DE ROUEN, 76128 MONT SAINT AIGNAN, FRANCE.

*E-mail address:* landim@impa.br

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE CERGY–PONTOISE, 2 AV. ADOLPHE CHAUVIN, B.P.222, PONTOISE 95.302 CERGY–PONTOISE CEDEX, FRANCE AND CENTRE DE MATHÉMATIQUES APPLIQUÉES, ECOLE POLYTECHNIQUE 91128 PALAISEAU CEDEX, FRANCE

*E-mail address:* olla@u-cergy.fr

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, 251 MERCER STREET, NEW YORK, NY, 10012, U.S.A.

*E-mail address:* varadhan@cims.nyu.edu