

Bosonization in a High-Density Fermi Gas

Upper Bound on the Correlation Energy

Niels Benedikter

joint work with

Phan Thành Nam, Marcello Porta, Benjamin Schlein, and Robert Seiringer



Quantum Many-Body Systems

Hamiltonian of N identical spinless particles on $[0, 2\pi]^3$ with periodic boundary conditions

$$H_N := \sum_{i=1}^N (-\Delta_i) + \sum_{1 \leq i < j \leq N} V(x_i - x_j) \quad \text{with } V : \mathbb{R}^3 \rightarrow \mathbb{R}$$

on the bosonic Hilbert space

$$L^2_{\text{symm}}(\mathbb{T}^{3N}) := \left\{ \psi \in L^2(\mathbb{T}^{3N}) \mid \psi(x_{\sigma(1)}, x_{\sigma(2)}, \dots) = \psi(x_1, x_2, \dots) \quad \forall \sigma \in S_N \right\}$$

or on the fermionic Hilbert space

$$L^2_{\text{antisymm}}(\mathbb{T}^{3N}) := \left\{ \psi \in L^2(\mathbb{T}^{3N}) \mid \psi(x_{\sigma(1)}, x_{\sigma(2)}, \dots) = \text{sgn}(\sigma) \psi(x_1, x_2, \dots) \quad \forall \sigma \in S_N \right\} .$$

Ground State Energy

What is the ground state energy

$$E_N := \inf_{\|\psi\|=1} \langle \psi, H_N \psi \rangle = \inf \text{spec}(H_N) ?$$

Defining the two- and one-particle reduced density matrices (r. d. m.)

$$\gamma^{(2)} := \frac{N!}{(N-2)!} \text{tr}_{3,4,\dots,N} |\psi\rangle\langle\psi|, \quad \gamma^{(1)} := \frac{1}{N-1} \text{tr}_2 \gamma^{(2)},$$

we always have

$$\langle \psi, H_N \psi \rangle = \text{tr} \left(-\Delta \gamma^{(1)} \right) + \frac{1}{2} \iint V(x_1 - x_2) \gamma^{(2)}(x_1, x_2; x_1, x_2) dx_1 dx_2.$$

Ground State Energy

What is the ground state energy

$$E_N := \inf_{\|\psi\|=1} \langle \psi, H_N \psi \rangle = \inf \text{spec}(H_N) ?$$

Defining the two- and one-particle reduced density matrices (r. d. m.)

$$\gamma^{(2)} := \frac{N!}{(N-2)!} \text{tr}_{3,4,\dots,N} |\psi\rangle\langle\psi|, \quad \gamma^{(1)} := \frac{1}{N-1} \text{tr}_2 \gamma^{(2)},$$

we always have

$$\langle \psi, H_N \psi \rangle = \text{tr} \left(-\Delta \gamma^{(1)} \right) + \frac{1}{2} \iint V(x_1 - x_2) \gamma^{(2)}(x_1, x_2; x_1, x_2) dx_1 dx_2.$$

So we simply minimize over $\gamma^{(2)}$? Unfortunately not: the set of all two-particle reduced density matrices is hard to characterize: **N-representability problem**.

Bosons

Bosonic Mean-Field Limit

The way out: [restrict to specific physical regimes](#).

Simplest: high density & weak interaction, s. th. we expect approximate mean-field behaviour:

$$H_N^{\text{mf}} = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \text{particle number } N \rightarrow \infty.$$

Bosonic Mean-Field Limit

The way out: **restrict to specific physical regimes.**

Simplest: high density & weak interaction, s. th. we expect approximate mean-field behaviour:

$$H_N^{\text{mf}} = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \text{particle number } N \rightarrow \infty.$$

As $N \rightarrow \infty$, the set of two-particle r. d. m. is characterized by Quantum de-Finetti theorem:

$$\frac{(N-k)!}{N!} \gamma^{(k)} \quad \longrightarrow \quad \int |u^{\otimes k}\rangle \langle u^{\otimes k}| d\mu(u) \quad \text{factorized, no quantum correlations.}$$

Bosonic Mean-Field Limit

The way out: **restrict to specific physical regimes.**

Simplest: high density & weak interaction, s. th. we expect approximate mean-field behaviour:

$$H_N^{\text{mf}} = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \text{particle number } N \rightarrow \infty.$$

As $N \rightarrow \infty$, the set of two-particle r. d. m. is characterized by Quantum de-Finetti theorem:

$$\frac{(N-k)!}{N!} \gamma^{(k)} \longrightarrow \int |u^{\otimes k}\rangle \langle u^{\otimes k}| d\mu(u) \quad \text{factorized, no quantum correlations.}$$

Implies convergence to Hartree functional [Lewin–Nam–Rougerie '13, ...]

$$E_N^{\text{mf}} \rightarrow N \inf_{\substack{u \in L^2(\mathbb{R}^3) \\ \|u\|=1}} \left[\int |\nabla u(x)|^2 dx + \int |u(x)|^2 V(x-y) |u(y)|^2 dx dy \right] =: N E^{\text{Hartree}}.$$

Correlation Corrections to the Hartree Functional

Next order correction: due to quantum correlations!

$$E_N^{\text{mf}} \rightarrow N E^{\text{Hartree}} + \mathcal{O}(1).$$

Correlation Corrections to the Hartree Functional

Next order correction: due to quantum correlations!

$$E_N^{\text{mf}} \rightarrow N E^{\text{Hartree}} + \mathcal{O}(1).$$

Bogoliubov theory [Grech–Seiringer '13, Pizzo '15]:

$$E_N^{\text{mf}} \rightarrow N E^{\text{Hartree}} - \frac{1}{2} \sum_{p \in \mathbb{Z}^3} \left[p^2 + \hat{V}(p) - \sqrt{p^4 + 2p^2 \hat{V}(p)} \right] + \mathcal{O}(N^{-1/2}).$$

Correlation Corrections to the Hartree Functional

Next order correction: due to quantum correlations!

$$E_N^{\text{mf}} \rightarrow N E^{\text{Hartree}} + \mathcal{O}(1).$$

Bogoliubov theory [Grech–Seiringer '13, Pizzo '15]:

$$E_N^{\text{mf}} \rightarrow N E^{\text{Hartree}} - \frac{1}{2} \sum_{p \in \mathbb{Z}^3} \left[p^2 + \hat{V}(p) - \sqrt{p^4 + 2p^2 \hat{V}(p)} \right] + \mathcal{O}(N^{-1/2}).$$

Remark: In thermodynamic limit the correlation energy is given by Lee–Huang–Yang formula

$$E(\rho) \rightarrow 4\pi\rho a \left[1 + \frac{128}{15\sqrt{\pi}} (\rho a^3)^{1/2} + \dots \right], \quad a = \text{scattering length of } V, \quad \rho = \text{density}$$

[Yau–Yin '09], [Giuliani–Seiringer '09], [Boccato–Brennecke–Cenatiempo–Schlein '18],
[Brietzke–Solovej '19], [Brietzke–Fournais–Solovej '19], [Fournais–Solovej '19]

Fermions

Fermionic Mean-Field Regime

Fermions have high kinetic energy (Fermi energy), to be tamed down in mean-field scaling

$$H_N^{\text{mf}} = \sum_{i=1}^N \left(-\hbar^2 \Delta_i \right) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \hbar = N^{-1/3}.$$

Fermionic Mean-Field Regime

Fermions have high kinetic energy (Fermi energy), to be tamed down in mean-field scaling

$$H_N^{\text{mf}} = \sum_{i=1}^N \left(-\hbar^2 \Delta_i \right) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \hbar = N^{-1/3}.$$

No Quantum de-Finetti theorem — set of two-particle r. d. m. is complicated [Klyachko '06].

Fermionic Mean-Field Regime

Fermions have high kinetic energy (Fermi energy), to be tamed down in mean-field scaling

$$H_N^{\text{mf}} = \sum_{i=1}^N \left(-\hbar^2 \Delta_i \right) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \hbar = N^{-1/3}.$$

No Quantum de-Finetti theorem — set of two-particle r. d. m. is complicated [Klyachko '06].

Special correlation estimate implies convergence to Hartree–Fock functional [Graf–Solovej '94]:

$$E_N^{\text{mf}} \rightarrow \inf_{\substack{\omega^2 = \omega \text{ on } L^2(\mathbb{T}^3) \\ \text{tr } \omega = N}} \left[\text{tr}(-\Delta \omega) + \iint \omega(x, x) V(x - y) \omega(y, y) - \iint |\omega(x, y)|^2 V(x - y) \right] =: E_N^{\text{HF}}.$$

Fermionic Mean-Field Regime

Fermions have high kinetic energy (Fermi energy), to be tamed down in mean-field scaling

$$H_N^{\text{mf}} = \sum_{i=1}^N \left(-\hbar^2 \Delta_i \right) + \frac{1}{N} \sum_{1 \leq i < j \leq N} V(x_i - x_j), \quad \hbar = N^{-1/3}.$$

No Quantum de-Finetti theorem — set of two-particle r. d. m. is complicated [Klyachko '06].

Special correlation estimate implies convergence to Hartree–Fock functional [Graf–Solovej '94]:

$$E_N^{\text{mf}} \rightarrow \inf_{\substack{\omega^2 = \omega \text{ on } L^2(\mathbb{T}^3) \\ \text{tr } \omega = N}} \left[\text{tr}(-\Delta \omega) + \iint \omega(x, x) V(x - y) \omega(y, y) - \iint |\omega(x, y)|^2 V(x - y) \right] =: E_N^{\text{HF}}.$$

[Wigner '34]: What is the next order, due to quantum correlations?

The Gell-Mann–Brueckner Formula

Originally jellium model considered: no scaling of couplings, Coulomb interaction, thermodynamic limit, and density $\rho \rightarrow \infty$.

The solution [Macke '50], [Bohm–Pines '53], [Gell-Mann–Brueckner '57], [Sawada–Brueckner–Fukuda–Brout '57] also explained screening and collective plasmon oscillations.

Random Phase Approximation

$$E^{\text{jellium}}(\rho) = \underbrace{C_{\text{TF}}\rho^{5/3} - C_{\text{D}}\rho^{4/3}}_{\text{Hartree-Fock energy of Fermi ball}} + \underbrace{C_{\text{BP}}\rho \log(\rho) + C_{\text{GB}}\rho}_{\text{correlation energy}} + o(\rho) \quad \text{as } \rho \rightarrow \infty.$$

The Gell-Mann–Brueckner Formula

Originally jellium model considered: no scaling of couplings, Coulomb interaction, thermodynamic limit, and density $\rho \rightarrow \infty$.

The solution [Macke '50], [Bohm–Pines '53], [Gell-Mann–Brueckner '57], [Sawada–Brueckner–Fukuda–Brout '57] also explained screening and collective plasmon oscillations.

Random Phase Approximation

$$E^{\text{jellium}}(\rho) = \underbrace{C_{\text{TF}}\rho^{5/3} - C_{\text{D}}\rho^{4/3}}_{\substack{\text{Hartree-Fock energy} \\ \text{of Fermi ball}}} + \underbrace{C_{\text{BP}}\rho \log(\rho) + C_{\text{GB}}\rho}_{\text{correlation energy}} + o(\rho) \quad \text{as } \rho \rightarrow \infty.$$

Mean-field scaling with regular interaction is slightly different:

$$E_N^{\text{mf}} = E_N^{\text{HF}} + \underbrace{E^{\text{BP}} + E^{\text{GB},1}}_{\sim N^{-1/3}} + \underbrace{E^{\text{GB},2}}_{\sim N^{-2/3}}.$$

How did Gell-Mann and Brueckner calculate the correlation energy?

The random phase approximation of Gell-Mann and Brueckner:

- 1 Notice: For Coulomb interaction, high orders are badly IR divergent,

$$\hat{V}(k)^n \sim |k|^{-2n} \quad \text{for } k \rightarrow 0.$$

How did Gell-Mann and Brueckner calculate the correlation energy?

The random phase approximation of Gell-Mann and Brueckner:

- 1 Notice: For Coulomb interaction, high orders are badly IR divergent,

$$\hat{V}(k)^n \sim |k|^{-2n} \quad \text{for } k \rightarrow 0.$$

- 2 Collect the most divergent term from each order of perturbation theory, finding

$$\text{with } x \sim \hat{V}(k): \quad x - \frac{x^2}{2} + \frac{x^3}{3} + \dots =$$

How did Gell-Mann and Brueckner calculate the correlation energy?

The random phase approximation of Gell-Mann and Brueckner:

- 1 Notice: For Coulomb interaction, high orders are badly IR divergent,

$$\hat{V}(k)^n \sim |k|^{-2n} \quad \text{for } k \rightarrow 0.$$

- 2 Collect the most divergent term from each order of perturbation theory, finding

$$\text{with } x \sim \hat{V}(k): \quad x - \frac{x^2}{2} + \frac{x^3}{3} + \dots = \log(1 + x).$$

Resummation of **all orders** of perturbation theory \rightsquigarrow regularization to log-divergence.

How did Gell-Mann and Brueckner calculate the correlation energy?

The random phase approximation of Gell-Mann and Brueckner:

- 1 Notice: For Coulomb interaction, high orders are badly IR divergent,

$$\hat{V}(k)^n \sim |k|^{-2n} \quad \text{for } k \rightarrow 0.$$

- 2 Collect the most divergent term from each order of perturbation theory, finding

$$\text{with } x \sim \hat{V}(k): \quad x - \frac{x^2}{2} + \frac{x^3}{3} + \dots = \log(1 + x).$$

Resummation of **all orders** of perturbation theory \rightsquigarrow regularization to log-divergence.

$$E^{\text{BP}} + E^{\text{GB},1} = \hbar \sum_{k \in \mathbb{Z}^3} |k| \left[\int_0^\infty \log \left(1 + \hat{V}(k) \left(1 - v \arctan v^{-1} \right) \right) dv - \frac{1}{4} \hat{V}(k) \right]$$

How did Gell-Mann and Brueckner calculate the correlation energy?

The random phase approximation of Gell-Mann and Brueckner:

- 1 Notice: For Coulomb interaction, high orders are badly IR divergent,

$$\hat{V}(k)^n \sim |k|^{-2n} \quad \text{for } k \rightarrow 0.$$

- 2 Collect the most divergent term from each order of perturbation theory, finding

$$\text{with } x \sim \hat{V}(k): \quad x - \frac{x^2}{2} + \frac{x^3}{3} + \dots = \log(1 + x).$$

Resummation of **all orders** of perturbation theory \rightsquigarrow regularization to log-divergence.

$$E^{\text{BP}} + E^{\text{GB},1} = \hbar \sum_{k \in \mathbb{Z}^3} |k| \left[\int_0^\infty \log \left(1 + \hat{V}(k) \left(1 - v \arctan v^{-1} \right) \right) dv - \frac{1}{4} \hat{V}(k) \right]$$

Remark: $E^{\text{GB},2}$ is much simpler, just second-order perturbation of exchange type.

Our Result:
Gell-Mann–Brueckner
Formula as Upper Bound

Upper Bound on Correlation Energy

Theorem: [B-Nam-Porta-Schlein-Seiringer, arXiv:1809.01902]

Let $\hat{V}(k)$ be non-negative and compactly supported. Then

$$E_N^{\text{mf}} \leq E_N^{\text{HF}} + E^{\text{BP}} + E^{\text{GB},1} + \mathcal{O}(\hbar N^{-1/27}).$$

Upper Bound on Correlation Energy

Theorem: [B-Nam-Porta-Schlein-Seiringer, arXiv:1809.01902]

Let $\hat{V}(k)$ be non-negative and compactly supported. Then

$$E_N^{\text{mf}} \leq E_N^{\text{HF}} + E^{\text{BP}} + E^{\text{GB},1} + \mathcal{O}(\hbar N^{-1/27}).$$

Remarks:

- [Hainzl-Porta-Rexze '18]: perturbative upper and lower bound to second order in \hat{V} .
- We use a trial state which in principle also captures $E^{\text{GB},2}$, but in the mean-field scaling this contribution is too small to be seen.

Preparation: Extracting the
Hartree–Fock Energy

Extracting the Hartree–Fock Energy

Hamiltonian in momentum representation, written with fermionic canonical operators:

$$H_N^{\text{mf}} := \hbar^2 \sum_{k \in \mathbb{Z}^3} |k|^2 a_k^* a_k + \frac{1}{N} \sum_{q, s, k \in \mathbb{Z}^3} \hat{V}(k) a_{q+k}^* a_{s-k}^* a_s a_q, \quad \hbar = N^{-1/3}$$

Extracting the Hartree–Fock Energy

Hamiltonian in momentum representation, written with fermionic canonical operators:

$$H_N^{\text{mf}} := \hbar^2 \sum_{k \in \mathbb{Z}^3} |k|^2 a_k^* a_k + \frac{1}{N} \sum_{q,s,k \in \mathbb{Z}^3} \hat{V}(k) a_{q+k}^* a_{s-k}^* a_s a_q, \quad \hbar = N^{-1/3}$$

Introduce the Slater determinant of N plane waves in the Fermi ball

$$\Psi_N := \bigwedge_{k \in \mathcal{B}_F} e^{ikx}, \quad \mathcal{B}_F := \left\{ k \in \mathbb{Z}^3 \mid |k| \leq N^{1/3} (3/4\pi)^{1/3} \right\}.$$

[Gontier–Hainzl–Lewin '18]: plane waves are very close to optimal Slater determinant:

$$\langle \Psi_N, H_N^{\text{mf}} \Psi_N \rangle = E_N^{\text{HF}} + \mathcal{O}(e^{-N^{1/6}}).$$

Extracting the Hartree–Fock Energy

Hamiltonian in momentum representation, written with fermionic canonical operators:

$$H_N^{\text{mf}} := \hbar^2 \sum_{k \in \mathbb{Z}^3} |k|^2 a_k^* a_k + \frac{1}{N} \sum_{q, s, k \in \mathbb{Z}^3} \hat{V}(k) a_{q+k}^* a_{s-k}^* a_s a_q, \quad \hbar = N^{-1/3}$$

Introduce the Slater determinant of N plane waves in the Fermi ball

$$\Psi_N := \bigwedge_{k \in \mathcal{B}_F} e^{ikx}, \quad \mathcal{B}_F := \left\{ k \in \mathbb{Z}^3 \mid |k| \leq N^{1/3} (3/4\pi)^{1/3} \right\}.$$

[Gontier–Hainzl–Lewin '18]: plane waves are very close to optimal Slater determinant:

$$\langle \Psi_N, H_N^{\text{mf}} \Psi_N \rangle = E_N^{\text{HF}} + \mathcal{O}(e^{-N^{1/6}}).$$

Goal: find $\tilde{\Psi}_N$ s. th. $\langle \tilde{\Psi}_N, H_N^{\text{mf}} \tilde{\Psi}_N \rangle = E_N^{\text{HF}} + E^{\text{BP}} + E^{\text{GB},1} + o(N^{-1/3})$.

Particle-Hole Transformation

Define the unitary map R on fermionic Fock space by

$$R\Omega := \Psi_N = \bigwedge_{k \in \mathcal{B}_F} e^{ikx}, \quad Ra_k^* R^* := \begin{cases} a_k & k \in \mathcal{B}_F \\ a_k^* & k \in \mathcal{B}_F^c \end{cases}$$

Particle-Hole Transformation

Define the unitary map R on fermionic Fock space by

$$R\Omega := \Psi_N = \bigwedge_{k \in \mathcal{B}_F} e^{ikx}, \quad Ra_k^* R^* := \begin{cases} a_k & k \in \mathcal{B}_F \\ a_k^* & k \in \mathcal{B}_F^c \end{cases}$$

Write $\tilde{\Psi}_N = R\xi$. Calculate $R^* H_N^{\text{mf}} R$ to get

$$\langle \tilde{\Psi}_N, H_N^{\text{mf}} \tilde{\Psi}_N \rangle = E_N^{\text{HF}} + \langle \xi, \underbrace{\left(\hbar^2 \sum_{p \in \mathcal{B}_F^c} p^2 a_p^* a_p - \hbar^2 \sum_{h \in \mathcal{B}_F} h^2 a_h^* a_h + Q \right)}_{=: H^{\text{kin}}} \xi \rangle + \mathcal{O}(N^{-1})$$

where Q is quartic in fermionic operators. Notice: $(H^{\text{kin}} + Q)\Omega = 0$.

Our task: construct a correlated trial state ξ .

Adding Correlations using
Bosonization

Collective Particle-Hole Pairs

The dominant part Q of the interaction can be expressed through collective pair operators

$$b_k^* := \sum_{\substack{p \in \mathcal{B}_F^c \\ h \in \mathcal{B}_F}} \delta_{p-h,k} a_p^* a_h^*$$

as a quadratic Hamiltonian

$$Q = \frac{1}{N} \sum_{k \in \mathbb{Z}^3} \hat{V}(k) (2b_k^* b_k + b_k^* b_{-k}^* + b_{-k} b_k) .$$

Collective Particle-Hole Pairs

The dominant part Q of the interaction can be expressed through collective pair operators

$$b_k^* := \sum_{\substack{p \in \mathcal{B}_F^c \\ h \in \mathcal{B}_F}} \delta_{p-h,k} a_p^* a_h^*$$

as a quadratic Hamiltonian

$$Q = \frac{1}{N} \sum_{k \in \mathbb{Z}^3} \hat{V}(k) (2b_k^* b_k + b_k^* b_{-k}^* + b_{-k} b_k) .$$

Idea: $[a_p^* a_h^*, a_{\tilde{p}}^* a_{\tilde{h}}^*] = 0$, pairs of fermions as bosons? The bad news:

$$(a_p^* a_h^*)^2 = 0 \quad (\text{Pauli principle!}) .$$

Collective Particle-Hole Pairs

The dominant part Q of the interaction can be expressed through collective pair operators

$$b_k^* := \sum_{\substack{p \in \mathcal{B}_F^c \\ h \in \mathcal{B}_F}} \delta_{p-h,k} a_p^* a_h^*$$

as a quadratic Hamiltonian

$$Q = \frac{1}{N} \sum_{k \in \mathbb{Z}^3} \hat{V}(k) (2b_k^* b_k + b_k^* b_{-k}^* + b_{-k} b_k) .$$

Idea: $[a_p^* a_h^*, a_{\tilde{p}}^* a_{\tilde{h}}^*] = 0$, pairs of fermions as bosons? The bad news:


$$(a_p^* a_h^*)^2 = 0 \quad (\text{Pauli principle!}) .$$

The good news: b_k^* are approximately bosonic **if we use the delocalization**:

In $\left(\sum_h a_{h+k}^* a_h^* \right)^2$ only the n diagonal summands out of all n^2 summands vanish.

Linearizing $H^{\text{kin}} = \hbar^2 \sum_p p^2 a_p^* a_p - \hbar^2 \sum_h h^2 a_h^* a_h$

Fermi ball \mathcal{B}_F




Localize to $M = M(N)$ patches near the Fermi surface,

$$b_{\alpha,k}^* := \frac{1}{n_{\alpha,k}} \sum_{\substack{h \in \mathcal{B}_F \cap \mathcal{B}_\alpha \\ p \in \mathcal{B}_F^c \cap \mathcal{B}_\alpha}} \delta_{p-h,k} a_p^* a_h^*$$

where $n_{\alpha,k} = \sqrt{\#\text{p-h pairs in patch } \alpha \text{ with momentum } k}$.

Linearizing $H^{\text{kin}} = \hbar^2 \sum_p p^2 a_p^* a_p - \hbar^2 \sum_h h^2 a_h^* a_h$

Fermi ball \mathcal{B}_F



Localize to $M = M(N)$ patches near the Fermi surface,

$$b_{\alpha,k}^* := \frac{1}{n_{\alpha,k}} \sum_{\substack{h \in \mathcal{B}_F \cap \mathcal{B}_\alpha \\ p \in \mathcal{B}_F^c \cap \mathcal{B}_\alpha}} \delta_{p-h,k} a_p^* a_h^*$$

where $n_{\alpha,k} = \sqrt{\#\text{p-h pairs in patch } \alpha \text{ with momentum } k}$.


Linearize kinetic energy around patch centers ω_α :

$$\begin{aligned} H^{\text{kin}} b_{\alpha,k}^* \Omega &\simeq 2\hbar \underbrace{|k \cdot \hat{\omega}_\alpha|}_{=: u_\alpha(k)^2} b_{\alpha,k}^* \Omega. \\ &=: u_\alpha(k)^2 \end{aligned}$$

By comparison to harmonic oscillator:

$$H^{\text{kin}} \simeq \sum_{k \in \mathbb{Z}^3} \sum_{\alpha} 2\hbar u_\alpha(k)^2 b_{\alpha,k}^* b_{\alpha,k}.$$

Effective Hamiltonian



Normalization constant = number of available modes:

$$n_{\alpha,k}^2 = \# \text{p-h pairs in patch } \alpha \text{ with momentum } k$$

$$\simeq \frac{4\pi N^{2/3}}{M} |k \cdot \hat{\omega}_\alpha| = \frac{4\pi N^{2/3}}{M} u_\alpha(k)^2.$$

In the quadratic Hamiltonian Q , decompose

$$b_k^* = \sum_{\alpha} n_{\alpha,k} b_{\alpha,k}^* + \text{lower order}.$$

$$H^{\text{eff}} = \hbar \sum_{k \in \mathbb{Z}^3} \left[\sum_{\alpha} u_{\alpha}(k)^2 b_{\alpha,k}^* b_{\alpha,k} + \frac{\hat{V}(k)}{M} \sum_{\alpha, \beta} \left(u_{\alpha}(k) u_{\beta}(k) b_{\alpha,k}^* b_{\beta,k} + u_{\alpha}(k) u_{\beta}(-k) b_{\alpha,k}^* b_{\beta,-k}^* + \text{h.c.} \right) \right]$$

Heuristics: Bosonic Approximation

For this slide only: Assume $b_{\alpha,k}^*$, $b_{\alpha,k}$ are exactly bosonic operators.

Then the ground state of H^{eff} is given by a Bogoliubov transformation:

$$\xi_{\text{gs}} = T\Omega, \quad T = \exp \left(\sum_{k \in \mathbb{Z}^3} \sum_{\alpha, \beta} K(k)_{\alpha, \beta} b_{\alpha, k}^* b_{\beta, -k}^* - \text{h.c.} \right) \quad (1)$$

$K(k)$ is an almost explicit $M \times M$ -matrix

and

$$\langle \xi_{\text{gs}}, H^{\text{eff}} \xi_{\text{gs}} \rangle \rightarrow E^{\text{BP}} + E^{\text{GB},1} \quad \text{as } M \rightarrow \infty.$$

To get a rigorous upper bound for the fermionic system:

Use (1) to define a trial state in fermionic Fock space.

Rigorous Analysis

Convergence to Bosonic Approximation

Lemma: We have approximately bosonic commutators:

$$[b_{\alpha,k}^*, b_{\beta,l}^*] = 0 = [b_{\alpha,k}, b_{\beta,l}] \quad \text{and} \quad [b_{\alpha,k}, b_{\beta,l}^*] = \delta_{\alpha,\beta} (\delta_{k,l} + \mathcal{E}_{\alpha}(k,l)),$$

where the operator $\mathcal{E}_{\alpha}(k,l)$ is bounded by

$$\|\mathcal{E}_{\alpha}(k,l)\psi\| \leq \frac{2}{n_{\alpha,k}n_{\alpha,l}} \|\mathcal{N}\psi\| \quad (\mathcal{N} = \text{fermionic number operator})$$

for all ψ in fermionic Fock space.

Approximate Bogoliubov Transformations

Proposition: With $K(k)$ from the bosonic approximation, let in fermionic Fock space

$$T_\lambda := \exp(\lambda B), \quad B := \sum_{k \in \mathbb{Z}^3} \sum_{\alpha, \beta} K(k)_{\alpha, \beta} b_{\alpha, k}^* b_{\beta, -k}^* - \text{h.c.}$$

Then T_λ acts as an approximate Bogoliubov transformation on $b_{\alpha, k}^*$ and $b_{\alpha, k}$, i. e.,

$$T_\lambda^* b_{\alpha, k} T_\lambda = \sum_{\beta=1}^M \cosh(\lambda K(k))_{\alpha, \beta} b_{\beta, k} + \sum_{\beta=1}^M \sinh(\lambda K(k))_{\alpha, \beta} b_{\beta, -k}^* + \mathfrak{E}_{\alpha, k}$$

where the error is bounded by

$$\left[\sum_{\alpha} \|\mathfrak{E}_{\alpha, k} \psi\|^2 \right]^{1/2} \leq \frac{C}{\min_{\alpha} n_{\alpha, k}^2} \|(\mathcal{N} + 2)^{3/2} T_\lambda \psi\| \quad \text{for all } \psi \text{ in fermionic Fock space.}$$

Lemma: The particle number on our trial state

$$\xi := T_{\lambda=1}\Omega$$

is bounded by

$$\langle \xi, (\mathcal{N} + 1)^3 \xi \rangle \leq C \quad \text{independent of } N.$$

Conclusion: We introduce a cutoff excluding patches with $u_\alpha(k)^2 \leq N^{-\delta}$;
thus the error terms are small,

$$\text{errors} \sim \frac{\langle \xi, (\mathcal{N} + 2)^3 \xi \rangle}{\min_\alpha n_{\alpha,k}^2} \leq \frac{C}{\frac{N^{2/3}}{M} u_\alpha(k)^2} \leq C \frac{M}{N^{2/3} N^{-\delta}},$$

\rightsquigarrow bosonic approximation is self-consistent for $M(N) \ll N^{2/3-\delta}$.

Lemma: The kinetic energy can be linearized as $H^{\text{kin}} = H^{\text{linear}} + \mathfrak{E}$, where

$$H^{\text{linear}} = \hbar \sum_{\alpha=1}^M \left[\sum_{p \in \mathcal{B}_F^c \cap B_\alpha} |p \cdot \hat{\omega}_\alpha| a_p^* a_p - \sum_{h \in \mathcal{B}_F \cap B_\alpha} |h \cdot \hat{\omega}_\alpha| a_h^* a_h \right]$$

and the error operator \mathfrak{E} is small compared to $\hbar = N^{-1/3}$ if $M(N) \gg N^{1/3}$; namely

$$|\langle \psi, \mathfrak{E} \psi \rangle| \leq \frac{C}{M} \langle \psi, \mathcal{N} \psi \rangle \quad \text{for all } \psi \text{ in fermionic Fock space.}$$

Lemma: The kinetic energy can be linearized as $H^{\text{kin}} = H^{\text{linear}} + \mathfrak{E}$, where

$$H^{\text{linear}} = \hbar \sum_{\alpha=1}^M \left[\sum_{p \in \mathcal{B}_F^c \cap B_\alpha} |p \cdot \hat{\omega}_\alpha| a_p^* a_p - \sum_{h \in \mathcal{B}_F \cap B_\alpha} |h \cdot \hat{\omega}_\alpha| a_h^* a_h \right]$$

and the error operator \mathfrak{E} is small compared to $\hbar = N^{-1/3}$ if $M(N) \gg N^{1/3}$; namely

$$|\langle \psi, \mathfrak{E} \psi \rangle| \leq \frac{C}{M} \langle \psi, \mathcal{N} \psi \rangle \quad \text{for all } \psi \text{ in fermionic Fock space.}$$

Lemma: We have

$$[H^{\text{linear}}, b_{\alpha,k}^*] = 2\hbar |k \cdot \hat{\omega}_\alpha| b_{\alpha,k}^* ;$$

in this (and only this) sense $H^{\text{linear}} \simeq \sum_{k \in \mathbb{Z}^3} \sum_{\alpha} 2\hbar u_\alpha(k)^2 b_{\alpha,k}^* b_{\alpha,k}$.

Proof of Main Theorem

Proof: We just have to calculate $\langle R\xi, H_N^{\text{mf}} R\xi \rangle \simeq \langle \Omega, T_{\lambda=1}^* (H^{\text{linear}} + Q) T_{\lambda=1} \Omega \rangle$.

- The interaction part Q is quadratic in b^* and b ;
just calculate the action of the approximate Bogoliubov transformation.

Proof of Main Theorem

Proof: We just have to calculate $\langle R\xi, H_N^{\text{mf}} R\xi \rangle \simeq \langle \Omega, T_{\lambda=1}^* (H^{\text{linear}} + Q) T_{\lambda=1} \Omega \rangle$.

- The interaction part Q is quadratic in b^* and b ;
just calculate the action of the approximate Bogoliubov transformation.
- The linearized kinetic energy H^{linear} is **not** quadratic in b^* and b ;
expand into commutators by applying once the Duhamel formula



$$\begin{aligned} \langle \xi, H^{\text{linear}} \xi \rangle &= \int_0^1 \langle \Omega, T_\lambda^* [H^{\text{linear}}, B] T_\lambda \Omega \rangle d\lambda \\ &= \int_0^1 \langle \Omega, T_\lambda^* \sum_{k \in \mathbb{Z}^3} \sum_{\alpha, \beta} K(k)_{\alpha, \beta} [H^{\text{linear}}, b_{\alpha, k}^* b_{\beta, -k}^* - \text{h.c.}] T_\lambda \Omega \rangle d\lambda \\ &= \int_0^1 \langle \Omega, T_\lambda^* \sum_{k \in \mathbb{Z}^3} \sum_{\alpha, \beta} K(k)_{\alpha, \beta} 2\hbar (|k \cdot \hat{\omega}_\alpha| + |k \cdot \hat{\omega}_\beta|) b_{\alpha, k}^* b_{\beta, -k}^* T_\lambda \Omega \rangle + \text{c. c.} \end{aligned}$$

and $T_\lambda^* b_{\alpha, k}^* T_\lambda$ is given by the approximate Bogoliubov transformation.

QED



Work in Progress

- Corresponding lower bound – gapless system!
- Coulomb interaction and the plasmon:



Work in Progress

- Corresponding lower bound – gapless system!
- Coulomb interaction and the plasmon:



Thank you!