

ISF App 28th January 2021

Exam : Introduction to Machine Learning¹ : Time 1h30

Exercise 1. [10pts] Select the correct answers

1. By assumption, the $(X^i, Y^i)_{i \in \llbracket 1, n \rrbracket}$ in the learning sample, in supervised learning are such :
 - a) the (X^i, Y^i) have all the same laws and are independent
 - b) the (X^i, Y^i) have the same laws but may be dependent
 - c) the X^i have all a normal distribution

2. if $(X^i)_{i \in \llbracket 1, n \rrbracket}$ is a learning sample without any labelling.
 - a) we are in the framework of supervised learning
 - b) we are in the framework of unsupervised learning
 - c) generally we try to derive a structure for the X^i

3. if $R_n(f_n)$ is the classification error and $R(f_n)$ is the prediction error then
 - a) $R(f_n) < R_n(f_n)$
 - b) $E(R(f_n)) \geq R_n(f_n)$
 - c) $R(f_n) \geq E[R_n(f_n)]$
 - d) $E(R(f_n)) \leq R_n(f_n)$

4. in Vapnik Chervonenkis's formula
 - a) $P\left(R(f_n) \leq R_n(f_n) + \phi_{n,\eta}\left(\frac{VC(\mathcal{F})}{n}\right)\right) \geq 1 - \eta$
 - b) $P\left(R_n(f_n) \leq R(f_n) + \phi_{n,\eta}\left(\frac{VC(\mathcal{F})}{n}\right)\right) \geq 1 - \eta$
 - c) $P\left(R_n(f_n) \leq R(f_n) + \phi_{n,\eta}\left(\frac{VC(\mathcal{F})}{n}\right)\right) \geq 1 - \frac{\eta}{n}$
 - a) $P\left(R(f_n) \leq R_n(f_n) + \phi_{n,\eta}\left(\frac{VC(\mathcal{F})}{n}\right)\right) \geq 1 - \frac{\eta}{n}$

5. in the Vapnik Chervonenkis's formula
 - a) $\phi_{n,\eta}(t)$ is a function which is decreasing with t
 - b) $\phi_{n,\eta}(t)$ is a function which is increasing with t
 - c) $\phi_{n,\eta}\left(\frac{1}{n}\right)$ tends to zero when n tends to ∞

6. in Structural Risk Minimisation with $\mathcal{F}_1 \subset \mathcal{F}_2 \cdots \subset \mathcal{F}_k \cdots$ the classifier $f_{n,k}$ from class \mathcal{F}_k is chosen if
 - a) it minimizes over k , $R_n(f_{n,k}) + \phi_{n,\eta}\left(\frac{VC(\mathcal{F}_k)}{n}\right)$
 - b) it minimizes over k , $R_n(f_{n,k})$
 - c) it minimizes over k , $R_n(f_{n,k}) - \phi_{n,\eta}\left(\frac{VC(\mathcal{F}_k)}{n}\right)$

1. Pierre Brugière University Paris Dauphine PSL

7. if x_1, x_2, \dots, x_n is a family of independent vectors then
- x_1, x_2, \dots, x_n can be classified in all possible ways with hyperplanes
 - 0 can be separated from x_1, x_2, \dots, x_n with a hyperplane
 - 0 can be separated from the convex envelope of x_1, x_2, \dots, x_n with a hyperplane
8. if a family of classifiers \mathcal{F} is defined by d parameters $d \geq 1$ then
- $VC(\mathcal{F}) < d + 2$
 - $VC(\mathcal{F}) = d + 1$
 - there may be some cases for which $VC(\mathcal{F}) = +\infty$
9. what is the distance between the point represented by the vector z and the hyperplane of equation $\langle w, x \rangle = b$
- $\frac{|b + \langle w, z \rangle|}{\|w\|^2}$
 - $\frac{|\langle w, z \rangle - b|}{\|w\|}$
 - $\frac{|\langle w, z \rangle - b|}{\|w\|^2}$
10. what is the distance between the two hyperplanes of equations :
- $\langle w, x \rangle = b$ and $\langle -w, x \rangle + c = 0$
- $\frac{|b - c|}{\|w\|^2}$
 - $\frac{|b + c|}{\|w\|}$
 - $\frac{|b - c|}{\|w\|}$
11. in \mathbf{R}^d
- the VC dimension of hyperplane classifiers of margin 1 is $d + 1$
 - the VC dimension of hyperplane classifiers within a ball of radius 1 is $d + 1$
 - the VC dimension of hyperplane classifiers of radius 1 and margin 0.1 is less than $d + 1$ when d is large
12. in a C-SVM
- all the support vectors are always on the borders of the separating hyperplanes
 - it is possible that some support vectors are not correctly classified
 - it is impossible that some support vectors lay between the two separating hyperplanes
13. which of these inequalities is correct :
- $\sup_{z \in \mathcal{Z}} \left[\inf_{y \in \mathcal{Y}} g(y, z) \right] \leq \inf_{y \in \mathcal{Y}} \left[\sup_{z \in \mathcal{Z}} g(y, z) \right]$
 - $\sup_{z \in \mathcal{Z}} \left[\inf_{y \in \mathcal{Y}} g(y, z) \right] \geq \inf_{y \in \mathcal{Y}} \left[\sup_{z \in \mathcal{Z}} g(y, z) \right]$

14. which of the following assertions are true :
- a) if the KKT conditions are satisfied the primal and dual problems have the same value
 - b) if the primal and dual problems have the same value the KKT conditions are satisfied
 - c) the KKT conditions enable to calculate some of the parameters of the separating hyperplanes
15. which of the following assertions are true for the Gaussian Kernel K_σ and associated transformation ϕ_σ :
- a) all the transformed points $\phi_\sigma(x_i)$ are on sphere or radius 1
 - b) for any sample $\{(x_i, y_i)\}_{i \in \llbracket 1, n \rrbracket}$ it is possible to find σ such that the classification with the Gaussian Kernel K_σ is perfect
 - c) a notion of margin can be associated to Gaussian Kernel classifiers
16. single class SVMs can be used
- a) to estimate the support of a distribution
 - b) to define different clusters amongst a sample
 - c) to solve a regression problem
17. if $K(\cdot, \cdot)$ is a kernel with values in \mathbf{R} and $(H, \langle \cdot, \cdot \rangle_{RK})$ is a Reproducing Kernel Hilbert Space for K then
- a) for all $f \in H$, $\langle K(x, \cdot), f \rangle_{RK} = f(x)$
 - b) $K(x_1, x_2)^2 \leq K(x_1, x_1)K(x_2, x_2)$
 - c) for all $f \in H$, $f(x)^2 \leq K(x, x)\langle f, f \rangle_{RK}$
18. in the discount factor curve construction problem studied in the course
- a) the regularisation term is linked to a scalar product on the function space
 - b) the regularisation term transforms a non parametric problem into a parametric problem
 - c) without the regularisation term the problem would not be well defined
19. in the discount factor curve construction problem studied in the course
- a) the spline functions are of degrees 2
 - b) the spline functions have knots on the cash flow dates of the instruments used as inputs
 - c) the coefficients of the spline functions are the solutions of a ridge regression problem
 - d) the coefficients of the spline functions are the solutions of a Lasso regression problem
20. in the Smith and Nelson model the regularisation term is of the form
- a) $\int f''^2(s)ds$
 - b) $\int f''^2(s) + \sigma^2 f'^2(s)ds$

c) $\int f''(s) + \sigma^2 f'(s) ds$

Problem [14pts]

Let $f(x, \lambda)$ be a function of (x, λ) . (x^*, λ^*) is called a saddle point of f iff

$$\forall(x, \lambda), f(x^*, \lambda) \leq f(x^*, \lambda^*) \leq f(x, \lambda^*)$$

1. [2pts] show that $\sup_{\lambda} \inf_x f(x, \lambda) \leq \inf_x \sup_{\lambda} f(x, \lambda)$
2. [2pts] show that if a saddle point (x^*, λ^*) exists then

$$\sup_{\lambda} \inf_x f(x, \lambda) = \inf_x \sup_{\lambda} f(x, \lambda)$$

3. [2pts] using 2., show that if a saddle point (x^*, λ^*) exists then

$$f(x^*, \lambda^*) = \inf_x \sup_{\lambda} f(x, \lambda) = \inf_x f(x, \lambda^*) = \sup_{\lambda} f(x^*, \lambda)$$

In this second part which is mostly independent from the first part, we

consider the problem $\begin{cases} \min_{x \in \mathbf{R}^n} \|x\|^2 \\ Ax = p \end{cases}$

where A is full row rank, and (as a consequence) $p \in \text{Im}(A)$

4. [1pt] write the Lagrangian $\mathcal{L}(x, \lambda)$
5. [1pt] for λ fixed, find $x(\lambda)$, as a function of A and λ which satisfies

$$\mathcal{L}(x(\lambda), \lambda) = \inf_x \mathcal{L}(x, \lambda)$$

6. [1pt] calculate $\mathcal{L}(x(\lambda), \lambda)$ as a function of A , λ and p
7. [2pts] find λ^* , as a function of A and λ which satisfies

$$\mathcal{L}(x(\lambda^*), \lambda^*) = \sup_{\lambda} \mathcal{L}(x(\lambda), \lambda)$$

8. [1pt] verify that $Ax(\lambda^*) = p$
9. [1pt] show that

$$\mathcal{L}(x(\lambda^*), \lambda^*) = \sup_{\lambda} \mathcal{L}(x(\lambda^*), \lambda)$$

10. [1pt] show that $(x(\lambda^*), \lambda^*)$ is a saddle point for \mathcal{L} .