

Champs magnétiques, interpolation et symétrie

Jean Dolbeault

<http://www.ceremade.dauphine.fr/~dolbeaul>

Ceremade, Université Paris-Dauphine

March 29, 2019

Séminaire de Mathématiques Appliquées
Collège de France

Outline

🟢 *Without magnetic fields: symmetry and symmetry breaking in interpolation inequalities*

- ▷ Gagliardo-Nirenberg-Sobolev inequalities on the sphere
- ▷ Keller-Lieb-Thirring inequalities on the sphere
- ▷ Caffarelli-Kohn-Nirenberg inequalities

🟢 *With magnetic fields in dimensions 2 and 3*

- ▷ Interpolation inequalities and spectral estimates
- ▷ Estimates, numerics; an open question on constant magnetic fields

🟢 *Magnetic rings: the case of \mathbb{S}^1*

- ▷ A one-dimensional magnetic interpolation inequality
- ▷ Consequences: Keller-Lieb-Thirring estimates, Aharonov-Bohm magnetic fields and a new Hardy inequality in \mathbb{R}^2

🟢 *Aharonov-Bohm magnetic fields in \mathbb{R}^2*

- ▷ Aharonov-Bohm effect
- ▷ Interpolation and Keller-Lieb-Thirring inequalities in \mathbb{R}^2
- ▷ Aharonov-Symmetry and symmetry breaking

A joint research program (mostly) with...


M.J. Esteban, Ceremade, Université Paris-Dauphine

▷ *symmetry, interpolation, Keller-Lieb-Thirring, magnetic fields*



M. Loss, Georgia Institute of Technology (Atlanta)

▷ *symmetry, interpolation, Keller-Lieb-Thirring, magnetic fields*



A. Laptev, Imperial College London

▷ *Keller-Lieb-Thirring, magnetic fields*



D. Bonheure, Université Libre de Bruxelles

▷ *Aharonov-Bohm magnetic fields*



Symmetry and symmetry breaking in interpolation inequalities *without magnetic field*

- Gagliardo-Nirenberg-Sobolev inequalities on the sphere
- Keller-Lieb-Thirring inequalities on the sphere
- Caffarelli-Kohn-Nirenberg inequalities on \mathbb{R}^2

A result of uniqueness on a classical example

On the sphere \mathbb{S}^d , let us consider the positive solutions of

$$-\Delta u + \lambda u = u^{p-1}$$

$$p \in [1, 2) \cup (2, 2^*] \text{ if } d \geq 3, \quad 2^* = \frac{2d}{d-2}$$

$$p \in [1, 2) \cup (2, +\infty) \text{ if } d = 1, 2$$

Theorem

If $\lambda \leq d$, $u \equiv \lambda^{1/(p-2)}$ is the unique solution

[Gidas & Spruck, 1981], [Bidaut-Véron & Véron, 1991]

Bifurcation point of view and symmetry breaking




Figure: $(p-2)\lambda \mapsto (p-2)\mu(\lambda)$ with $d=3$

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu(\lambda) \|u\|_{L^p(\mathbb{S}^d)}^2$$

Taylor expansion of $u = 1 + \varepsilon \varphi_1$ as $\varepsilon \rightarrow 0$ with $-\Delta \varphi_1 = d \varphi_1$

$$\mu(\lambda) < \lambda \quad \text{if and only if} \quad \lambda > \frac{d}{p-2}$$

▷ The inequality holds with $\mu(\lambda) = \lambda = \frac{d}{p-2}$ [Bakry & Emery, 1985]

[Beckner, 1993], [Bidaut-Véron & Véron, 1991, Corollary 6.1]

The Bakry-Emery method on the sphere

Entropy functional

$$\mathcal{E}_p[\rho] := \frac{1}{p-2} \left[\int_{\mathbb{S}^d} \rho^{\frac{2}{p}} d\mu - \left(\int_{\mathbb{S}^d} \rho d\mu \right)^{\frac{2}{p}} \right] \quad \text{if } p \neq 2$$

$$\mathcal{E}_2[\rho] := \int_{\mathbb{S}^d} \rho \log \left(\frac{\rho}{\|\rho\|_{L^1(\mathbb{S}^d)}} \right) d\mu$$

Fisher information functional

$$\mathcal{I}_p[\rho] := \int_{\mathbb{S}^d} |\nabla \rho^{\frac{1}{p}}|^2 d\mu$$

[Bakry & Emery, 1985] *carré du champ* method: use the heat flow

$$\frac{\partial \rho}{\partial t} = \Delta \rho$$

and observe that $\frac{d}{dt} \mathcal{E}_p[\rho] = -\mathcal{I}_p[\rho]$

$$\frac{d}{dt} \left(\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho] \right) \leq 0 \quad \implies \quad \mathcal{I}_p[\rho] \geq d \mathcal{E}_p[\rho]$$

with $\rho = |u|^p$, if $p \leq 2^\# := \frac{2d^2+1}{(d-1)^2}$


The evolution under the fast diffusion flow

To overcome the limitation $p \leq 2^\#$, one can consider a nonlinear diffusion of fast diffusion / porous medium type

$$\frac{\partial \rho}{\partial t} = \Delta \rho^m$$

[Demange], [JD, Esteban, Kowalczyk, Loss]: for any $p \in [1, 2^*]$

$$\mathcal{K}_p[\rho] := \frac{d}{dt} \left(\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho] \right) \leq 0$$



(p, m) admissible region, $d = 5$

References

JD, M. J. Esteban, M. Kowalczyk, and M. Loss. Improved interpolation inequalities on the sphere. Discrete and Continuous Dynamical Systems Series S, 7 (4): 695-724, 2014.

JD, M.J. Esteban, and M. Loss. Interpolation inequalities on the sphere: linear vs. nonlinear flows. Annales de la faculté des sciences de Toulouse Sér. 6, 26 (2): 351-379, 2017

Keller-Lieb-Thirring inequalities on the sphere

- The Keller-Lieb-Thirring inequality is equivalent to an interpolation inequality of Gagliardo-Nirenberg-Sobolev type
- We measure a quantitative deviation with respect to the semi-classical regime due to finite size effects

Joint work with M.J. Esteban and A. Laptev

An introduction to (Keller)-Lieb-Thirring inequalities in \mathbb{R}^d

$(\lambda_k)_{k \geq 1}$: eigenvalues of the Schrödinger operator $H = -\Delta - V$ on \mathbb{R}^d

• Euclidean case [Keller, 1961]

$$|\lambda_1|^\gamma \leq L_{\gamma,d}^1 \int_{\mathbb{R}^d} V_+^{\gamma + \frac{d}{2}}$$

[Lieb-Thirring, 1976]

$$\sum_{k \geq 1} |\lambda_k|^\gamma \leq L_{\gamma,d} \int_{\mathbb{R}^d} V_+^{\gamma + \frac{d}{2}}$$

$\gamma \geq 1/2$ if $d = 1$, $\gamma > 0$ if $d = 2$ and $\gamma \geq 0$ if $d \geq 3$ [Weidl], [Cwikel], [Rosenbljum], [Aizenman], [Laptev-Weidl], [Helffer], [Robert], [JD-Felmer-Loss-Paturel], [JD-Laptev-Loss]... [Frank, Hundertmark, Jex, Nam]

• Compact manifolds: log Sobolev case: [Federbusch], [Rothaus];
 case $\gamma = 0$ (Rozenbljum-Lieb-Cwikel inequality): [Levin-Solomyak];
 [Lieb], [Levin], [Ouabaz-Poupaud]... [Ilyin]



▷ How does one take into account the finite size effects on \mathbb{S}^d ?

Hölder duality and link with interpolation inequalities

Let $p = \frac{q}{q-2}$. Consider the Schrödinger energy

$$\begin{aligned} \int_{\mathbb{S}^d} |\nabla u|^2 - \int_{\mathbb{S}^d} V |u|^2 &\geq \int_{\mathbb{S}^d} |\nabla u|^2 - \mu \|u\|_{L^q(\mathbb{S}^d)}^2 \\ &\geq -\lambda(\mu) \|u\|_{L^2(\mathbb{S}^d)}^2 \quad \text{if } \mu = \|V_+\|_{L^p(\mathbb{S}^d)} \end{aligned}$$

● We deduce from $\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu(\lambda) \|u\|_{L^q(\mathbb{S}^d)}^2$ that
 $\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 - \mu(\lambda) \|u\|_{L^q(\mathbb{S}^d)}^2 \geq -\lambda \|u\|_{L^2(\mathbb{S}^d)}^2$



A Keller-Lieb-Thirring inequality on the sphere

Let $d \geq 1$, $p \in [\max\{1, d/2\}, +\infty)$ and $\mu_* := \frac{d}{2}(p-1)$

Theorem (JD-Esteban-Laptev)

There exists a convex increasing function λ s.t. $\lambda(\mu) = \mu$ if $\mu \in [0, \mu_]$ and $\lambda(\mu) > \mu$ if $\mu \in (\mu_*, +\infty)$ and, for any $p < d/2$,*

$$|\lambda_1(-\Delta - V)| \leq \lambda(\|V\|_{L^p(\mathbb{S}^d)}) \quad \forall V \in L^p(\mathbb{S}^d)$$

This estimate is optimal

For large values of μ , we have

$$\lambda(\mu)^{p-\frac{d}{2}} = L_{p-\frac{d}{2}, d}^1(\kappa_{q,d} \mu)^p (1 + o(1))$$

If $p = d/2$ and $d \geq 3$, the inequality holds with $\lambda(\mu) = \mu$ iff $\mu \in [0, \mu_]$*

A Keller-Lieb-Thirring inequality: second formulation

Let $d \geq 1$, $\gamma = p - d/2$

Corollary (JD-Esteban-Laptev)

$$|\lambda_1(-\Delta - V)|^\gamma \lesssim L_{\gamma,d}^1 \int_{\mathbb{S}^d} V^{\gamma+\frac{d}{2}} \quad \text{as } \mu = \|V\|_{L^{\gamma+\frac{d}{2}}(\mathbb{S}^d)} \rightarrow \infty$$

if either $\gamma > \max\{0, 1 - d/2\}$ or $\gamma = 1/2$ and $d = 1$

However, if $\mu = \|V\|_{L^{\gamma+\frac{d}{2}}(\mathbb{S}^d)} \leq \mu_*$, then we have

$$|\lambda_1(-\Delta - V)|^{\gamma+\frac{d}{2}} \leq \int_{\mathbb{S}^d} V^{\gamma+\frac{d}{2}}$$

for any $\gamma \geq \max\{0, 1 - d/2\}$ and this estimate is optimal

$L_{\gamma,d}^1$ is the optimal constant in the Euclidean one bound state ineq.

$$|\lambda_1(-\Delta - \phi)|^\gamma \leq L_{\gamma,d}^1 \int_{\mathbb{R}^d} \phi_+^{\gamma+\frac{d}{2}} dx$$

References

- JD, M. J. Esteban, and A. Laptev. Spectral estimates on the sphere. *Analysis & PDE*, 7 (2): 435-460, 2014
- JD, M.J. Esteban, A. Laptev, M. Loss. Spectral properties of Schrödinger operators on compact manifolds: Rigidity, flows, interpolation and spectral estimates. *Comptes Rendus Mathématique*, 351 (11-12): 437-440, 2013

Caffarelli-Kohn-Nirenberg, symmetry and symmetry breaking results, and weighted nonlinear flows

Joint work with M.J. Esteban and M. Loss

Critical Caffarelli-Kohn-Nirenberg inequality

Let $\mathcal{D}_{a,b} := \left\{ v \in L^p(\mathbb{R}^d, |x|^{-b} dx) : |x|^{-a} |\nabla v| \in L^2(\mathbb{R}^d, dx) \right\}$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx \quad \forall v \in \mathcal{D}_{a,b}$$

holds under conditions on a and b

$$p = \frac{2d}{d-2+2(b-a)} \quad (\text{critical case})$$

▷ *An optimal function among radial functions:*


$$v_\star(x) = \left(1 + |x|^{(p-2)(a_c-a)} \right)^{-\frac{2}{p-2}} \quad \text{and} \quad C_{a,b}^\star = \frac{\| |x|^{-b} v_\star \|_p^2}{\| |x|^{-a} \nabla v_\star \|_2^2}$$

Question: $C_{a,b} = C_{a,b}^\star$ (symmetry) or $C_{a,b} > C_{a,b}^\star$ (symmetry breaking) ?

Critical CKN: range of the parameters

Figure: $d = 3$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx$$



$$p = \frac{2d}{d-2+2(b-a)}$$

$$a < a_c := (d-2)/2$$

$$a \leq b \leq a+1 \text{ if } d \geq 3,$$

$$a + 1/2 < b \leq a+1 \text{ if } d = 1$$

$$\text{and } a < b \leq a+1 < 1,$$

$$a < b \leq a+1 < 1 \text{ if } d = 2$$

[Il'in (1961)]

[Glaser, Martin, Grosse, Thirring (1976)]


[Caffarelli, Kohn, Nirenberg (1984)]

[F. Catrina, Z.-Q. Wang (2001)]

Linear instability of radial minimizers: the Felli-Schneider curve

The Felli & Schneider curve

$$b_{\text{FS}}(a) := \frac{d(a_c - a)}{2\sqrt{(a_c - a)^2 + d - 1}} + a - a_c$$



[Smets], [Smets, Willem], [Catrina, Wang], [Felli, Schneider]


The functional

$$C_{a,b}^* \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx - \left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p}$$

is linearly instable at $v = v_*$.

Symmetry *versus* symmetry breaking: the sharp result in the critical case

[JD, Esteban, Loss (2016)]



Theorem

Let $d \geq 2$ and $p < 2^*$. If either $a \in [0, a_c)$ and $b > 0$, or $a < 0$ and $b \geq b_{\text{FS}}(a)$, then the optimal functions for the critical Caffarelli-Kohn-Nirenberg inequalities are radially symmetric

The symmetry proof in one slide

• A change of variables: $v(|x|^{\alpha-1}x) = w(x)$, $D_\alpha v = \left(\alpha \frac{\partial v}{\partial s}, \frac{1}{s} \nabla_\omega v\right)$

$$\|v\|_{L^{2p,d-n}(\mathbb{R}^d)} \leq K_{\alpha,n,p} \|D_\alpha v\|_{L^{2,d-n}(\mathbb{R}^d)}^\vartheta \|v\|_{L^{p+1,d-n}(\mathbb{R}^d)}^{1-\vartheta} \quad \forall v \in H_{d-n,d-n}^p(\mathbb{R}^d)$$

• Concavity of the Rényi entropy power: with

$$\mathcal{L}_\alpha = -D_\alpha^* D_\alpha = \alpha^2 \left(u'' + \frac{n-1}{s} u'\right) + \frac{1}{s^2} \Delta_\omega u \quad \text{and} \quad \frac{\partial u}{\partial t} = \mathcal{L}_\alpha u^m$$

$$\begin{aligned} & - \frac{d}{dt} \mathcal{G}[u(t, \cdot)] \left(\int_{\mathbb{R}^d} u^m d\mu \right)^{1-\sigma} \\ & \geq (1-m)(\sigma-1) \int_{\mathbb{R}^d} u^m \left| \mathcal{L}_\alpha P - \frac{\int_{\mathbb{R}^d} u |D_\alpha P|^2 d\mu}{\int_{\mathbb{R}^d} u^m d\mu} \right|^2 d\mu \\ & + 2 \int_{\mathbb{R}^d} \left(\alpha^4 \left(1 - \frac{1}{n}\right) \left| P'' - \frac{P'}{s} - \frac{\Delta_\omega P}{\alpha^2 (n-1) s^2} \right|^2 + \frac{2\alpha^2}{s^2} \left| \nabla_\omega P' - \frac{\nabla_\omega P}{s} \right|^2 \right) u^m d\mu \\ & + 2 \int_{\mathbb{R}^d} \left((n-2) (\alpha_{\text{FS}}^2 - \alpha^2) |\nabla_\omega P|^2 + c(n, m, d) \frac{|\nabla_\omega P|^4}{P^2} \right) u^m d\mu \end{aligned}$$

• Elliptic regularity and the Emden-Fowler transformation: justifying the integrations by parts

The variational problem on the cylinder

▷ *With the Emden-Fowler transformation*

$$v(r, \omega) = r^{a-a_c} \varphi(s, \omega) \quad \text{with} \quad r = |x|, \quad s = -\log r \quad \text{and} \quad \omega = \frac{x}{r}$$

the variational problem becomes

$$\Lambda \mapsto \mu(\Lambda) := \min_{\varphi \in H^1(C)} \frac{\|\partial_s \varphi\|_{L^2(C)}^2 + \|\nabla_\omega \varphi\|_{L^2(C)}^2 + \Lambda \|\varphi\|_{L^2(C)}^2}{\|\varphi\|_{L^p(C)}^2}$$

is a concave increasing function


Restricted to symmetric functions, the variational problem becomes

$$\mu_\star(\Lambda) := \min_{\varphi \in H^1(\mathbb{R})} \frac{\|\partial_s \varphi\|_{L^2(\mathbb{R}^d)}^2 + \Lambda \|\varphi\|_{L^2(\mathbb{R}^d)}^2}{\|\varphi\|_{L^p(\mathbb{R}^d)}^2} = \mu_\star(1) \Lambda^\alpha$$

Symmetry means $\mu(\Lambda) = \mu_\star(\Lambda)$

Symmetry breaking means $\mu(\Lambda) < \mu_\star(\Lambda)$

Numerical results



Parametric plot of the branch of optimal functions for $p = 2.8$, $d = 5$. Non-symmetric solutions bifurcate from symmetric ones at a bifurcation point Λ_1 computed by V. Felli and M. Schneider. The branch behaves for large values of Λ as shown by F. Catrina and Z.-Q. Wang

Three references

📄 Lecture notes on *Symmetry and nonlinear diffusion flows...*
a course on entropy methods (see webpage)

📄 [JD, Maria J. Esteban, and Michael Loss] *Symmetry and symmetry breaking: rigidity and flows in elliptic PDEs*
... the elliptic point of view: Proc. Int. Cong. of Math., Rio de Janeiro, 3: 2279-2304, 2018.

📄 [JD, Maria J. Esteban, and Michael Loss] *Interpolation inequalities, nonlinear flows, boundary terms, optimality and linearization...* the parabolic point of view
Journal of elliptic and parabolic equations, 2: 267-295, 2016.

Magnetic interpolation inequalities in the Euclidean space

- ▷ Three interpolation inequalities and their dual forms
- ▷ Estimates in dimension $d = 2$ for constant magnetic fields
 - Lower estimates
 - Upper estimates and numerical results
 - A linear stability result (numerical) and an open question
- 🟢 Warning: assumptions are not repeated
- 🟢 Estimates are given only in the case $p > 2$ but similar estimates hold in the other cases

Joint work with M.J. Esteban, A. Laptev and M. Loss

Magnetic interpolation inequalities

$$\|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}^2 + \alpha \|\psi\|_{L^2(\mathbb{R}^d)}^2 \geq \mu_{\mathbf{B}}(\alpha) \|\psi\|_{L^p(\mathbb{R}^d)}^2 \quad \forall \psi \in H_{\mathbf{A}}^1(\mathbb{R}^d)$$

for any $\alpha \in (-\Lambda[\mathbf{B}], +\infty)$ and any $p \in (2, 2^*)$,

$$\|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}^2 + \beta \|\psi\|_{L^p(\mathbb{R}^d)}^2 \geq \nu_{\mathbf{B}}(\beta) \|\psi\|_{L^2(\mathbb{R}^d)}^2 \quad \forall \psi \in H_{\mathbf{A}}^1(\mathbb{R}^d)$$

for any $\beta \in (0, +\infty)$ and any $p \in (1, 2)$

$$\|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}^2 \geq \gamma \int_{\mathbb{R}^d} |\psi|^2 \log \left(\frac{|\psi|^2}{\|\psi\|_{L^2(\mathbb{R}^d)}^2} \right) dx + \xi_{\mathbf{B}}(\gamma) \|\psi\|_{L^2(\mathbb{R}^d)}^2$$

(limit case corresponding to $p = 2$) for any $\gamma \in (0, +\infty)$

$$C_p := \begin{cases} \min_{u \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla u\|_{L^2(\mathbb{R}^d)}^2 + \|u\|_{L^2(\mathbb{R}^d)}^2}{\|u\|_{L^p(\mathbb{R}^d)}^2} & \text{if } p \in (2, 2^*) \\ \min_{u \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla u\|_{L^2(\mathbb{R}^d)}^2 + \|u\|_{L^p(\mathbb{R}^d)}^2}{\|u\|_{L^2(\mathbb{R}^d)}^2} & \text{if } p \in (1, 2) \end{cases}$$

$$\mu_0(1) = C_p \text{ if } p \in (2, 2^*), \quad \nu_0(1) = C_p \text{ if } p \in (1, 2)$$

$$\xi_0(\gamma) = \gamma \log(\pi e^2/\gamma) \text{ if } p = 2$$

Technical assumptions

$\mathbf{A} \in L_{\text{loc}}^{\alpha}(\mathbb{R}^d)$, $\alpha > 2$ if $d = 2$ or $\alpha = 3$ if $d = 3$ and

$$\lim_{\sigma \rightarrow +\infty} \sigma^{d-2} \int_{\mathbb{R}^d} |\mathbf{A}(x)|^2 e^{-\sigma|x|} dx = 0 \quad \text{if } p \in (2, 2^*)$$

$$\lim_{\sigma \rightarrow +\infty} \frac{\sigma^{\frac{d}{2}-1}}{\log \sigma} \int_{\mathbb{R}^d} |\mathbf{A}(x)|^2 e^{-\sigma|x|^2} dx = 0 \quad \text{if } p = 2$$

$$\lim_{\sigma \rightarrow +\infty} \sigma^{d-2} \int_{|x| < 1/\sigma} |\mathbf{A}(x)|^2 dx \quad \text{if } p \in (1, 2)$$

These estimates can be found in [Esteban, Lions, 1989]

A statement

Theorem

$p \in (2, 2^*)$: $\mu_{\mathbf{B}}$ is monotone increasing on $(-\Lambda[\mathbf{B}], +\infty)$, concave and

$$\lim_{\alpha \rightarrow (-\Lambda[\mathbf{B}])_+} \mu_{\mathbf{B}}(\alpha) = 0 \quad \text{and} \quad \lim_{\alpha \rightarrow +\infty} \mu_{\mathbf{B}}(\alpha) \alpha^{\frac{d-2}{2} - \frac{d}{p}} = C_p$$

$p \in (1, 2)$: $\nu_{\mathbf{B}}$ is monotone increasing on $(0, +\infty)$, concave and

$$\lim_{\beta \rightarrow 0_+} \nu_{\mathbf{B}}(\beta) = \Lambda[\mathbf{B}] \quad \text{and} \quad \lim_{\beta \rightarrow +\infty} \nu_{\mathbf{B}}(\beta) \beta^{-\frac{2p}{2p+d(2-p)}} = C_p$$

$\xi_{\mathbf{B}}$ is continuous on $(0, +\infty)$, concave, $\xi_{\mathbf{B}}(0) = \Lambda[\mathbf{B}]$ and

$$\xi_{\mathbf{B}}(\gamma) = \frac{d}{2} \gamma \log \left(\frac{\pi e^2}{\gamma} \right) (1 + o(1)) \quad \text{as} \quad \gamma \rightarrow +\infty$$

Constant magnetic fields: equality is achieved

Nonconstant magnetic fields: only partial answers are known




Figure: Case $d = 2$, $p = 3$, $B = 1$: plot of $\alpha \mapsto (2\pi)^{\frac{2}{p}-1} \mu_B(\alpha)$




Figure: Case $d = 2$, $p = 1.4$, $B = 1$: plot of $\beta \mapsto \nu_B(\beta)$

The horizontal axis is measured in units of $(2\pi)^{1-\frac{2}{p}} \beta$

Magnetic Keller-Lieb-Thirring inequalities

$\lambda_{\mathbf{A},V}$ is the principal eigenvalue of $-\Delta_{\mathbf{A}} + V$

$\alpha_{\mathbf{B}} : (0, +\infty) \rightarrow (-\Lambda, +\infty)$ is the inverse function of $\alpha \mapsto \mu_{\mathbf{B}}(\alpha)$

Corollary

(i) For any $q = p/(p-2) \in (d/2, +\infty)$ and any potential $V \in L^q_+(\mathbb{R}^d)$

$$\lambda_{\mathbf{A},V} \geq -\alpha_{\mathbf{B}}(\|V\|_{L^q(\mathbb{R}^d)})$$

$$\lim_{\mu \rightarrow 0_+} \alpha_{\mathbf{B}}(\mu) = \Lambda \text{ and } \lim_{\mu \rightarrow +\infty} \alpha_{\mathbf{B}}(\mu) \mu^{\frac{2(q+1)}{d-2-2q}} = -C_p^{\frac{2(q+1)}{d-2-2q}}$$

(ii) For any $q = p/(2-p) \in (1, +\infty)$ and any $0 < W^{-1} \in L^q(\mathbb{R}^d)$

$$\lambda_{\mathbf{A},W} \geq \nu_{\mathbf{B}}\left(\|W^{-1}\|_{L^q(\mathbb{R}^d)}^{-1}\right)$$

(iii) For any $\gamma > 0$ and any $W \geq 0$ s.t. $e^{-W/\gamma} \in L^1(\mathbb{R}^d)$

$$\lambda_{\mathbf{A},W} \geq \xi_{\mathbf{B}}(\gamma) - \gamma \log \left(\int_{\mathbb{R}^d} e^{-W/\gamma} dx \right)$$

Symmetry in non-magnetic interpolation inequalities

Magnetic interpolation in the Euclidean space

Magnetic rings: the one-dimensional periodic case

Symmetry in Aharonov-Bohm magnetic fields

Three interpolation inequalities and their dual forms

Proofs for general magnetic fields

Estimates in dimension $d = 2$ for constant magnetic fields

Numerical results and the symmetry issue

Proofs

Interpolation without magnetic field...

Assume that $p > 2$ and let C_p denote the best constant in

$$\|\nabla u\|_{L^2(\mathbb{R}^d)}^2 + \|u\|_{L^2(\mathbb{R}^d)}^2 \geq C_p \|u\|_{L^p(\mathbb{R}^d)}^2 \quad \forall u \in H^1(\mathbb{R}^d)$$

By scaling, if we test the inequality by $u(\cdot/\lambda)$, we find that

$$\|\nabla u\|_{L^2(\mathbb{R}^d)}^2 + \lambda^2 \|u\|_{L^2(\mathbb{R}^d)}^2 \geq C_p \lambda^{2-d(1-\frac{2}{p})} \|u\|_{L^p(\mathbb{R}^d)}^2 \quad \forall u \in H^1(\mathbb{R}^d) \quad \forall \lambda > 0$$

An optimization on $\lambda > 0$ shows that the best constant in the scale-invariant inequality

$$\|\nabla u\|_{L^2(\mathbb{R}^d)}^{d(1-\frac{2}{p})} \|u\|_{L^2(\mathbb{R}^d)}^{2-d(1-\frac{2}{p})} \geq S_p \|u\|_{L^p(\mathbb{R}^d)}^2 \quad \forall u \in H^1(\mathbb{R}^d)$$

is given by

$$S_p = \frac{1}{2^p} (2p - d(p-2))^{1-d\frac{p-2}{2p}} (d(p-2))^{\frac{d(p-2)}{2p}} C_p$$

... and with magnetic field

Proposition

Let $d = 2$ or 3 . For any $p \in (2, +\infty)$, any $\alpha > -\Lambda = -\Lambda[\mathbf{B}] < 0$

$$\mu_{\mathbf{B}}(\alpha) \geq \mu_{\text{interp}}(\alpha) := \begin{cases} S_p(\alpha + \Lambda) \Lambda^{-d \frac{p-2}{2p}} & \text{if } \alpha \in \left[-\Lambda, \frac{\Lambda(2p-d(p-2))}{d(p-2)} \right] \\ C_p \alpha^{1-d \frac{p-2}{2p}} & \text{if } \alpha \geq \frac{\Lambda(2p-d(p-2))}{d(p-2)} \end{cases}$$

Diamagnetic inequality: $\|\nabla|\psi|\|_{L^2(\mathbb{R}^d)} \leq \|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}$

Non-magnetic inequality with $\lambda = \frac{\alpha + \Lambda t}{1-t}$, $t \in [0, 1]$

$$\begin{aligned} \|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}^2 + \alpha \|\psi\|_{L^2(\mathbb{R}^d)}^2 &\geq t \left(\|\nabla_{\mathbf{A}}\psi\|_{L^2(\mathbb{R}^d)}^2 - \Lambda \|\psi\|_{L^2(\mathbb{R}^d)}^2 \right) \\ &\quad + (1-t) \left(\|\nabla|\psi|\|_{L^2(\mathbb{R}^d)}^2 + \frac{\alpha + \Lambda t}{1-t} \|\psi\|_{L^2(\mathbb{R}^d)}^2 \right) \\ &\geq C_p (1-t)^{\frac{d(p-2)}{2p}} (\alpha + t\Lambda)^{1-d \frac{p-2}{2p}} \|\psi\|_{L^p(\mathbb{R}^d)}^2 \end{aligned}$$

and optimize on $t \in [\max\{0, -\alpha/\Lambda\}, 1]$

The special case of constant magnetic field in dimension $d = 2$

Constant magnetic field, $d = 2$...

Assume that $\mathbf{B} = (0, B)$ is constant, $d = 2$ and choose

$$\mathbf{A}_1 = \frac{B}{2} x_2, \quad \mathbf{A}_2 = -\frac{B}{2} x_1 \quad \forall x = (x_1, x_2) \in \mathbb{R}^2$$

Proposition

[Loss, Thaller, 1997] Consider a constant magnetic field with field strength B in two dimensions. For every $c \in [0, 1]$, we have

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq (1 - c^2) \int_{\mathbb{R}^2} |\nabla \psi|^2 dx + c B \int_{\mathbb{R}^2} \psi^2 dx$$

and equality holds with $\psi = u e^{iS}$ and $u > 0$ if and only if

$$(-\partial_2 u^2, \partial_1 u^2) = \frac{2u^2}{c} (\mathbf{A} + \nabla S)$$

... a computation ($d = 2$, constant magnetic field)

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx &= \int_{\mathbb{R}^2} |\nabla u|^2 dx + \int_{\mathbb{R}^2} |\mathbf{A} + \nabla S|^2 u^2 dx \\ &= (1 - c^2) \int_{\mathbb{R}^2} |\nabla u|^2 dx + \underbrace{\int_{\mathbb{R}^2} (c^2 |\nabla u|^2 + |\mathbf{A} + \nabla S|^2 u^2) dx}_{\geq \int_{\mathbb{R}^2} 2c |\nabla u| |\mathbf{A} + \nabla S| u dx} \end{aligned}$$

with equality only if $c |\nabla u| = |\mathbf{A} + \nabla S| u$

$$2 |\nabla u| |\mathbf{A} + \nabla S| u = |\nabla u^2| |\mathbf{A} + \nabla S| \geq (\nabla u^2)^\perp \cdot (\mathbf{A} + \nabla S)$$

$$\text{where } (\nabla u^2)^\perp := (-\partial_2 u^2, \partial_1 u^2)$$

Equality case: $(-\partial_2 u^2, \partial_1 u^2) = \gamma (\mathbf{A} + \nabla S)$ for $\gamma = 2u^2/c$

Integration by parts yields

$$\int_{\mathbb{R}^2} (c^2 |\nabla u|^2 + |\mathbf{A} + \nabla S|^2 u^2) dx \geq B c \int_{\mathbb{R}^2} u^2 dx$$

... a lower estimate ($d = 2$, constant magnetic field)

Proposition

Consider a constant magnetic field with field strength B in two dimensions. Given any $p \in (2, +\infty)$, and any $\alpha > -B$, we have

$$\mu_{\mathbf{B}}(\alpha) \geq C_p (1 - c^2)^{1 - \frac{2}{p}} (\alpha + c B)^{\frac{2}{p}} =: \mu_{\text{LT}}(\alpha)$$

with

$$c = c(p, \eta) = \frac{\sqrt{\eta^2 + p - 1} - \eta}{p - 1} = \frac{1}{\eta + \sqrt{\eta^2 + p - 1}} \in (0, 1)$$

and $\eta = \alpha(p - 2)/(2B)$

Upper estimate (1): $d = 2$, constant magnetic field

For every integer $k \in \mathbb{N}$ we introduce the special symmetry class

$$\psi(x) = \left(\frac{x_2 + i x_1}{|x|} \right)^k v(|x|) \quad \forall x = (x_1, x_2) \in \mathbb{R}^2 \quad (\mathcal{C}_k)$$

[Esteban, Lions, 1989]: if $\psi \in \mathcal{C}_k$, then

$$\frac{1}{2\pi} \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx = \int_0^{+\infty} |v'|^2 r dr + \int_0^{+\infty} \left(\frac{k}{r} - \frac{Br}{2} \right)^2 |v|^2 r dr$$

and optimality is achieved in \mathcal{C}_k

Test function $v_\sigma(r) = e^{-r^2/(2\sigma)}$: an optimization on $\sigma > 0$ provides an explicit expression of $\mu_{\text{Gauss}}(\alpha)$ such that

Proposition

If $p > 2$, then

$$\mu_{\mathbf{B}}(\alpha) \leq \mu_{\text{Gauss}}(\alpha) \quad \forall \alpha > -\Lambda[\mathbf{B}]$$

This estimate is not optimal because v_σ does not solve the Euler-Lagrange equations

Upper estimate (2): $d = 2$, constant magnetic field

A more numerical point of view. The Euler-Lagrange equation in \mathcal{C}_0 is

$$-v'' - \frac{v'}{r} + \left(\frac{B^2}{4} r^2 + \alpha \right) v = \mu_{\text{EL}}(\alpha) \left(\int_0^{+\infty} |v|^p r \, dr \right)^{\frac{2}{p}-1} |v|^{p-2} v$$

We can restrict the problem to positive solutions such that

$$\mu_{\text{EL}}(\alpha) = \left(\int_0^{+\infty} |v|^p r \, dr \right)^{1-\frac{2}{p}}$$

and then we have to solve the reduced problem

$$-v'' - \frac{v'}{r} + \left(\frac{B^2}{4} r^2 + \alpha \right) v = |v|^{p-2} v$$

Numerical results and the symmetry issue




Figure: Case $d = 2$, $p = 3$, $B = 1$

Upper estimates: $\alpha \mapsto \mu_{\text{Gauss}}(\alpha)$, $\mu_{\text{EL}}(\alpha)$

Lower estimates: $\alpha \mapsto \mu_{\text{interp}}(\alpha)$, $\mu_{\text{LT}}(\alpha)$

The exact value associated with μ_B lies in the grey area.

Plots represent the curves $\log_{10}(\mu/\mu_{\text{EL}})$

Asymptotics (1): *Lowest Landau Level*

Proposition

Let $d = 2$ and consider a constant magnetic field with field strength B . If ψ_α is a minimizer for $\mu_B(\alpha)$ such that $\|\psi_\alpha\|_{L^p(\mathbb{R}^d)} = 1$, then there exists a non trivial $\varphi_\alpha \in \text{LLL}$ such that

$$\lim_{\alpha \rightarrow (-B)_+} \|\psi_\alpha - \varphi_\alpha\|_{H_A^1(\mathbb{R}^2)} = 0$$

Let $\psi_\alpha \in H_A^1(\mathbb{R}^2)$ be an optimal function such that $\|\psi_\alpha\|_{L^p(\mathbb{R}^d)} = 1$ and let us decompose it as $\psi_\alpha = \varphi_\alpha + \chi_\alpha$, where $\varphi_\alpha \in \text{LLL}$ and χ_α is in the orthogonal of LLL

$$\mu_B(\alpha) \geq (\alpha+B) \|\varphi_\alpha\|_{L^2(\mathbb{R}^d)}^2 + (\alpha+3B) \|\chi_\alpha\|_{L^2(\mathbb{R}^d)}^2 \geq (\alpha+3B) \|\chi_\alpha\|_{L^2(\mathbb{R}^d)}^2 \sim 2B \|\chi_\alpha\|_{L^2(\mathbb{R}^d)}^2$$

as $\alpha \rightarrow (-B)_+$ because $\|\nabla \chi_\alpha\|_{L^2(\mathbb{R}^d)}^2 \geq 3B \|\chi_\alpha\|_{L^2(\mathbb{R}^d)}^2$

Since $\lim_{\alpha \rightarrow (-B)_+} \mu_B(\alpha) = 0$, $\lim_{\alpha \rightarrow (-B)_+} \|\chi_\alpha\|_{L^2(\mathbb{R}^d)} = 0$ and

$$\mu_B(\alpha) = (\alpha+B) \|\varphi_\alpha\|_{L^2(\mathbb{R}^d)}^2 + \|\nabla_A \chi_\alpha\|_{L^2(\mathbb{R}^d)}^2 + \alpha \|\chi_\alpha\|_{L^2(\mathbb{R}^d)}^2 \geq \frac{2}{3} \|\nabla_A \chi_\alpha\|_{L^2(\mathbb{R}^d)}^2$$

concludes the proof

Asymptotics (2): *semi-classical regime*

Let us consider the small magnetic field regime. We assume that the magnetic potential is given by

$$\mathbf{A}_1 = \frac{B}{2}x_2, \quad \mathbf{A}_2 = -\frac{B}{2}x_1 \quad \forall x = (x_1, x_2) \in \mathbb{R}^2$$

if $d = 2$. In dimension $d = 3$, we choose $\mathbf{A} = \frac{B}{2}(-x_2, x_1, 0)$ and observe that the constant magnetic field is $\mathbf{B} = (0, 0, B)$, while the spectral gap is $\Lambda[\mathbf{B}] = B$.

Proposition

Let $d = 2$ or 3 and consider a constant magnetic field \mathbf{B} of intensity B with magnetic potential \mathbf{A}

For any $p \in (2, 2^)$ and any fixed α and $\mu > 0$, we have*

$$\lim_{\varepsilon \rightarrow 0_+} \mu_\varepsilon \mathbf{B}(\alpha) = C_p \alpha^{\frac{d}{p} - \frac{d-2}{2}}$$

Consider any function $\psi \in H_{\mathbf{A}}^1(\mathbb{R}^d)$ and let $\psi(x) = \chi(\sqrt{\varepsilon}x)$, $\sqrt{\varepsilon} \mathbf{A}(x/\sqrt{\varepsilon}) = \mathbf{A}(x)$ with our conventions on \mathbf{A}

Numerical stability of radial optimal functions

Let us denote by ψ_0 an optimal function in (\mathcal{C}_0) such that

$$-\psi_0'' - \frac{\psi_0'}{r} + \left(\frac{B^2}{4} r^2 + \alpha\right) \psi_0 = |\psi_0|^{p-2} \psi_0$$

and consider the test function

$$\psi_\varepsilon = \psi_0 + \varepsilon e^{i\theta} v$$

where $v = v(r)$ and $e^{i\theta} = (x_1 + i x_2)/r$

As $\varepsilon \rightarrow 0_+$, the leading order term is

$$2\pi \left[\int_{\mathbb{R}^2} |v'|^2 dx + \int_{\mathbb{R}^2} \left(\left(\frac{1}{r} - \frac{Br}{2} \right)^2 + \alpha \right) |v|^2 dx - \frac{p}{2} \int_0^{+\infty} |\psi_0|^{p-2} v^2 r dr \right] \varepsilon^2$$

and we have to solve the eigenvalue problem

$$-v'' - \frac{v'}{r} + \left(\left(\frac{1}{r} - \frac{Br}{2} \right)^2 + \alpha \right) v - \frac{p}{2} |\psi_0|^{p-2} v = \mu v$$




Figure: Case $p = 3$ and $B = 1$: plot of the eigenvalue μ as a function of α . A careful investigation shows that μ is always positive, including in the limiting case as $\alpha \rightarrow (-B)_+$, thus proving the numerical stability of the optimal function in \mathcal{C}_0 with respect to perturbations in \mathcal{C}_1 .

An open question of symmetry

🟢 [Bonheure, Nys, Van Schaftingen, 2016] for a fixed $\alpha > 0$ and for \mathbf{B} small enough, the optimal functions are radially symmetric functions, *i.e.*, belong to \mathcal{C}_0


This regime is equivalent to the regime as $\alpha \rightarrow +\infty$ for a given \mathbf{B} , at least if the magnetic field is constant

🟢 Numerically our upper and lower bounds are (in dimension $d = 2$, for a constant magnetic field) numerically extremely close

🟢 The optimal function in \mathcal{C}_0 is linearly stable with respect to perturbations in \mathcal{C}_1

▷ *Prove that the optimality case is achieved among radial function if $d = 2$ and \mathbf{B} is a constant magnetic field*

Reference

 JD, M.J. Esteban, A. Laptev, M. Loss. Interpolation inequalities and spectral estimates for magnetic operators. Annales Henri Poincaré, 19 (5): 1439-1463, May 2018

Magnetic rings

▷ A magnetic interpolation inequality on \mathbb{S}^1 : with $p > 2$

$$\|\psi' + i a \psi\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|\psi\|_{L^2(\mathbb{S}^1)}^2 \geq \mu_{a,p}(\alpha) \|\psi\|_{L^p(\mathbb{S}^1)}^2$$

▷ Consequences

- A Keller-Lieb-Thirring inequality
- A new Hardy inequality for Aharonov-Bohm magnetic fields in \mathbb{R}^2

Joint work with M.J. Esteban, A. Laptev and M. Loss

Magnetic flux, a reduction

Assume that $a : \mathbb{R} \rightarrow \mathbb{R}$ is a 2π -periodic function such that its restriction to $(-\pi, \pi] \approx \mathbb{S}^1$ is in $L^1(\mathbb{S}^1)$ and define the space

$$X_a := \{ \psi \in C_{\text{per}}(\mathbb{R}) : \psi' + i a \psi \in L^2(\mathbb{S}^1) \}$$

🟢 A standard change of gauge (see e.g. [Ilyin, Laptev, Loss, Zelik, 2016])

$$\psi(s) \mapsto e^{i \int_{-\pi}^s (a(\sigma) - \bar{a}) d\sigma} \psi(s)$$

where $\bar{a} := \int_{-\pi}^{\pi} a(s) d\sigma$ is the *magnetic flux*, reduces the problem to

a is a constant function

🟢 For any $k \in \mathbb{Z}$, ψ by $s \mapsto e^{iks} \psi(s)$ shows that $\mu_{a,p}(\alpha) = \mu_{k+a,p}(\alpha)$

$$a \in [0, 1]$$

🟢 $\mu_{a,p}(\alpha) = \mu_{1-a,p}(\alpha)$ because

$$|\psi' + i a \psi|^2 = |\chi' + i(1-a)\chi|^2 = |\bar{\psi}' - i a \bar{\psi}|^2 \text{ if } \chi(s) = e^{-is} \overline{\psi(s)}$$

$$a \in [0, 1/2]$$

Optimal interpolation

We want to characterize the *optimal constant* in the inequality

$$\|\psi' + i a \psi\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|\psi\|_{L^2(\mathbb{S}^1)}^2 \geq \mu_{a,p}(\alpha) \|\psi\|_{L^p(\mathbb{S}^1)}^2$$

written for any $p > 2$, $a \in (0, 1/2]$, $\alpha \in (-a^2, +\infty)$, $\psi \in X_a$

$$\mu_{a,p}(\alpha) := \inf_{\psi \in X_a \setminus \{0\}} \frac{\int_{-\pi}^{\pi} (|\psi' + i a \psi|^2 + \alpha |\psi|^2) d\sigma}{\|\psi\|_{L^p(\mathbb{S}^1)}^2}$$

$p = -2 = 2d/(d-2)$ with $d = 1$ [Exner, Harrell, Loss, 1998]

$p = +\infty$ [Galunov, Olienik, 1995] [Ilyin, Laptev, Loss, Zelik, 2016]

$\lim_{\alpha \rightarrow -a^2} \mu_{a,p}(\alpha) = 0$ [JD, Esteban, Laptev, Loss, 2016]

Using a Fourier series $\psi(s) = \sum_{k \in \mathbb{Z}} \psi_k e^{iks}$, we obtain that

$$\|\psi' + i a \psi\|_{L^2(\mathbb{S}^1)}^2 = \sum_{k \in \mathbb{Z}} (a + k)^2 |\psi_k|^2 \geq a^2 \|\psi\|_{L^2(\mathbb{S}^1)}^2$$

$\psi \mapsto \|\psi' + i a \psi\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|\psi\|_{L^2(\mathbb{S}^1)}^2$ is coercive for any $\alpha > -a^2$

An interpolation result for the magnetic ring

Theorem

For any $p > 2$, $a \in \mathbb{R}$, and $\alpha > -a^2$, $\mu_{a,p}(\alpha)$ is achieved and
 (i) if $a \in [0, 1/2]$ and $a^2(p+2) + \alpha(p-2) \leq 1$, then $\mu_{a,p}(\alpha) = a^2 + \alpha$
 and equality is achieved only by the constant functions
 (ii) if $a \in [0, 1/2]$ and $a^2(p+2) + \alpha(p-2) > 1$, then $\mu_{a,p}(\alpha) < a^2 + \alpha$
 and equality is not achieved by the constant functions
 If $\alpha > -a^2$, $a \mapsto \mu_{a,p}(\alpha)$ is monotone increasing on $(0, 1/2)$




Figure: $\alpha \mapsto \mu_{a,p}(\alpha)$ with $p = 4$ and (left) $a = 0.45$ or (right) $a = 0.2$

The proof: how to eliminate the phase

Reformulations of the interpolation problem (1/3)

Any minimizer $\psi \in X_a$ of $\mu_{a,p}(\alpha)$ satisfies the Euler-Lagrange equation

$$(H_a + \alpha) \psi = |\psi|^{p-2} \psi, \quad H_a \psi = - \left(\frac{d}{ds} + i a \right)^2 \psi \quad (*)$$

up to a multiplication by a constant and $v(s) = \psi(s) e^{ias}$ satisfies the condition

$$v(s + 2\pi) = e^{2i\pi a} v(s) \quad \forall s \in \mathbb{R}$$

Hence

$$\mu_{a,p}(\alpha) = \min_{v \in Y_a \setminus \{0\}} Q_{p,\alpha}[v]$$

where $Y_a := \{v \in C(\mathbb{R}) : v' \in L^2(\mathbb{S}^1), (*) \text{ holds}\}$ and

$$Q_{p,\alpha}[v] := \frac{\|v'\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|v\|_{L^2(\mathbb{S}^1)}^2}{\|v\|_{L^p(\mathbb{S}^1)}^2}$$

Reformulations of the interpolation problem (2/3)

With $v = u e^{i\phi}$ the boundary condition becomes

$$u(\pi) = u(-\pi), \quad \phi(\pi) = 2\pi(a + k) + \phi(-\pi) \quad (**)$$

for some $k \in \mathbb{Z}$, and $\|v'\|_{L^2(\mathbb{S}^1)}^2 = \|u'\|_{L^2(\mathbb{S}^1)}^2 + \|u\phi'\|_{L^2(\mathbb{S}^1)}^2$

Hence

$$\mu_{a,p}(\alpha) = \min_{(u,\phi) \in Z_a \setminus \{0\}} \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + \|u\phi'\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^1)}^2}{\|u\|_{L^p(\mathbb{S}^1)}^2}$$

where $Z_a := \{(u, \phi) \in C(\mathbb{R})^2 : u', u\phi' \in L^2(\mathbb{S}^1), (**) \text{ holds}\}$

Reformulations of the interpolation problem (3/3)

We use the Euler-Lagrange equations

$$-u'' + |\phi'|^2 u + \alpha u = |u|^{p-2} u \quad \text{and} \quad (\phi' u^2)' = 0$$

Integrating the second equation, and *assuming that u never vanishes*, we find a constant L such that $\phi' = L/u^2$. Taking $(*)$ into account, we deduce from

$$L \int_{-\pi}^{\pi} \frac{ds}{u^2} = \int_{-\pi}^{\pi} \phi' ds = 2\pi (a + k)$$

that

$$\|u \phi'\|_{L^2(\mathbb{S}^1)}^2 = L^2 \int_{-\pi}^{\pi} \frac{d\sigma}{u^2} = \frac{(a + k)^2}{\|u^{-1}\|_{L^2(\mathbb{S}^1)}^2}$$

Hence

$$\phi(s) - \phi(0) = \frac{a + k}{\|u^{-1}\|_{L^2(\mathbb{S}^1)}^2} \int_{-\pi}^s \frac{ds}{u^2}$$

Let us define

$$\mathcal{Q}_{a,p,\alpha}[u] := \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \alpha \|u\|_{L^2(\mathbb{S}^1)}^2}{\|u\|_{L^p(\mathbb{S}^1)}^2}$$

Lemma

For any $a \in (0, 1/2)$, $p > 2$, $\alpha > -a^2$,

$$\mu_{a,p}(\alpha) = \min_{u \in H^1(\mathbb{S}^1) \setminus \{0\}} \mathcal{Q}_{a,p,\alpha}[u]$$

is achieved by a function $u > 0$

Proofs

• The existence proof is done on the original formulation of the problem using the diamagnetic inequality

• $\psi(s) e^{ias} = v_1(s) + i v_2(s)$, solves

$$-v_j'' + \alpha v_j = (v_1^2 + v_2^2)^{\frac{p}{2}-1} v_j, \quad j = 1, 2$$

and the Wronskian $w = (v_1 v_2' - v_1' v_2)$ is constant so that $\psi(s) = 0$ is incompatible with the twisted boundary condition

• if $a^2(p+2) + \alpha(p-2) \leq 1$, then $\mu_{a,p}(\alpha) = a^2 + \alpha$ because

$$\begin{aligned} \|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \alpha \|u\|_{L^2(\mathbb{S}^1)}^2 &= (1-4a^2) \|u'\|_{L^2(\mathbb{S}^1)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^1)}^2 \\ &\quad + 4a^2 \left(\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{1}{4} \|u^{-1}\|_{L^2(\mathbb{S}^1)}^2 \right) \end{aligned}$$

if $a^2(p+2) + \alpha(p-2) > 1$, the test function $u_\varepsilon := 1 + \varepsilon w_1$

$$\mathcal{Q}_{a,p,\alpha}[u_\varepsilon] = a^2 + \alpha + (1 - a^2(p+2) - \alpha(p-2)) \varepsilon^2 + o(\varepsilon^2)$$

proves the linear instability of the constants and $\mu_{a,p}(\alpha) < a^2 + \alpha$

$$\mathcal{Q}_{a,p,\alpha}[u] := \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \alpha \|u\|_{L^2(\mathbb{S}^1)}^2}{\|u\|_{L^p(\mathbb{S}^1)}^2},$$

$$\mu_{a,p}(\alpha) = \min_{u \in H^1(\mathbb{S}^1) \setminus \{0\}} \mathcal{Q}_{a,p,\alpha}[u]$$

$$\mathcal{Q}_{p,\alpha}[u] = \mathcal{Q}_{a=0,p,\alpha}[u], \quad \nu_p(\alpha) := \inf_{v \in H_0^1(\mathbb{S}^1) \setminus \{0\}} \mathcal{Q}_{p,\alpha}[v]$$

Proposition

$\forall p > 2, \alpha > -a^2$, we have $\mu_{a,p}(\alpha) < \mu_{1/2,p}(\alpha) \leq \nu_p(\alpha) = \mu_{1/2,p}(\alpha)$




Figure: $p = 4, \alpha = 0, a = 0.40, 0.41, \dots 0.49; u'' + u^{p-1} = 0$

Consequences: Keller-Lieb-Thirring inequalities and Hardy inequalities for Aharonov-Bohm magnetic fields

A Keller-Lieb-Thirring inequality

Magnetic Schrödinger operator $H_a - \varphi = -\left(\frac{d}{ds} + i a\right)^2 \psi - \varphi$

• The function $\alpha \mapsto \mu_{a,p}(\alpha)$ is monotone increasing, concave, and therefore has an inverse, denoted by $\alpha_{a,p} : \mathbb{R}^+ \rightarrow (-a^2, +\infty)$, which is monotone increasing, and convex

Corollary

Let $p > 2$, $a \in [0, 1/2]$, $q = p/(p-2)$ and assume that φ is a non-negative function in $L^q(\mathbb{S}^1)$. Then

$$\lambda_1(H_a - \varphi) \geq -\alpha_{a,p}(\|\varphi\|_{L^q(\mathbb{S}^1)})$$

and $\alpha_{a,p}(\mu) = \mu - a^2$ iff $4a^2 + \mu(p-2) \leq 1$ (optimal φ is constant)

Equality is achieved

Aharonov-Bohm magnetic fields

On the two-dimensional Euclidean space \mathbb{R}^2 , let us introduce the polar coordinates $(r, \vartheta) \in [0, +\infty) \times \mathbb{S}^1$ of $\mathbf{x} \in \mathbb{R}^2$ and consider a magnetic potential \mathbf{a} in a transversal (Poincaré) gauge, or Poincaré gauge

$$(\mathbf{a}, \mathbf{e}_r) = 0 \quad \text{and} \quad (\mathbf{a}, \mathbf{e}_\vartheta) = a_\vartheta(r, \vartheta)$$

Magnetic Schrödinger energy

$$\int_{\mathbb{R}^2} |(i \nabla + \mathbf{a}) \Psi|^2 d\mathbf{x} = \int_0^{+\infty} \int_{-\pi}^{\pi} \left(|\partial_r \Psi|^2 + \frac{1}{r^2} |\partial_\vartheta \Psi + i r a_\vartheta \Psi|^2 \right) r d\vartheta dr$$

Aharonov-Bohm magnetic fields: $a_\vartheta(r, \vartheta) = a/r$ for some constant $a \in \mathbb{R}$ (a is the magnetic flux), with magnetic field $b = \text{curl } \mathbf{a}$

$$\int_{\mathbb{R}^2} |(i \nabla + \mathbf{a}) \Psi|^2 d\mathbf{x} \geq \tau \int_{\mathbb{R}^2} \frac{\varphi(\mathbf{x}/|\mathbf{x}|)}{|\mathbf{x}|^2} |\Psi|^2 d\mathbf{x} \quad \forall \varphi \in L^q(\mathbb{S}^1), \quad q \in (1, +\infty)$$

$$\implies \tau = \tau(a, \|\varphi\|_{L^q(\mathbb{S}^1)}) \quad ?$$

Hardy inequalities

[Hoffmann-Ostenhof, Laptev, 2015] proved Hardy's inequality

$$\int_{\mathbb{R}^d} |\nabla \Psi|^2 d\mathbf{x} \geq \tau \int_{\mathbb{R}^d} \frac{\varphi(\mathbf{x}/|\mathbf{x}|)}{|\mathbf{x}|^2} |\Psi|^2 d\mathbf{x}$$

where the constant τ depends on the value of $\|\varphi\|_{L^q(\mathbb{S}^{d-1})}$ and $d \geq 3$

Aharonov-Bohm vector potential in dimension $d = 2$

$$\mathbf{a}(\mathbf{x}) = a \left(\frac{x_2}{|\mathbf{x}|^2}, \frac{-x_1}{|\mathbf{x}|^2} \right), \quad \mathbf{x} = (x_1, x_2) \in \mathbb{R}^2, \quad a \in \mathbb{R}$$

and recall the inequality [Laptev, Weidl, 1999]

$$\int_{\mathbb{R}^2} |(i \nabla + \mathbf{a}) \Psi|^2 d\mathbf{x} \geq \min_{k \in \mathbb{Z}} (a - k)^2 \int_{\mathbb{R}^2} \frac{|\Psi|^2}{|\mathbf{x}|^2} d\mathbf{x}$$

A new Hardy inequality

$$\int_{\mathbb{R}^2} |(i \nabla + \mathbf{a}) \Psi|^2 d\mathbf{x} \geq \tau \int_{\mathbb{R}^2} \frac{\varphi(\mathbf{x}/|\mathbf{x}|)}{|\mathbf{x}|^2} |\Psi|^2 d\mathbf{x} \quad \forall \varphi \in L^q(\mathbb{S}^1), \quad q \in (1, +\infty)$$

Corollary

Let $p > 2$, $a \in [0, 1/2]$, $q = p/(p-2)$ and assume that φ is a non-negative function in $L^q(\mathbb{S}^1)$. Then the inequality holds with $\tau > 0$ given by

$$\alpha_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^1)}) = 0$$

Moreover, $\tau = a^2 / \|\varphi\|_{L^q(\mathbb{S}^1)}$ if $4a^2 + \|\varphi\|_{L^q(\mathbb{S}^1)}(p-2) \leq 1$

For any $a \in (0, 1/2)$, by taking φ constant, small enough in order that $4a^2 + \|\varphi\|_{L^q(\mathbb{S}^1)}(p-2) \leq 1$, we recover the inequality

$$\int_{\mathbb{R}^2} |(i \nabla + \mathbf{a}) \Psi|^2 d\mathbf{x} \geq a^2 \int_{\mathbb{R}^2} \frac{|\Psi|^2}{|\mathbf{x}|^2} d\mathbf{x}$$

Proofs (Keller-Lieb-Thirring inequality)

Hölder's inequality

$$\|\psi' + i a \psi\|_{L^2(\mathbb{S}^d)}^2 - \int_{-\pi}^{\pi} \varphi |\psi|^2 d\sigma \geq \|\psi' + i a \psi\|_{L^2(\mathbb{S}^d)}^2 - \mu \|\psi\|_{L^p(\mathbb{S}^d)}^2$$

where $\mu = \|\varphi\|_{L^q(\mathbb{S}^d)}$ and $\frac{1}{q} + \frac{2}{p} = 1$: choose $\mu_{a,p}(\alpha) = \mu$

$$\|\psi' + i a \psi\|_{L^2(\mathbb{S}^d)}^2 - \mu \|\psi\|_{L^p(\mathbb{S}^d)}^2 \geq -\alpha \|\psi\|_{L^2(\mathbb{S}^d)}^2$$

Proofs (Hardy inequality)

Let $\tau \geq 0$, $\mathbf{x} = (r, \vartheta) \in \mathbb{R}^2$ be polar coordinates in \mathbb{R}^2

$$\begin{aligned} & \int_{\mathbb{R}^2} \left(|(i \nabla + \mathbf{a}) \Psi|^2 - \tau \frac{\varphi}{|\mathbf{x}|^2} |\Psi|^2 \right) d\mathbf{x} \\ &= \int_0^\infty \int_{\mathbb{S}^1} \left(\underbrace{r |\partial_r \Psi|^2}_{\geq 0} + \frac{1}{r} |\partial_\vartheta \Psi + i a \Psi|^2 - \tau \frac{\varphi}{r} |\Psi|^2 \right) d\vartheta dr \\ &\geq \lambda_1 (H_a - \tau \varphi) \int_0^\infty \int_{\mathbb{S}^1} \frac{1}{r} |\Psi|^2 d\vartheta dr \\ &\geq -\alpha_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^d)}) \int_0^\infty \int_{\mathbb{S}^1} \frac{1}{r} |\Psi|^2 d\vartheta dr \end{aligned}$$

• If $\tau = 0$, then $\alpha_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^d)}) = \alpha_{a,p}(0) = -a^2$

• $\alpha_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^d)}) > 0$ for τ large

$\implies \exists ! \tau > 0$ such that $\alpha_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^d)}) = 0$

Comments

- ▷ The region $\alpha^2(p+2) + \alpha(p-2) < 1$ is exactly the set where the constant functions are linearly stable critical points
- ▷ The proof of the *rigidity result* is based
 - neither on the *carré du champ* method, at least directly
 - nor on a Fourier representation of the operator as it was the case in earlier proofs ($p = +\infty$, or $p > 2$ and $\alpha = 0$)
- ▷ Magnetic rings: see [Bonnaillie-Noël, Hérau, Raymond, 2017]
- ▷ Deducing *Hardy's inequality* applied with *Aharonov-Bohm* magnetic fields from a *Keller-Lieb-Thirring inequality* is an extension of [Hoffmann-Ostenhof, Laptev, 2015] to the magnetic case
- ▷ Our results are not limited to the semi-classical regime

Symmetry in Aharonov-Bohm magnetic fields

- Aharonov-Bohm effect
- Interpolation and Keller-Lieb-Thirring inequalities in \mathbb{R}^2
 - ▷ Statements
 - ▷ Constants and numerics
- Symmetry and symmetry breaking

Joint work with D. Bonheure, M.J. Esteban, A. Laptev, & M. Loss

Aharonov-Bohm effect

A major difference between classical mechanics and quantum mechanics is that particles are described by a non-local object, the wave function. In 1959 Y. Aharonov and D. Bohm proposed a series of experiments intended to put in evidence such phenomena which are nowadays called *Aharonov-Bohm effects*

One of the proposed experiments relies on a long, thin solenoid which produces a magnetic field such that the region in which the magnetic field is non-zero can be approximated by a line in dimension $d = 3$ and by a point in dimension $d = 2$

▷ [Physics today, 2009] *“The notion, introduced 50 years ago, that electrons could be affected by electromagnetic potentials without coming in contact with actual force fields was received with a skepticism that has spawned a flourishing of experimental tests and expansions of the original idea.”* Problem solved by considering appropriate weak solutions !

▷ Is the wave function a physical object or is the modulus the only relevant quantity ? Decisive experiments have been done only 20 ▶

The interpolation inequality

Let us consider an Aharonov-Bohm vector potential

$$\mathbf{A}(x) = \frac{a}{|x|^2} (x_2, -x_1), \quad x = (x_1, x_2) \in \mathbb{R}^2 \setminus \{0\}, \quad a \in \mathbb{R}$$

Magnetic Hardy inequality [Laptev, Weidl, 1999]

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \min_{k \in \mathbb{Z}} (a - k)^2 \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx$$

where $\nabla_{\mathbf{A}} \psi := \nabla \psi + i \mathbf{A} \psi$, so that, with $\psi = |\psi| e^{iS}$

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx = \int_{\mathbb{R}^2} \left[(\partial_r |\psi|)^2 + (\partial_r S)^2 |\psi|^2 + \frac{1}{r^2} (\partial_\theta S + A)^2 |\psi|^2 \right] dx$$

Magnetic interpolation inequality

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \lambda \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx \geq \mu(\lambda) \left(\int_{\mathbb{R}^2} \frac{|\psi|^p}{|x|^2} dx \right)^{2/p}$$

▷ Symmetrization: [Erdős, 1996], [Boulenger, Lenzmann], [Lenzmann, Sok]

A magnetic Hardy-Sobolev inequality

Theorem

Let $a \in [0, 1/2]$ and $p > 2$. For any $\lambda > -a^2$, there is an optimal, monotone increasing, concave function $\lambda \mapsto \mu(\lambda)$ which is such that

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \lambda \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx \geq \mu(\lambda) \left(\int_{\mathbb{R}^2} \frac{|\psi|^p}{|x|^2} dx \right)^{2/p}$$

If $\lambda \leq \lambda_{\star} = 4 \frac{1 - 4a^2}{p^2 - 4} - a^2$ equality is achieved by

$$\psi(x) = (|x|^{\alpha} + |x|^{-\alpha})^{-\frac{2}{p-2}} \quad \forall x \in \mathbb{R}^2, \quad \text{with} \quad \alpha = \frac{p-2}{2} \sqrt{\lambda + a^2}$$

If $\lambda > \lambda_{\bullet}$ with

$$\lambda_{\bullet} := \frac{8 \left(\sqrt{p^4 - a^2 (p-2)^2 (p+2) (3p-2) + 2} \right) - 4p(p+4)}{(p-2)^3 (p+2)} - a^2$$

A magnetic Keller-Lieb-Thirring estimate

Let $q \in (1, +\infty)$ and denote by $L_\star^q(\mathbb{R}^2)$ the space defined using the weighted norm $\|\phi\|_q := \left(\int_{\mathbb{R}^2} |\phi|^q |x|^{2(q-1)} dx \right)^{1/q}$

Theorem

Let $a \in (0, 1/2)$, $q \in (1, \infty)$ and $\phi \in L_\star^q(\mathbb{R}^2)$: $\mu \mapsto \lambda(\mu)$ is a convex monotone increasing function such that $\lim_{\mu \rightarrow 0^+} \lambda(\mu) = -\min_{k \in \mathbb{Z}} (a - k)^2$ and

$$\lambda_1(-\Delta_{\mathbf{A}} - \phi) \geq -\lambda \left(\|\phi\|_q \right)$$

There is an explicit $\mu_\star > 0$ such that the equality case is achieved for any $\mu \leq \mu_\star$ by

$$\phi(x) = (|x|^\alpha + |x|^{-\alpha})^{-2} \quad \forall x \in \mathbb{R}^2, \quad \text{with} \quad \alpha = \frac{p-2}{2} \sqrt{\lambda(\mu) + a^2}$$

There is an explicit $\mu_\bullet > \mu_\star$ such that the equality case is achieved only by non-radial functions if $\mu > \mu_\bullet$.

Constants are explicit...

- For $a = 1/2$, we shall see that $\mu_{\bullet} = \mu_{\star} = -1/4$
- The function $\lambda \mapsto \mu(\lambda)$ is the inverse of $\mu \mapsto \lambda(\mu)$ and

$$\mu_{\star} = h(\lambda_{\star}) \quad \text{and} \quad \mu_{\bullet} = h(\lambda_{\bullet})$$

with

$$h(\lambda) := \frac{p}{2} (2\pi)^{1-\frac{2}{p}} (\lambda + a^2)^{1+\frac{2}{p}} \left(\frac{2\sqrt{\pi} \Gamma(\frac{p}{p-2})}{(p-2) \Gamma(\frac{p}{p-2} + \frac{1}{2})} \right)^{1-\frac{2}{p}}$$




Figure: Case $p = 4$

Symmetry breaking region: $\lambda > \lambda_{\bullet}(a)$

Symmetry breaking region: $\lambda < \lambda_{\star}$




Figure: The curve $a \mapsto \lambda_{\bullet}(a) - \lambda_{\star}(a)$

Lemma

Let $a \in [0, 1/2]$ and $\psi = u e^{iS} \in C^1 \cap H_{\mathbf{A}}^1$ such that $|\psi| > 0$

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \int_{\mathbb{R}^2} \left(|\partial_r u|^2 + \frac{1}{r^2} |\partial_\theta u|^2 + \frac{1}{r^2} \frac{a^2}{\int_{\mathbb{S}^2} u^{-2} d\sigma} \right) dx$$

Equality holds if and only if $\partial_r S \equiv 0$ and

$$\partial_\theta S = a - \frac{a}{u^2} \frac{1}{\int_{\mathbb{S}^2} u^{-2} d\sigma}$$

When u does not depend on θ , equality is achieved iff S is constant

Lemma

For all $a \in [0, 1/2]$ and $\psi \in H^1(\mathbb{S}^1)$ with $u = |\psi|$, we have

$$\int_{\mathbb{S}^2} |\partial_\theta \psi - i a \psi|^2 d\sigma \geq (1 - 4a^2) \int_{\mathbb{S}^2} |\partial_\theta u|^2 d\sigma + a^2 \int_{\mathbb{S}^2} u^2 d\sigma$$

Proof (1/3): the inequality with a non-optimal constant

Diamagnetic inequality: $\|\nabla_{\mathbf{A}} \psi\|_{L^2(\mathbb{R}^2)} \geq \|\nabla u\|_{L^2(\mathbb{R}^2)}$, $u = |\psi|$

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \lambda \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx \\ \geq t \left(\|\nabla_{\mathbf{A}} \psi\|_{L^2(\mathbb{R}^2)}^2 - a^2 \int_{\mathbb{R}^2} \frac{u^2}{|x|^2} dx \right) \\ + (1-t) \left(\|\nabla u\|_{L^2(\mathbb{R}^2)}^2 + \frac{\lambda + a^2 t}{1-t} \int_{\mathbb{R}^2} \frac{u^2}{|x|^2} dx \right) \end{aligned}$$

With $a^2 = \frac{\lambda + a^2 t}{1-t}$, $t \in (0, 1)$ such that $\lambda + a^2 t > 0$: existence of a positive constant $\mu(\lambda)$

Proof (2/3): optimal estimate in the symmetry range

With $a \in [0, 1/2]$, $\psi \in H^1(\mathbb{R}^2)$ and $u = |\psi|$

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \int_{\mathbb{R}^2} |\partial_r u|^2 dx + (1 - 4a^2) \int_{\mathbb{R}^2} \frac{1}{r^2} |\partial_\theta u|^2 dx + a^2 \int_{\mathbb{R}^2} u^2 dx$$

The relaxed inequality

$$\int_{\mathbb{R}^2} \left(|\partial_r u|^2 + \frac{1-4a^2}{r^2} |\partial_\theta u|^2 \right) dx + (\lambda + a^2) \int_{\mathbb{R}^2} \frac{|u|^2}{|x|^2} dx \geq \mu_{\text{rel}}(\lambda) \left(\int_{\mathbb{R}^2} \frac{|u|^p}{|x|^2} dx \right)^{\frac{2}{p}}$$

is rewritten on the cylinder $\mathcal{C} := \mathbb{R} \times \mathbb{S}^1$ using the Emden-Fowler transformation as

$$\begin{aligned} \int_{\mathcal{C}} (|\partial_s w|^2 + (1 - 4a^2) |\partial_\theta w|^2) dy + (\lambda + a^2) \int_{\mathcal{C}} |w|^2 dy \\ \geq (2\pi)^{\frac{2}{p}-1} \mu_{\text{rel}}(\lambda) \left(\int_{\mathcal{C}} |w|^p dy \right)^{\frac{2}{p}} \end{aligned}$$

If $(\lambda + a^2)(p^2 - 4) \leq 4(1 - 4a^2) \iff \lambda \leq \lambda_*$, the minimizer is symmetric

Proof (3/3): symmetry breaking range

$$\mathcal{E}_{a,\lambda}[\psi] := \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \lambda \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx - \mu \left(\int_{\mathbb{R}^2} \frac{|\psi|^p}{|x|^2} dx \right)^{2/p}$$

$$\mu = (2\pi \int_C |w_\star|^p dy)^{1-2/p}, \quad w_\star(s) = \zeta_\star (\cosh(\omega s))^{-\frac{2}{p-2}}$$

$$s = -\log r \text{ and } \psi_\varepsilon(r, \theta) := (w_\star(s) + \varepsilon \varphi(s, \theta)) e^{i\varepsilon \chi(s, \theta)}$$

$$\mathcal{E}_{a,\lambda}[\psi_\varepsilon] = \varepsilon^2 \mathcal{Q}[\varphi, \chi] + o(\varepsilon^2)$$

$$\begin{aligned} \mathcal{Q}[\varphi, \chi] &= \int_C w_\star^2 (|\partial_s \chi|^2 + |\partial_\theta \chi - a|^2 - a^2) dy - 4a \int_C w_\star \varphi \partial_\theta \chi dy \\ &\quad + \int_C (|\partial_s \varphi|^2 + |\partial_\theta \varphi|^2 + (\lambda + a^2) \varphi^2) dy \\ &\quad - (p-1) \int_C |w_\star|^{p-2} |\varphi|^2 dy \end{aligned}$$

$$\varphi(s, \theta) = \frac{\cos \theta}{\cosh(\omega s)^{\frac{p}{p-2}}}, \quad \chi(s, \theta) = \frac{\zeta_\star}{\zeta_\star} \frac{\sin \theta}{\cosh(\omega s)}:$$

$$\mathcal{Q}[\varphi, \chi] < 0 \implies \lambda > \lambda_\bullet$$

These slides can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Conferences/>
▷ Lectures

The papers can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Preprints/list/>
▷ Preprints / papers

For final versions, use **Dolbeault** as login and **Jean** as password

Thank you for your attention !

- 1 Symmetry in non-magnetic interpolation inequalities
 - Interpolation on the sphere
 - Keller-Lieb-Thirring inequalities on the sphere
 - CKN inequalities, symmetry breaking and weighted nonlinear flows
- 2 Magnetic interpolation in the Euclidean space
 - Three interpolation inequalities and their dual forms
 - Proofs for general magnetic fields
 - Estimates in dimension $d = 2$ for constant magnetic fields
 - Numerical results and the symmetry issue
- 3 Magnetic rings: the one-dimensional periodic case
 - Magnetic interpolation on the circle
 - Proof: how to eliminate the phase
 - Consequences: Keller-Lieb-Thirring and Hardy inequalities
- 4 Symmetry in Aharonov-Bohm magnetic fields
 - Aharonov-Bohm effect
 - Interpolation and Keller-Lieb-Thirring inequalities
 - Symmetry and symmetry breaking