

Symétrie et brisure de symétrie dans les inégalités de Caffarelli-Kohn-Nirenberg

Jean Dolbeault

<http://www.ceremade.dauphine.fr/~dolbeault>

Ceremade, Université Paris-Dauphine

11 décembre 2015

Collège de France
Chaire d'EDP et Applications (Pierre-Louis Lions)
Séminaire de Mathématiques Appliquées

Outline

- An introduction to symmetry and symmetry breaking results in weighted elliptic PDEs
- Caffarelli-Kohn-Nirenberg inequalities
 - ▷ the symmetry issue
 - ▷ the result
- The proof
 - ▷ a change of variables and a Sobolev type inequality
 - ▷ the fast diffusion flow and the nonlinear Fisher information
 - ▷ regularity, decay and integrations by parts
- Concavity of the Rényi entropy powers: role of the nonlinear flow
- The Bakry-Emery method: curvature, linear and nonlinear flows (on the sphere)
- Constraints and improvements

In collaboration with M.J. Esteban and M. Loss


An introduction to symmetry and symmetry breaking results in weighted elliptic PDEs

▷ *The typical issue is the competition between a potential or a weight and a nonlinearity*



The *mexican hat* potential

Let us consider a nonlinear Schrödinger equation in presence of a radial external potential with a minimum which is not at the origin

$$-\Delta u + V(x) u - f(u) = 0$$





A one-dimensional potential $V(x)$



A two-dimensional potential $V(x)$ with mexican hat shape



Radial solutions to $-\Delta u + V(x)u - F'(u) = 0$



... give rise to a radial density of energy $x \mapsto V|u|^2 + F(u)$

symmetry breaking

... but in some cases minimal energy solutions



... give rise to a non-radial density of energy $x \mapsto V|u|^2 + F(u)$

Symmetry breaking may occur...

- ▷ in presence of *non-cooperative* potentials
- ▷ in weighted equations or weighted variational problems
- ▷ in phase transition problems
- ▷ in evolution equations (instability of symmetric solutions)
- ▷ in various models of mathematical physics and quantum field theory
etc.

Symmetry and symmetry breaking

Proving symmetry breaking

- ▷ Linear instability of radial critical point (perturbation of a radial solution and energy descent)
- ▷ direct energy estimates (use of special test functions, comparison with the solutions to another problem “at infinity”)

Methods for proving symmetry

Classical methods (a non exhaustive list)

- Alexandrov moving planes and the result of [B. Gidas, W. Ni, L. Nirenberg (1979, 1980)]

$$-\Delta u = f(|x|, u) \quad \text{in } \mathbb{R}^d \quad d \geq 3$$

If f is of class C^1 , $\frac{\partial f}{\partial r} < 0$, $u \geq 0$ is of class C^2 and sufficiently decaying at infinity, then u is a radial function and $\frac{\partial u}{\partial r} < 0$.

- Reflexion with respect planes and unique continuation [O. Lopes]
- Symmetrization methods: Schwarz, Steiner, etc.
- A priori estimates, direct energy estimates
- Uniqueness or rigidity: [B. Gidas, J. Spruck], [M.-F. Bidaut-Véron, L. Véron, 1991]
- Probabilistic methods (Markov processes), semi-group theory and *carré du champ* methods [D. Bakry, M. Emery, 1984]

▷ *A new method based on entropy functionals and evolution under the action of a nonlinear flow: flow interpretation, non-compact case*

Caffarelli-Kohn-Nirenberg inequalities

▷ *Nonlinear flows (fast diffusion equation) can be used as a tool for the investigation of sharp functional inequalities*

Caffarelli-Kohn-Nirenberg inequalities and the symmetry breaking issue

Let $\mathcal{D}_{a,b} := \left\{ v \in L^p(\mathbb{R}^d, |x|^{-b} dx) : |x|^{-a} |\nabla v| \in L^2(\mathbb{R}^d, dx) \right\}$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx \quad \forall v \in \mathcal{D}_{a,b}$$

hold under the conditions that $a \leq b \leq a + 1$ if $d \geq 3$, $a < b \leq a + 1$ if $d = 2$, $a + 1/2 < b \leq a + 1$ if $d = 1$, and $a < a_c := (d - 2)/2$

$$p = \frac{2d}{d - 2 + 2(b - a)}$$

▷ *An optimal function among radial functions:*


$$v_\star(x) = \left(1 + |x|^{(p-2)(a_c-a)} \right)^{-\frac{2}{p-2}} \quad \text{and} \quad C_{a,b}^\star = \frac{\| |x|^{-b} v_\star \|_p^2}{\| |x|^{-a} \nabla v_\star \|_2^2}$$

Question: $C_{a,b} = C_{a,b}^\star$ (symmetry) or $C_{a,b} > C_{a,b}^\star$ (symmetry breaking) ?

CKN: range of the parameters

Figure: $d = 3$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx$$



$a \leq b \leq a + 1$ if $d \geq 3$

$a < b \leq a + 1$ if $d = 2$, $a + 1/2 < b \leq a + 1$ if $d = 1$

and $a < a_c := (d - 2)/2$

$$p = \frac{2d}{d - 2 + 2(b - a)}$$


[Glaser, Martin, Grosse, Thirring (1976)]

[F. Catrina, Z.-Q. Wang (2001)]

Symmetry and symmetry breaking in Caffarelli-Kohn-Nirenberg inequalities


Proving symmetry breaking

[F. Catrina, Z.-Q. Wang], [V. Felli, M. Schneider (2003)]



[J.D., Esteban, Loss, Tarantello, 2009] There is a curve which separates the symmetry region from the symmetry breaking region, which is parametrized by a function $p \mapsto a + b$

Moving planes and symmetrization techniques




[Chou, Chu], [Horiuchi]

[Betta, Brock, Mercaldo, Posteraro]

+ Perturbation results: [CS Lin, ZQ Wang], [Smets, Willem], [JD, Esteban, Tarantello 2007], [J.D., Esteban, Loss, Tarantello, 2009]

Linear instability of radial minimizers: the Felli-Schneider curve




[Catrina, Wang], [Felli, Schneider] The functional

$$C_{a,b}^* \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx - \left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p}$$


is linearly instable at $v = v_*$

Direct spectral estimates



[J.D., Esteban, Loss, 2011]: sharp interpolation on the sphere and a Keller-Lieb-Thirring spectral estimate on the line


Numerical results



Parametric plot of the branch of optimal functions for $p = 2.8$, $d = 5$. Non-symmetric solutions bifurcate from symmetric ones at a bifurcation point computed by V. Felli and M. Schneider. The branch behaves for large values of Λ as predicted by F. Catrina and Z.-Q. Wang

Other evidences


- Further numerical results [J.D., Esteban, 2012] (coarse / refined / self-adaptive grids)



- Formal commutation of the non-symmetric branch near the bifurcation point [J.D., Esteban, 2013]
- Asymptotic energy estimates [J.D., Esteban, 2013]

Symmetry *versus* symmetry breaking: the sharp result

A result based on entropies and nonlinear flows



[J.D., Esteban, Loss, 2015]: <http://arxiv.org/abs/1506.03664>

The symmetry result

The Felli & Schneider curve is defined by

$$b_{\text{FS}}(a) := \frac{d(a_c - a)}{2\sqrt{(a_c - a)^2 + d - 1}} + a - a_c$$

Theorem

Let $d \geq 2$ and $p < 2^$. If either $a \in [0, a_c)$ and $b > 0$, or $a < 0$ and $b \geq b_{\text{FS}}(a)$, then the optimal functions for the Caffarelli-Kohn-Nirenberg inequalities are radially symmetric*

The Emden-Fowler transformation and the cylinder

▷ *With an Emden-Fowler transformation, Caffarelli-Kohn-Nirenberg inequalities on the Euclidean space are equivalent to Gagliardo-Nirenberg inequalities on a cylinder*

$$v(r, \omega) = r^{a-a_c} \varphi(s, \omega) \quad \text{with} \quad r = |x|, \quad s = -\log r \quad \text{and} \quad \omega = \frac{x}{r}$$

With this transformation, the Caffarelli-Kohn-Nirenberg inequalities can be rewritten as

$$\|\partial_s \varphi\|_{L^2(\mathcal{C})}^2 + \|\nabla_\omega \varphi\|_{L^2(\mathcal{C})}^2 + \Lambda \|\varphi\|_{L^2(\mathcal{C})}^2 \geq \mu(\Lambda) \|\varphi\|_{L^p(\mathcal{C})}^2 \quad \forall \varphi \in H^1(\mathcal{C})$$

where $\Lambda := (a_c - a)^2$, $\mathcal{C} = \mathbb{R} \times \mathbb{S}^{d-1}$ and the optimal constant $\mu(\Lambda)$ is

$$\mu(\Lambda) = \frac{1}{C_{a,b}} \quad \text{with} \quad a = a_c \pm \sqrt{\Lambda} \quad \text{and} \quad b = \frac{d}{p} \pm \sqrt{\Lambda}$$

Generalizations and comments

Generalized Caffarelli-Kohn-Nirenberg inequalities (CKN)

Let $2^* = \infty$ if $d = 1$ or $d = 2$, $2^* = 2d/(d - 2)$ if $d \geq 3$ and define

$$\vartheta(p, d) := \frac{d(p - 2)}{2p}$$

[Caffarelli-Kohn-Nirenberg-84] Let $d \geq 1$. For any $\theta \in [\vartheta(p, d), 1]$, with $p = \frac{2d}{d-2+2(b-a)}$, there exists a positive constant $C_{\text{CKN}}(\theta, p, a)$ such that

$$\left(\int_{\mathbb{R}^d} \frac{|u|^p}{|x|^{bp}} dx \right)^{\frac{2}{p}} \leq C_{\text{CKN}}(\theta, p, a) \left(\int_{\mathbb{R}^d} \frac{|\nabla u|^2}{|x|^{2a}} dx \right)^{\theta} \left(\int_{\mathbb{R}^d} \frac{|u|^2}{|x|^{2(a+1)}} dx \right)^{1-\theta}$$

In the radial case, with $\Lambda = (a - a_c)^2$, the best constant when the inequality is restricted to radial functions is $C_{\text{CKN}}^*(\theta, p, a)$ and

$$C_{\text{CKN}}(\theta, p, a) \geq C_{\text{CKN}}^*(\theta, p, a) = C_{\text{CKN}}^*(\theta, p) \Lambda^{\frac{p-2}{2p} - \theta}$$

$$C_{\text{CKN}}^*(\theta, p) = \left[\frac{2\pi^{d/2}}{\Gamma(d/2)} \right]^2 \frac{p-1}{p} \left[\frac{(p-2)^2}{2+(2\theta-1)p} \right]^{\frac{p-2}{2p}} \left[\frac{2+(2\theta-1)p}{2p\theta} \right]^{\theta} \left[\frac{4}{p+2} \right]^{\frac{6-p}{2p}} \left[\frac{\Gamma(\frac{2}{p-2} + \frac{1}{2})}{\sqrt{\pi} \Gamma(\frac{2}{p-2})} \right]^{\frac{p-2}{p}}$$

Implementing the method of Catrina-Wang / Felli-Schneider

Among functions $w \in H^1(\mathcal{C})$ which depend only on s , the minimum of

$$\mathcal{J}[w] := \int_{\mathcal{C}} (|\nabla w|^2 + \frac{1}{4} (d-2-2a)^2 |w|^2) dx - [\mathcal{C}^*(\theta, p, a)]^{-\frac{1}{\theta}} \frac{(\int_{\mathcal{C}} |w|^p dx)^{\frac{2}{p\theta}}}{(\int_{\mathcal{C}} |w|^2 dx)^{\frac{1-\theta}{\theta}}}$$

is achieved by $\bar{w}(y) := [\cosh(\lambda s)]^{-\frac{2}{p-2}}$, $y = (s, \omega) \in \mathbb{R} \times \mathbb{S} = \mathcal{C}$ with

$\lambda := \frac{1}{4} (d-2-2a) (p-2) \sqrt{\frac{p+2}{2p\theta-(p-2)}}$ as a solution of

$$\lambda^2 (p-2)^2 w'' - 4w + 2p |w|^{p-2} w = 0$$

Spectrum of $\mathcal{L} := -\Delta + \kappa \bar{w}^{p-2} + \mu$ is given for $\sqrt{1 + 4\kappa/\lambda^2} \geq 2j + 1$

by $\lambda_{i,j} = \mu + i(d+i-2) - \frac{\lambda^2}{4} \left(\sqrt{1 + 4\kappa/\lambda^2} - (1+2j) \right)^2 \quad \forall i, j \in \mathbb{N}$

- The eigenspace of \mathcal{L} corresponding to $\lambda_{0,0}$ is generated by \bar{w}
- The eigenfunction $\phi_{(1,0)}$ associated to $\lambda_{1,0}$ is not radially symmetric and such that $\int_{\mathcal{C}} \bar{w} \phi_{(1,0)} dx = 0$ and $\int_{\mathcal{C}} \bar{w}^{p-1} \phi_{(1,0)} dx = 0$
- If $\lambda_{1,0} < 0$, optimal functions for (CKN) cannot be radially symmetric and $\mathcal{C}(\theta, p, a) > \mathcal{C}^*(\theta, p, a)$

A parametrization of the solutions

All optimal functions can be computed

▷ by solving (numerically)

$$-\Delta u + \mu u = u^{p-1}$$

▷ by computing $\Lambda = \Lambda^\theta(\mu)$ (reparametrization)

and the corresponding optimal constant is given by

$$J^\theta(\mu) := Q_{\Lambda^\theta(\mu)}^\theta[u_\mu]$$


where

$$Q_\Lambda^\theta[u] := \frac{\left(\|\nabla u\|_{\mathcal{L}^2(C)}^2 + \Lambda \|u\|_{\mathcal{L}^2(C)}^2\right)^\theta \|u\|_{\mathcal{L}^2(C)}^{2(1-\theta)}}{\|u\|_{\mathcal{L}^p(C)}^2}$$


If u_μ is symmetric, then

$$J^\theta(\mu) = J^\theta(1) \mu^\alpha$$


Parametric plot of $\mu \mapsto (\Lambda^\theta(\mu), J^\theta(\mu))$ for $p = 2.8$, $d = 5$, $\theta = 1$




Parametric plot of $\mu \mapsto (\Lambda^\theta(\mu), J^\theta(\mu))$ for $p = 2.8$, $d = 5$, $\theta = 0.8$




Parametric plot of $\mu \mapsto (\Lambda^\theta(\mu), J^\theta(\mu))$ for $p = 2.8$, $d = 5$, $\theta = 0.72$




Enlargement for $p = 2.8$, $d = 5$, $\theta = 0.95$




Enlargement for $p = 2.8$, $d = 5$, $\theta = 0.72$




Critical case $\theta = \vartheta(p, d)$




Parametric plot of $\mu \mapsto (\Lambda^\theta(\mu), J^\theta(\mu))$ for $p = 3.15$, $d = 5$, $\theta = 1$




Parametric plot of $\mu \mapsto (\Lambda^\theta(\mu), J^\theta(\mu))$ for $p = 3.15$, $d = 5$, $\theta = 0.95$



Case $p = 3.15$, $d = 5$, $\theta = \vartheta(3.15, 5) \approx 0.9127$



Local and asymptotic criteria for $\theta = \vartheta(p, d)$ – numerical



- Local criterion: based on an expansion of the solutions near the bifurcation point, it decides whether the branch goes to the right or to the left.
- Asymptotic criterion: based on the energy of the branch as $\Lambda \rightarrow +\infty$ and an analysis in a semi-classical regime

The main steps of the proof

- A change of variables: an equivalent inequality of Sobolev type
- The fast diffusion flow and the nonlinear Fisher information
- Proving the decay along the flow
- The justification of the integration by parts: decay estimates on the cylinder

A change of variables

With $(r = |x|, \omega = x/r) \in \mathbb{R}^+ \times \mathbb{S}^{d-1}$, the Caffarelli-Kohn-Nirenberg inequality is

$$\left(\int_0^\infty \int_{\mathbb{S}^{d-1}} |v|^p r^{d-bp} \frac{dr}{r} d\omega \right)^{\frac{2}{p}} \leq C_{a,b} \int_0^\infty \int_{\mathbb{S}^{d-1}} |\nabla v|^2 r^{d-2a} \frac{dr}{r} d\omega$$

Change of variables $r \mapsto r^\alpha$, $v(r, \omega) = w(r^\alpha, \omega)$

$$\begin{aligned} & \alpha^{1-\frac{2}{p}} \left(\int_0^\infty \int_{\mathbb{S}^{d-1}} |w|^p r^{\frac{d-bp}{\alpha}} \frac{dr}{r} d\omega \right)^{\frac{2}{p}} \\ & \leq C_{a,b} \int_0^\infty \int_{\mathbb{S}^{d-1}} \left(\alpha^2 \left| \frac{\partial w}{\partial r} \right|^2 + \frac{1}{r^2} |\nabla_\omega w|^2 \right) r^{\frac{d-2a-2}{\alpha} + 2} \frac{dr}{r} d\omega \end{aligned}$$

Choice of α

$$n = \frac{d-bp}{\alpha} = \frac{d-2a-2}{\alpha} + 2$$

Then $p = \frac{2n}{n-2}$ is the critical Sobolev exponent associated with n

A Sobolev type inequality

The parameters α and n vary in the ranges $0 < \alpha < \infty$ and $d < n < \infty$ and the *Felli-Schneider curve* in the (α, n) variables is given by

$$\alpha = \sqrt{\frac{d-1}{n-1}} =: \alpha_{\text{FS}}$$

With

$$Dw = \left(\alpha \frac{\partial w}{\partial r}, \frac{1}{r} \nabla_{\omega} w \right), \quad d\mu := r^{n-1} dr d\omega$$

the inequality becomes

$$\alpha^{1-\frac{2}{p}} \left(\int_{\mathbb{R}^d} |w|^p d\mu \right)^{\frac{2}{p}} \leq C_{a,b} \int_{\mathbb{R}^d} |Dw|^2 d\mu$$

Proposition

Let $d \geq 2$. Optimality is achieved by radial functions and $C_{a,b} = C_{a,b}^*$ if $\alpha \leq \alpha_{\text{FS}}$

Gagliardo-Nirenberg inequalities on general cylinders: similar

Notations

When there is no ambiguity, we will omit the index ω and from now on write that $\nabla = \nabla_\omega$ denotes the gradient with respect to the angular variable $\omega \in \mathbb{S}^{d-1}$ and that Δ is the Laplace-Beltrami operator on \mathbb{S}^{d-1} . We define the self-adjoint operator \mathcal{L} by

$$\mathcal{L} w := -D^* D w = \alpha^2 w'' + \alpha^2 \frac{n-1}{r} w' + \frac{\Delta w}{r^2}$$

The fundamental property of \mathcal{L} is the fact that

$$\int_{\mathbb{R}^d} w_1 \mathcal{L} w_2 d\mu = - \int_{\mathbb{R}^d} Dw_1 \cdot Dw_2 d\mu \quad \forall w_1, w_2 \in \mathcal{D}(\mathbb{R}^d)$$

▷ Heuristics: we look for a monotonicity formula along a well chosen nonlinear flow, based on the analogy with the decay of the Fisher information along the fast diffusion flow in \mathbb{R}^d

Fisher information

$$\text{Let } u^{\frac{1}{2}-\frac{1}{n}} = |w| \iff u = |w|^p, p = \frac{2n}{n-2}$$

$$\mathcal{I}[u] := \int_{\mathbb{R}^d} u |\text{Dp}|^2 d\mu, \quad \text{p} = \frac{m}{1-m} u^{m-1} \quad \text{and} \quad m = 1 - \frac{1}{n}$$

Here \mathcal{I} is the *Fisher information* and p is the *pressure function*

Proposition

With $\Lambda = 4\alpha^2/(p-2)^2$ and for some explicit numerical constant κ , we have

$$\kappa \mu(\Lambda) = \inf \{ \mathcal{I}[u] : \|u\|_{L^1(\mathbb{R}^d, d\mu)} = 1 \}$$

▷ Optimal solutions (solutions of the elliptic PDE) are (constrained) critical point of \mathcal{I}

The fast diffusion equation

$$\frac{\partial u}{\partial t} = \mathcal{L} u^m, \quad m = 1 - \frac{1}{n}$$

Barenblatt self-similar solutions

$$u_*(t, r, \omega) = t^{-n} \left(c_* + \frac{r^2}{2(n-1)\alpha^2 t^2} \right)^{-n}$$

Lemma

Barenblatt solutions realize the minimum of \mathcal{I} among radial functions:

$$\kappa \mu_*(\Lambda) = \mathcal{I}[u_*(t, \cdot)] \quad \forall t > 0$$

▷ Strategy:

1) prove that $\frac{d}{dt} \mathcal{I}[u(t, \cdot)] \leq 0$,

2) prove that $\frac{d}{dt} \mathcal{I}[u(t, \cdot)] = 0$ means that $u = u_*$ up to a time shift

Decay of the Fisher information along the flow ?

The *pressure function* $p = \frac{m}{1-m} u^{m-1}$ satisfies

$$\frac{\partial p}{\partial t} = \frac{1}{n} p \mathcal{L} p - |Dp|^2$$

$$\mathcal{Q}[p] := \frac{1}{2} \mathcal{L} |Dp|^2 - Dp \cdot D\mathcal{L} p$$

$$\mathcal{K}[p] := \int_{\mathbb{R}^d} k[p] p^{1-n} d\mu = \int_{\mathbb{R}^d} \left(\mathcal{Q}[p] - \frac{1}{n} (\mathcal{L} p)^2 \right) p^{1-n} d\mu$$

Lemma

If u solves the weighted fast diffusion equation, then

$$\frac{d}{dt} \mathcal{I}[u(t, \cdot)] = -2(n-1)^{n-1} \mathcal{K}[p]$$

If u is a critical point, then $\mathcal{K}[p] = 0$

▷ Boundary terms ! Regularity !

Proving decay (1/2)

$$k[p] := Q(p) - \frac{1}{n} (\mathcal{L} p)^2 = \frac{1}{2} \mathcal{L} |Dp|^2 - Dp \cdot D \mathcal{L} p - \frac{1}{n} (\mathcal{L} p)^2$$

$$k_{\mathfrak{M}}[p] := \frac{1}{2} \Delta |\nabla p|^2 - \nabla p \cdot \nabla \Delta p - \frac{1}{n-1} (\Delta p)^2 - (n-2) \alpha^2 |\nabla p|^2$$

Lemma

Let $n \neq 1$ be any real number, $d \in \mathbb{N}$, $d \geq 2$, and consider a function $p \in C^3((0, \infty) \times \mathfrak{M})$, where (\mathfrak{M}, g) is a smooth, compact Riemannian manifold. Then we have

$$k[p] = \alpha^4 \left(1 - \frac{1}{n}\right) \left[p'' - \frac{p'}{r} - \frac{\Delta p}{\alpha^2 (n-1) r^2} \right]^2 + 2\alpha^2 \frac{1}{r^2} \left| \nabla p' - \frac{\nabla p}{r} \right|^2 + \frac{1}{r^4} k_{\mathfrak{M}}[p]$$

Proving decay (2/2)

Lemma

Assume that $d \geq 3$, $n > d$ and $\mathfrak{M} = \mathbb{S}^{d-1}$. For some $\zeta_\star > 0$ we have

$$\int_{\mathbb{S}^{d-1}} k_{\mathfrak{M}}[p] p^{1-n} d\omega \geq (\lambda_\star - (n-2)\alpha^2) \int_{\mathbb{S}^{d-1}} |\nabla p|^2 p^{1-n} d\omega \\ + \zeta_\star (n-d) \int_{\mathbb{S}^{d-1}} |\nabla p|^4 p^{1-n} d\omega$$

Proof based on the Bochner-Lichnerowicz-Weitzenböck formula

Corollary

Let $d \geq 2$ and assume that $\alpha \leq \alpha_{\text{FS}}$. Then for any nonnegative function $u \in L^1(\mathbb{R}^d)$ with $\mathcal{I}[u] < +\infty$ and $\int_{\mathbb{R}^d} u d\mu = 1$, we have

$$\mathcal{I}[u] \geq \mathcal{I}_\star$$

When $\mathfrak{M} = \mathbb{S}^{d-1}$, $\lambda_\star = (n-2) \frac{d-1}{n-1}$

A perturbation argument

- If u is a critical point of \mathcal{I} under the mass constraint $\int_{\mathbb{R}^d} u \, d\mu = 1$, then

$$o(\varepsilon) = \mathcal{I}[u + \varepsilon \mathcal{L} u^m] - \mathcal{I}[u] = -2(n-1)^{n-1} \varepsilon \mathcal{K}[p] + o(\varepsilon)$$

because $\varepsilon \mathcal{L} u^m$ is an admissible perturbation (formal). Indeed, we know that

$$\int_{\mathbb{R}^d} (u + \varepsilon \mathcal{L} u^m) \, d\mu = \int_{\mathbb{R}^d} u \, d\mu = 1$$

but positivity of $u + \varepsilon \mathcal{L} u^m$ is an issue: compute

$$0 = D\mathcal{I}[u] \cdot \mathcal{L} u^m = -\mathcal{K}[p]$$

- Regularity issues (uniform decay of various derivatives up to order 3) and boundary terms
- If $\alpha \leq \alpha_{\text{FS}}$, then $\mathcal{K}[p] = 0$ implies that $u = u_*$

The justification of the integration by parts: decay estimates on the cylinder

After the Emden-Fowler transformation, a critical point satisfies the Euler-Lagrange equation

$$-\partial_s^2 \varphi - \Delta_\omega \varphi + \Lambda \varphi = \varphi^{p-1} \quad \text{in } \mathcal{C} = \mathbb{R} \times \mathcal{M}$$

(up to a multiplication by a constant; $\mathcal{M} = \mathbb{S}^{d-1}$ e.g.)

Proposition

For all $(s, \omega) \in \mathcal{C}$, we have $C_1 e^{-\sqrt{\Lambda}|s|} \leq \varphi(s, \omega) \leq C_2 e^{-\sqrt{\Lambda}|s|}$

$$|\varphi'(s, \omega)|, |\varphi''(s, \omega)|, |\nabla \varphi(s, \omega)|, |\Delta \varphi(s, \omega)| \leq C_2 e^{-\sqrt{\Lambda}|s|}$$

$$\int_{\mathcal{M}} |\mathbf{p}'(r, \omega)|^2 d\nu_g \leq O(1), \quad \int_{\mathcal{M}} |\nabla \mathbf{p}(r, \omega)|^2 d\nu_g \leq O(r^2),$$

$$\int_{\mathcal{M}} |\mathbf{p}''(r, \omega)|^2 d\nu_g \leq O(1/r^2)$$

$$\int_{\mathcal{M}} \left| \nabla \mathbf{p}'(r, \omega) - \frac{1}{r} \nabla \mathbf{p}(r, \omega) \right|^2 d\nu_g \leq O(1),$$

$$\int_{\mathcal{M}} |\Delta \mathbf{p}(r, \omega)|^2 d\nu_g \leq O(1/r^2)$$

Some comments on the method

The flow interpretation is very useful to organize the computations. It unifies *rigidity* methods and *carré du champ* (or Γ_2) techniques in a common framework which has a clear variational interpretation and opens a door for improvements

However, the proof is done at a purely *variational* level: we consider a critical point and test it against a special test function built on top of the solution, in order to obtain another identity involving sum of squares: each of these square has to be equal to zero, which provides *additional equations* and allow to identify the critical point with a known one (uniqueness result)

In a sense, this method relates with similar estimates:

- ▷ *Nehari manifolds*: one tests the equation with the function u itself. At variational level, this amounts to test the homogeneity of the energy (if any).
- ▷ *Pohozaev's method*: one tests the equation with $x \cdot \nabla u$, which corresponds to a local dilation. At variational level, this amounts to test the energy under scalings (eventually localized) and remarkably, this is a method to produce uniqueness results [Schmitt], [Schaeffer]
- ▷ Here we test the equation with Δu^m , which has to do with *convexity* (displacement convexity) and produces uniqueness (rigidity) results.

Two ingredients for the proof

- ▷ Rényi entropy powers and fast diffusion
- ▷ Flows on the sphere

Rényi entropy powers and fast diffusion

- ▷ Rényi entropy powers, the entropy approach without rescaling: [Savaré, Toscani]: scalings, nonlinearity and a concavity property inspired by information theory
- ▷ faster rates of convergence: [Carrillo, Toscani], [JD, Toscani]

The fast diffusion equation in original variables

Consider the nonlinear diffusion equation in \mathbb{R}^d , $d \geq 1$

$$\frac{\partial u}{\partial t} = \Delta u^m$$

with initial datum $u(x, t = 0) = u_0(x) \geq 0$ such that $\int_{\mathbb{R}^d} u_0 dx = 1$ and $\int_{\mathbb{R}^d} |x|^2 u_0 dx < +\infty$. The large time behavior of the solutions is governed by the source-type Barenblatt solutions

$$u_\star(t, x) := \frac{1}{(\kappa t^{1/\mu})^d} \mathcal{B}_\star\left(\frac{x}{\kappa t^{1/\mu}}\right)$$

where

$$\mu := 2 + d(m - 1), \quad \kappa := \left| \frac{2\mu m}{m - 1} \right|^{1/\mu}$$

and \mathcal{B}_\star is the Barenblatt profile

$$\mathcal{B}_\star(x) := \begin{cases} (C_\star - |x|^2)_+^{1/(m-1)} & \text{if } m > 1 \\ (C_\star + |x|^2)^{1/(m-1)} & \text{if } m < 1 \end{cases}$$

The Rényi entropy power F

The *entropy* is defined by

$$E := \int_{\mathbb{R}^d} u^m dx$$

and the *Fisher information* by

$$I := \int_{\mathbb{R}^d} u |\nabla p|^2 dx \quad \text{with} \quad p = \frac{m}{m-1} u^{m-1}$$

If u solves the fast diffusion equation, then

$$E' = (1 - m) I$$

To compute I' , we will use the fact that

$$\frac{\partial p}{\partial t} = (m-1) p \Delta p + |\nabla p|^2$$

$$F := E^\sigma \quad \text{with} \quad \sigma = \frac{\mu}{d(1-m)} = 1 + \frac{2}{1-m} \left(\frac{1}{d} + m - 1 \right) = \frac{2}{d} \frac{1}{1-m} - 1$$

has a linear growth asymptotically as $t \rightarrow +\infty$

The concavity property

Theorem

[Toscani-Savaré] Assume that $m \geq 1 - \frac{1}{d}$ if $d > 1$ and $m > 0$ if $d = 1$. Then $F(t)$ is increasing, $(1 - m)F''(t) \leq 0$ and

$$\lim_{t \rightarrow +\infty} \frac{1}{t} F(t) = (1 - m) \sigma \lim_{t \rightarrow +\infty} E^{\sigma-1} I = (1 - m) \sigma E_{\star}^{\sigma-1} I_{\star}$$

[Dolbeault-Toscani] The inequality

$$E^{\sigma-1} I \geq E_{\star}^{\sigma-1} I_{\star}$$

is equivalent to the Gagliardo-Nirenberg inequality

$$\|\nabla w\|_{L^2(\mathbb{R}^d)}^{\theta} \|w\|_{L^{q+1}(\mathbb{R}^d)}^{1-\theta} \geq C_{\text{GN}} \|w\|_{L^{2q}(\mathbb{R}^d)}$$

if $1 - \frac{1}{d} \leq m < 1$. Hint: $u^{m-1/2} = \frac{w}{\|w\|_{L^{2q}(\mathbb{R}^d)}}$, $q = \frac{1}{2m-1}$

The proof

Lemma

If u solves $\frac{\partial u}{\partial t} = \Delta u^m$ with $\frac{1}{d} \leq m < 1$, then

$$I' = \frac{d}{dt} \int_{\mathbb{R}^d} u |\nabla p|^2 dx = -2 \int_{\mathbb{R}^d} u^m \left(\|D^2 p\|^2 + (m-1) (\Delta p)^2 \right) dx$$

$$\|D^2 p\|^2 = \frac{1}{d} (\Delta p)^2 + \left\| D^2 p - \frac{1}{d} \Delta p \text{Id} \right\|^2$$

$$\begin{aligned} \frac{1}{\sigma(1-m)} E^{2-\sigma} (E^\sigma)'' &= (1-m)(\sigma-1) \left(\int_{\mathbb{R}^d} u |\nabla p|^2 dx \right)^2 \\ &\quad - 2 \left(\frac{1}{d} + m - 1 \right) \int_{\mathbb{R}^d} u^m dx \int_{\mathbb{R}^d} u^m (\Delta p)^2 dx \\ &\quad - 2 \int_{\mathbb{R}^d} u^m dx \int_{\mathbb{R}^d} u^m \left\| D^2 p - \frac{1}{d} \Delta p \text{Id} \right\|^2 dx \end{aligned}$$

Flows on the sphere

▷ The heat flow introduced by D. Bakry and M. Emery (*carré du champ* method) does not cover all exponents up to the critical one

[Bakry, Emery, 1984]

[Bidaut-Véron, Véron, 1991], [Bakry, Ledoux, 1996]

[Demange, 2008], [JD, Esteban, Loss, 2014 & 2015]

The interpolation inequalities

On the d -dimensional sphere, let us consider the interpolation inequality

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \frac{d}{p-2} \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \frac{d}{p-2} \|u\|_{L^p(\mathbb{S}^d)}^2 \quad \forall u \in H^1(\mathbb{S}^d, d\mu)$$

where the measure $d\mu$ is the uniform probability measure on $\mathbb{S}^d \subset \mathbb{R}^{d+1}$ corresponding to the measure induced by the Lebesgue measure on \mathbb{R}^{d+1} , and the exponent $p \geq 1$, $p \neq 2$, is such that

$$p \leq 2^* := \frac{2d}{d-2}$$

if $d \geq 3$. We adopt the convention that $2^* = \infty$ if $d = 1$ or $d = 2$. The case $p = 2$ corresponds to the logarithmic Sobolev inequality

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 \geq \frac{d}{2} \int_{\mathbb{S}^d} |u|^2 \log \left(\frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \right) d\nu_g \quad \forall u \in H^1(\mathbb{S}^d, d\mu) \setminus \{0\}$$

The Bakry-Emery method

Entropy functional

$$\mathcal{E}_p[\rho] := \frac{1}{p-2} \left[\int_{\mathbb{S}^d} \rho^{\frac{2}{p}} d\nu_g - \left(\int_{\mathbb{S}^d} \rho d\nu_g \right)^{\frac{2}{p}} \right] \quad \text{if } p \neq 2$$

$$\mathcal{E}_2[\rho] := \int_{\mathbb{S}^d} \rho \log \left(\frac{\rho}{\|\rho\|_{L^1(\mathbb{S}^d)}} \right) d\nu_g$$

Fisher information functional

$$\mathcal{I}_p[\rho] := \int_{\mathbb{S}^d} |\nabla \rho^{\frac{1}{p}}|^2 d\nu_g$$

Bakry-Emery (carré du champ): use the heat flow $\frac{\partial \rho}{\partial t} = \Delta \rho$ where Δ denotes the Laplace-Beltrami operator on \mathbb{S}^d , and compute

$$\frac{d}{dt} \mathcal{E}_p[\rho] = -\mathcal{I}_p[\rho] \quad \text{and} \quad \frac{d}{dt} \mathcal{I}_p[\rho] \leq -d \mathcal{I}_p[\rho]$$

$\frac{d}{dt} (\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho]) \leq 0 \implies \mathcal{I}_p[\rho] \geq d \mathcal{E}_p[\rho]$ with $\rho = |u|^p$, if

$$p \leq 2^\# := \frac{2d^2+1}{(d-1)^2}$$


The evolution under the fast diffusion flow

To overcome the limitation $p \leq 2^\#$, one can consider a nonlinear diffusion of fast diffusion / porous medium type

$$\frac{\partial \rho}{\partial t} = \Delta \rho^m. \quad (1)$$

[Demange], [JD, Esteban, Kowalczyk, Loss]: for any $p \in [1, 2^*]$

$$\mathcal{K}_p[\rho] := \frac{d}{dt} \left(\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho] \right) \leq 0,$$



(p, m) admissible region, $d = 5$

Sobolev's inequality

The stereographic projection of $\mathbb{S}^d \subset \mathbb{R}^d \times \mathbb{R} \ni (\rho, \phi, z)$ onto \mathbb{R}^d :
to $\rho^2 + z^2 = 1$, $z \in [-1, 1]$, $\rho \geq 0$, $\phi \in \mathbb{S}^{d-1}$ we associate $x \in \mathbb{R}^d$ such
that $r = |x|$, $\phi = \frac{x}{|x|}$

$$z = \frac{r^2 - 1}{r^2 + 1} = 1 - \frac{2}{r^2 + 1}, \quad \rho = \frac{2r}{r^2 + 1}$$

and transform any function u on \mathbb{S}^d into a function v on \mathbb{R}^d using

$$u(y) = \left(\frac{r}{\rho}\right)^{\frac{d-2}{2}} v(x) = \left(\frac{r^2+1}{2}\right)^{\frac{d-2}{2}} v(x) = (1-z)^{-\frac{d-2}{2}} v(x)$$

• $p = 2^*$, $S_d = \frac{1}{4} d(d-2) |\mathbb{S}^d|^{2/d}$: Euclidean Sobolev inequality

$$\int_{\mathbb{R}^d} |\nabla v|^2 dx \geq S_d \left[\int_{\mathbb{R}^d} |v|^{\frac{2d}{d-2}} dx \right]^{\frac{d-2}{d}} \quad \forall v \in \mathcal{D}^{1,2}(\mathbb{R}^d)$$

Schwarz symmetrization and the ultraspherical setting

$(\xi_0, \xi_1, \dots, \xi_d) \in \mathbb{S}^d$, $\xi_d = z$, $\sum_{i=0}^d |\xi_i|^2 = 1$ [Smets-Willem]

Lemma

Up to a rotation, any minimizer of \mathcal{Q} depends only on $\xi_d = z$

- Let $d\sigma(\theta) := \frac{(\sin \theta)^{d-1}}{Z_d} d\theta$, $Z_d := \sqrt{\pi} \frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d+1}{2})}$: $\forall v \in H^1([0, \pi], d\sigma)$

$$\frac{p-2}{d} \int_0^\pi |v'(\theta)|^2 d\sigma + \int_0^\pi |v(\theta)|^2 d\sigma \geq \left(\int_0^\pi |v(\theta)|^p d\sigma \right)^{\frac{2}{p}}$$

- Change of variables $z = \cos \theta$, $v(\theta) = f(z)$

$$\frac{p-2}{d} \int_{-1}^1 |f'|^2 \nu d\nu_d + \int_{-1}^1 |f|^2 d\nu_d \geq \left(\int_{-1}^1 |f|^p d\nu_d \right)^{\frac{2}{p}}$$

where $\nu_d(z) dz = d\nu_d(z) := Z_d^{-1} \nu^{\frac{d}{2}-1} dz$, $\nu(z) := 1 - z^2$

The ultraspherical operator

With $d\nu_d = Z_d^{-1} \nu^{\frac{d}{2}-1} dz$, $\nu(z) := 1 - z^2$, consider the space $L^2((-1, 1), d\nu_d)$ with scalar product

$$\langle f_1, f_2 \rangle = \int_{-1}^1 f_1 f_2 d\nu_d, \quad \|f\|_{L^p(\mathbb{S}^d)} = \left(\int_{-1}^1 f^p d\nu_d \right)^{\frac{1}{p}}$$

The self-adjoint *ultraspherical* operator is

$$\mathcal{L}f := (1 - z^2)f'' - dzf' = \nu f'' + \frac{d}{2}\nu' f'$$

which satisfies $\langle f_1, \mathcal{L}f_2 \rangle = - \int_{-1}^1 f_1' f_2' \nu d\nu_d$

Proposition

Let $p \in [1, 2) \cup (2, 2^*]$, $d \geq 1$. For any $f \in H^1([-1, 1], d\nu_d)$,

$$-\langle f, \mathcal{L}f \rangle = \int_{-1}^1 |f'|^2 \nu d\nu_d \geq d \frac{\|f\|_{L^p(\mathbb{S}^d)}^2 - \|f\|_{L^2(\mathbb{S}^d)}^2}{p - 2}$$

Heat flow and the Bakry-Emery method

With $g = f^p$, i.e. $f = g^\alpha$ with $\alpha = 1/p$

$$\text{(Ineq.) } -\langle f, \mathcal{L} f \rangle = -\langle g^\alpha, \mathcal{L} g^\alpha \rangle =: \mathcal{I}[g] \geq d \frac{\|g\|_{L^1(\mathbb{S}^d)}^{2\alpha} - \|g^{2\alpha}\|_{L^1(\mathbb{S}^d)}}{p-2} =: \mathcal{F}[g]$$

Heat flow

$$\frac{\partial g}{\partial t} = \mathcal{L} g$$

$$\frac{d}{dt} \|g\|_{L^1(\mathbb{S}^d)} = 0, \quad \frac{d}{dt} \|g^{2\alpha}\|_{L^1(\mathbb{S}^d)} = -2(p-2) \langle f, \mathcal{L} f \rangle = 2(p-2) \int_{-1}^1 |f'|^2 \nu \, d\nu$$

which finally gives

$$\frac{d}{dt} \mathcal{F}[g(t, \cdot)] = -\frac{d}{p-2} \frac{d}{dt} \|g^{2\alpha}\|_{L^1(\mathbb{S}^d)} = -2d \mathcal{I}[g(t, \cdot)]$$

$$\text{Ineq.} \iff \frac{d}{dt} \mathcal{F}[g(t, \cdot)] \leq -2d \mathcal{F}[g(t, \cdot)] \iff \frac{d}{dt} \mathcal{I}[g(t, \cdot)] \leq -2d \mathcal{I}[g(t, \cdot)]$$

The equation for $g = f^p$ can be rewritten in terms of f as

$$\frac{\partial f}{\partial t} = \mathcal{L} f + (p-1) \frac{|f'|^2}{f} \nu$$

$$-\frac{1}{2} \frac{d}{dt} \int_{-1}^1 |f'|^2 \nu \, d\nu_d = \frac{1}{2} \frac{d}{dt} \langle f, \mathcal{L} f \rangle = \langle \mathcal{L} f, \mathcal{L} f \rangle + (p-1) \left\langle \frac{|f'|^2}{f} \nu, \mathcal{L} f \right\rangle$$

$$\begin{aligned} \frac{d}{dt} \mathcal{I}[g(t, \cdot)] + 2d \mathcal{I}[g(t, \cdot)] &= \frac{d}{dt} \int_{-1}^1 |f'|^2 \nu \, d\nu_d + 2d \int_{-1}^1 |f'|^2 \nu \, d\nu_d \\ &= -2 \int_{-1}^1 \left(|f''|^2 + (p-1) \frac{d}{d+2} \frac{|f'|^4}{f^2} - 2(p-1) \frac{d-1}{d+2} \frac{|f'|^2 f''}{f} \right) \nu^2 \, d\nu_d \end{aligned}$$

is nonpositive if

$$|f''|^2 + (p-1) \frac{d}{d+2} \frac{|f'|^4}{f^2} - 2(p-1) \frac{d-1}{d+2} \frac{|f'|^2 f''}{f}$$

is pointwise nonnegative, which is granted if

$$\left[(p-1) \frac{d-1}{d+2} \right]^2 \leq (p-1) \frac{d}{d+2} \iff p \leq \frac{2d^2+1}{(d-1)^2} = 2^\# < \frac{2d}{d-2} = 2^*$$

... up to the critical exponent: a proof in two slides

$$\left[\frac{d}{dz}, \mathcal{L} \right] u = (\mathcal{L} u)' - \mathcal{L} u' = -2z u'' - d u'$$

$$\int_{-1}^1 (\mathcal{L} u)^2 d\nu_d = \int_{-1}^1 |u''|^2 \nu^2 d\nu_d + d \int_{-1}^1 |u'|^2 \nu d\nu_d$$

$$\int_{-1}^1 (\mathcal{L} u) \frac{|u'|^2}{u} \nu d\nu_d = \frac{d}{d+2} \int_{-1}^1 \frac{|u'|^4}{u^2} \nu^2 d\nu_d - 2 \frac{d-1}{d+2} \int_{-1}^1 \frac{|u'|^2 u''}{u} \nu^2 d\nu_d$$

On $(-1, 1)$, let us consider the *porous medium (fast diffusion)* flow

$$u_t = u^{2-2\beta} \left(\mathcal{L} u + \kappa \frac{|u'|^2}{u} \nu \right)$$

If $\kappa = \beta(p-2) + 1$, the L^p norm is conserved

$$\frac{d}{dt} \int_{-1}^1 u^{\beta p} d\nu_d = \beta p (\kappa - \beta(p-2) - 1) \int_{-1}^1 u^{\beta(p-2)} |u'|^2 \nu d\nu_d = 0$$

$$f = u^\beta, \|f'\|_{L^2(\mathbb{S}^d)}^2 + \frac{d}{p-2} \left(\|f\|_{L^2(\mathbb{S}^d)}^2 - \|f\|_{L^p(\mathbb{S}^d)}^2 \right) \geq 0 ?$$

$$\begin{aligned} \mathcal{A} := & \int_{-1}^1 |u''|^2 \nu^2 d\nu_d - 2 \frac{d-1}{d+2} (\kappa + \beta - 1) \int_{-1}^1 u'' \frac{|u'|^2}{u} \nu^2 d\nu_d \\ & + \left[\kappa(\beta - 1) + \frac{d}{d+2} (\kappa + \beta - 1) \right] \int_{-1}^1 \frac{|u'|^4}{u^2} \nu^2 d\nu_d \end{aligned}$$

\mathcal{A} is nonnegative for some β if

$$\frac{8d^2}{(d+2)^2} (p-1)(2^* - p) \geq 0$$

\mathcal{A} is a sum of squares if $p \in (2, 2^*)$ for an arbitrary choice of β in a certain interval (depending on p and d)

$$\mathcal{A} = \int_{-1}^1 \left| u'' - \frac{p+2}{6-p} \frac{|u'|^2}{u} \right|^2 \nu^2 d\nu_d \geq 0 \quad \text{if } p = 2^* \text{ and } \beta = \frac{4}{6-p}$$

The rigidity point of view

Which computation have we done ? $u_t = u^{2-2\beta} \left(\mathcal{L} u + \kappa \frac{|u'|^2}{u} \nu \right)$

$$-\mathcal{L} u - (\beta - 1) \frac{|u'|^2}{u} \nu + \frac{\lambda}{p-2} u = \frac{\lambda}{p-2} u^\kappa$$

Multiply by $\mathcal{L} u$ and integrate

$$\dots \int_{-1}^1 \mathcal{L} u u^\kappa d\nu_d = -\kappa \int_{-1}^1 u^\kappa \frac{|u'|^2}{u} d\nu_d$$

Multiply by $\kappa \frac{|u'|^2}{u}$ and integrate

$$\dots = +\kappa \int_{-1}^1 u^\kappa \frac{|u'|^2}{u} d\nu_d$$

The two terms cancel and we are left only with the two-homogenous terms

Constraints and improvements

Integral constraints

Proposition

For any $p \in (2, 2^\#)$, the inequality

$$\int_{-1}^1 |f'|^2 \nu \, d\nu_d + \frac{\lambda}{p-2} \|f\|_2^2 \geq \frac{\lambda}{p-2} \|f\|_p^2$$

$$\forall f \in H^1((-1, 1), d\nu_d) \text{ s.t. } \int_{-1}^1 z |f|^p \, d\nu_d = 0$$

holds with

$$\lambda \geq d + \frac{(d-1)^2}{d(d+2)} (2^\# - p) (\lambda^* - d)$$

Antipodal symmetry

With the additional restriction of *antipodal symmetry*, that is

$$u(-x) = u(x) \quad \forall x \in \mathbb{S}^d$$

Theorem


If $p \in (1, 2) \cup (2, 2^*)$, we have

$$\int_{\mathbb{S}^d} |\nabla u|^2 d\nu_g \geq \frac{d}{p-2} \left[1 + \frac{(d^2-4)(2^*-p)}{d(d+2)+p-1} \right] \left(\|u\|_{L^p(\mathbb{S}^d)}^2 - \|u\|_{L^2(\mathbb{S}^d)}^2 \right)$$

for any $u \in H^1(\mathbb{S}^d, d\mu)$ with antipodal symmetry. The limit case $p = 2$ corresponds to the improved logarithmic Sobolev inequality


$$\int_{\mathbb{S}^d} |\nabla u|^2 d\nu_g \geq \frac{d(d+3)^2}{2(d+1)^2} \int_{\mathbb{S}^d} |u|^2 \log \left(\frac{|u|^2}{\|u\|_{L^2(\mathbb{S}^d)}^2} \right) d\nu_g$$

The larger picture: branches of antipodal solutions



Case $d = 5$, $p = 3$: values of the shooting parameter a as a function of λ

The optimal constant in the antipodal framework



Numerical computation of the optimal constant when $d = 5$ and $1 \leq p \leq 10/3 \approx 3.33$. The limiting value of the constant is numerically found to be equal to $\lambda_\star = 2^{1-2/p} d \approx 6.59754$ with $d = 5$ and $p = 10/3$

Conclusion

- ▷ The *flow interpretation* is a powerful method to organize computations which are otherwise nasty
- ▷ Outcomes of the method: various improvements. We are able to deal with *non-compact cases and weights* but the price to pay for controlling the boundary terms is high
- ▷ A better understanding of the variational structure of the problem opens a new direction for research on *constrained inequalities*, but generic difficulties (non-monotone branches) have to be understood

▷ Papers

<http://www.ceremade.dauphine.fr/~dolbeaul/Preprints/list/>

▷ Computations

<http://www.ceremade.dauphine.fr/~dolbeaul/NumericalGallery/list/>

▷ These slides can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Conferences/>

Thank you for your attention !