

Symétrie et unicité par des méthodes de flots non-linéaires

Jean Dolbeault

<http://www.ceremade.dauphine.fr/~dolbeault>

Ceremade, Université Paris-Dauphine

8 février 2019

Université de Monastir

A result of uniqueness on a classical example

On the sphere \mathbb{S}^d , let us consider the positive solutions of

$$-\Delta u + \lambda u = u^{p-1}$$

$$p \in [1, 2) \cup (2, 2^*] \text{ if } d \geq 3, \quad 2^* = \frac{2d}{d-2}$$

$$p \in [1, 2) \cup (2, +\infty) \text{ if } d = 1, 2$$

Theorem

If $\lambda \leq d$, $u \equiv \lambda^{1/(p-2)}$ is the unique solution

[Gidas & Spruck, 1981], [Bidaut-Véron & Véron, 1991]

Bifurcation point of view




Figure: $(p-2)\lambda \mapsto (p-2)\mu(\lambda)$ with $d=3$

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu(\lambda) \|u\|_{L^p(\mathbb{S}^d)}^2$$

Taylor expansion of $u = 1 + \varepsilon \varphi_1$ as $\varepsilon \rightarrow 0$ with $-\Delta \varphi_1 = d \varphi_1$

$$\mu(\lambda) < \lambda \quad \text{if and only if} \quad \lambda > \frac{d}{p-2}$$

▷ The inequality holds with $\mu(\lambda) = \lambda = \frac{d}{p-2}$ [Bakry & Emery, 1985]
[Beckner, 1993], [Bidaud-Véron & Véron, 1991, Corollary 6.1]

Inequalities without weights and fast diffusion equations: optimality and uniqueness of the critical points

- The Bakry-Emery method (compact manifolds)
 - ▷ The Fokker-Planck equation
 - ▷ The Bakry-Emery method on the sphere: a parabolic method
 - ▷ The Moser-Trudiger-Onofri inequality (on a compact manifold)
- Fast diffusion equations on the Euclidean space (without weights)
 - ▷ Euclidean space: Rényi entropy powers
 - ▷ Euclidean space: self-similar variables and relative entropies
 - ▷ The role of the spectral gap

The Fokker-Planck equation

The linear Fokker-Planck (FP) equation

$$\frac{\partial u}{\partial t} = \Delta u + \nabla \cdot (u \nabla \phi)$$

on a domain $\Omega \subset \mathbb{R}^d$, with no-flux boundary conditions

$$(\nabla u + u \nabla \phi) \cdot \nu = 0 \quad \text{on } \partial\Omega$$

is equivalent to the Ornstein-Uhlenbeck (OU) equation

$$\frac{\partial v}{\partial t} = \Delta v - \nabla \phi \cdot \nabla v =: \mathcal{L} v$$

(Bakry, Emery, 1985), (Arnold, Markowich, Toscani, Unterreiter, 2001)

With mass normalized to 1, the unique stationary solution of (FP) is

$$u_s = \frac{e^{-\phi}}{\int_{\Omega} e^{-\phi} dx} \iff v_s = 1$$

The Bakry-Emery method

With $d\gamma = u_s dx$ and ν such that $\int_{\Omega} \nu d\gamma = 1$, $q \in (1, 2]$, the q -entropy is defined by

$$\mathcal{E}_q[\nu] := \frac{1}{q-1} \int_{\Omega} (\nu^q - 1 - q(\nu - 1)) d\gamma$$

Under the action of (OU), with $w = \nu^{q/2}$, $\mathcal{I}_q[\nu] := \frac{4}{q} \int_{\Omega} |\nabla w|^2 d\gamma$,

$$\frac{d}{dt} \mathcal{E}_q[\nu(t, \cdot)] = -\mathcal{I}_q[\nu(t, \cdot)] \quad \text{and} \quad \frac{d}{dt} (\mathcal{I}_q[\nu] - 2\lambda \mathcal{E}_q[\nu]) \leq 0$$

$$\text{with } \lambda := \inf_{w \in H^1(\Omega, d\gamma) \setminus \{0\}} \frac{\int_{\Omega} (2 \frac{q-1}{q} \|\text{Hess } w\|^2 + \text{Hess } \phi: \nabla w \otimes \nabla w) d\gamma}{\int_{\Omega} |\nabla w|^2 d\gamma}$$

Proposition

(Bakry, Emery, 1984) (JD, Nazaret, Savaré, 2008) *Let Ω be convex. If $\lambda > 0$ and ν is a solution of (OU), then $\mathcal{I}_q[\nu(t, \cdot)] \leq \mathcal{I}_q[\nu(0, \cdot)] e^{-2\lambda t}$ and $\mathcal{E}_q[\nu(t, \cdot)] \leq \mathcal{E}_q[\nu(0, \cdot)] e^{-2\lambda t}$ for any $t \geq 0$ and, as a consequence,*

$$\mathcal{I}_q[\nu] \geq 2\lambda \mathcal{E}_q[\nu] \quad \forall \nu \in H^1(\Omega, d\gamma)$$

A proof of the interpolation inequality by the *carré du champ* method

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 \geq \frac{d}{p-2} \left(\|u\|_{L^p(\mathbb{S}^d)}^2 - \|u\|_{L^2(\mathbb{S}^d)}^2 \right) \quad \forall u \in H^1(\mathbb{S}^d)$$

$$p \in [1, 2) \cup (2, 2^*] \text{ if } d \geq 3, \quad 2^* = \frac{2d}{d-2}$$

$$p \in [1, 2) \cup (2, +\infty) \text{ if } d = 1, 2$$

The Bakry-Emery method on the sphere

Entropy functional

$$\mathcal{E}_p[\rho] := \frac{1}{p-2} \left[\int_{\mathbb{S}^d} \rho^{\frac{2}{p}} d\mu - \left(\int_{\mathbb{S}^d} \rho d\mu \right)^{\frac{2}{p}} \right] \quad \text{if } p \neq 2$$

$$\mathcal{E}_2[\rho] := \int_{\mathbb{S}^d} \rho \log \left(\frac{\rho}{\|\rho\|_{L^1(\mathbb{S}^d)}} \right) d\mu$$

Fisher information functional

$$\mathcal{I}_p[\rho] := \int_{\mathbb{S}^d} |\nabla \rho^{\frac{1}{p}}|^2 d\mu$$

[Bakry & Emery, 1985] *carré du champ* method: use the heat flow

$$\frac{\partial \rho}{\partial t} = \Delta \rho$$

and observe that $\frac{d}{dt} \mathcal{E}_p[\rho] = -\mathcal{I}_p[\rho]$,

$$\frac{d}{dt} \left(\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho] \right) \leq 0 \quad \implies \quad \mathcal{I}_p[\rho] \geq d \mathcal{E}_p[\rho]$$

with $\rho = |u|^p$, if $p \leq 2^\# := \frac{2d^2+1}{(d-1)^2}$


The evolution under the fast diffusion flow

To overcome the limitation $p \leq 2^\#$, one can consider a nonlinear diffusion of fast diffusion / porous medium type

$$\frac{\partial \rho}{\partial t} = \Delta \rho^m$$


(Demange), (JD, Esteban, Kowalczyk, Loss): for any $p \in [1, 2^*]$

$$\mathcal{K}_p[\rho] := \frac{d}{dt} \left(\mathcal{I}_p[\rho] - d \mathcal{E}_p[\rho] \right) \leq 0$$



(p, m) admissible region, $d = 5$

Cylindrical coordinates, Schwarz symmetrization, stereographic projection...



... and the ultra-spherical operator

Change of variables $z = \cos \theta$, $\nu(\theta) = f(z)$, $d\nu_d := \nu^{\frac{d}{2}-1} dz / Z_d$,
 $\nu(z) := 1 - z^2$

The self-adjoint *ultraspherical* operator is

$$\mathcal{L} f := (1 - z^2) f'' - d z f' = \nu f'' + \frac{d}{2} \nu' f'$$

which satisfies $\langle f_1, \mathcal{L} f_2 \rangle = - \int_{-1}^1 f_1' f_2' \nu d\nu_d$

Proposition

Let $p \in [1, 2) \cup (2, 2^*]$, $d \geq 1$. For any $f \in H^1([-1, 1], d\nu_d)$,

$$- \langle f, \mathcal{L} f \rangle = \int_{-1}^1 |f'|^2 \nu d\nu_d \geq d \frac{\|f\|_{L^p(\mathbb{S}^d)}^2 - \|f\|_{L^2(\mathbb{S}^d)}^2}{p - 2}$$

The heat equation $\frac{\partial g}{\partial t} = \mathcal{L} g$ for $g = f^p$ can be rewritten in terms of f as

$$\frac{\partial f}{\partial t} = \mathcal{L} f + (p-1) \frac{|f'|^2}{f} \nu$$

$$-\frac{1}{2} \frac{d}{dt} \int_{-1}^1 |f'|^2 \nu d\nu_d = \frac{1}{2} \frac{d}{dt} \langle f, \mathcal{L} f \rangle = \langle \mathcal{L} f, \mathcal{L} f \rangle + (p-1) \left\langle \frac{|f'|^2}{f} \nu, \mathcal{L} f \right\rangle$$

$$\begin{aligned} \frac{d}{dt} \mathcal{I}[g(t, \cdot)] + 2d \mathcal{I}[g(t, \cdot)] &= \frac{d}{dt} \int_{-1}^1 |f'|^2 \nu d\nu_d + 2d \int_{-1}^1 |f'|^2 \nu d\nu_d \\ &= -2 \int_{-1}^1 \left(|f''|^2 + (p-1) \frac{d}{d+2} \frac{|f'|^4}{f^2} - 2(p-1) \frac{d-1}{d+2} \frac{|f'|^2 f''}{f} \right) \nu^2 d\nu_d \end{aligned}$$

is nonpositive if

$$|f''|^2 + (p-1) \frac{d}{d+2} \frac{|f'|^4}{f^2} - 2(p-1) \frac{d-1}{d+2} \frac{|f'|^2 f''}{f}$$

is pointwise nonnegative, which is granted if

$$\left[(p-1) \frac{d-1}{d+2} \right]^2 \leq (p-1) \frac{d}{d+2} \iff p \leq \frac{2d^2+1}{(d-1)^2} = 2^\# < \frac{2d}{d-2} = 2^*$$

The elliptic point of view (nonlinear flow)

$$\frac{\partial u}{\partial t} = u^{2-2\beta} \left(\mathcal{L} u + \kappa \frac{|u'|^2}{u} \nu \right), \quad \kappa = \beta(p-2) + 1$$

$$- \mathcal{L} u - (\beta-1) \frac{|u'|^2}{u} \nu + \frac{\lambda}{p-2} u = \frac{\lambda}{p-2} u^\kappa$$

Multiply by $\mathcal{L} u$ and integrate

$$\dots \int_{-1}^1 \mathcal{L} u u^\kappa d\nu_d = - \kappa \int_{-1}^1 u^\kappa \frac{|u'|^2}{u} d\nu_d$$

Multiply by $\kappa \frac{|u'|^2}{u}$ and integrate

$$\dots = + \kappa \int_{-1}^1 u^\kappa \frac{|u'|^2}{u} d\nu_d$$

The two terms cancel and we are left only with

$$\int_{-1}^1 \left| u'' - \frac{p+2}{6-p} \frac{|u'|^2}{u} \right|^2 \nu^2 d\nu_d = 0 \quad \text{if } p = 2^* \text{ and } \beta = \frac{4}{6-p}$$

The Moser-Trudinger-Onofri inequality on Riemannian manifolds

Joint work with G. Jankowiak and M.J. Esteban

- Extension to compact Riemannian manifolds of dimension 2...



We shall also denote by \mathfrak{R} the Ricci tensor, by $H_g u$ the Hessian of u and by

$$L_g u := H_g u - \frac{g}{d} \Delta_g u$$

the trace free Hessian. Let us denote by $M_g u$ the trace free tensor

$$M_g u := \nabla u \otimes \nabla u - \frac{g}{d} |\nabla u|^2$$

We define

$$\lambda_\star := \inf_{u \in H^2(\mathfrak{M}) \setminus \{0\}} \frac{\int_{\mathfrak{M}} \left[\|L_g u - \frac{1}{2} M_g u\|^2 + \mathfrak{R}(\nabla u, \nabla u) \right] e^{-u/2} d\nu_g}{\int_{\mathfrak{M}} |\nabla u|^2 e^{-u/2} d\nu_g}$$

Theorem

Assume that $d = 2$ and $\lambda_\star > 0$. If u is a smooth solution to

$$-\frac{1}{2} \Delta_g u + \lambda = e^u$$

then u is a constant function if $\lambda \in (0, \lambda_\star)$

The Moser-Trudinger-Onofri inequality on \mathfrak{M}

$$\frac{1}{4} \|\nabla u\|_{L^2(\mathfrak{M})}^2 + \lambda \int_{\mathfrak{M}} u \, d\nu_g \geq \lambda \log \left(\int_{\mathfrak{M}} e^u \, d\nu_g \right) \quad \forall u \in H^1(\mathfrak{M})$$

for some constant $\lambda > 0$. Let us denote by λ_1 the first positive eigenvalue of $-\Delta_g$

Corollary

If $d = 2$, then the MTO inequality holds with $\lambda = \Lambda := \min\{4\pi, \lambda_\star\}$. Moreover, if Λ is strictly smaller than $\lambda_1/2$, then the optimal constant in the MTO inequality is strictly larger than Λ

The flow

$$\frac{\partial f}{\partial t} = \Delta_g(e^{-f/2}) - \frac{1}{2} |\nabla f|^2 e^{-f/2}$$

$$\begin{aligned} \mathcal{G}_\lambda[f] := \int_{\mathfrak{M}} \|L_g f - \frac{1}{2} M_g f\|^2 e^{-f/2} d\nu_g + \int_{\mathfrak{M}} \Re(\nabla f, \nabla f) e^{-f/2} d\nu_g \\ - \lambda \int_{\mathfrak{M}} |\nabla f|^2 e^{-f/2} d\nu_g \end{aligned}$$

Then for any $\lambda \leq \lambda_\star$ we have

$$\begin{aligned} \frac{d}{dt} \mathcal{F}_\lambda[f(t, \cdot)] &= \int_{\mathfrak{M}} \left(-\frac{1}{2} \Delta_g f + \lambda\right) \left(\Delta_g(e^{-f/2}) - \frac{1}{2} |\nabla f|^2 e^{-f/2}\right) d\nu_g \\ &= -\mathcal{G}_\lambda[f(t, \cdot)] \end{aligned}$$

Since \mathcal{F}_λ is nonnegative and $\lim_{t \rightarrow \infty} \mathcal{F}_\lambda[f(t, \cdot)] = 0$, we obtain that

$$\mathcal{F}_\lambda[u] \geq \int_0^\infty \mathcal{G}_\lambda[f(t, \cdot)] dt$$

Weighted Moser-Trudinger-Onofri inequalities on the two-dimensional Euclidean space

On the Euclidean space \mathbb{R}^2 , given a general probability measure μ does the inequality

$$\frac{1}{16\pi} \int_{\mathbb{R}^2} |\nabla u|^2 dx \geq \lambda \left[\log \left(\int_{\mathbb{R}^2} e^u d\mu \right) - \int_{\mathbb{R}^2} u d\mu \right]$$

hold for some $\lambda > 0$? Let

$$\Lambda_\star := \inf_{x \in \mathbb{R}^2} \frac{-\Delta \log \mu}{8\pi \mu}$$

Theorem

Assume that μ is a radially symmetric function. Then any radially symmetric solution to the EL equation is a constant if $\lambda < \Lambda_\star$ and the inequality holds with $\lambda = \Lambda_\star$ if equality is achieved among radial functions

Euclidean space: Rényi entropy powers and fast diffusion

● The Euclidean space without weights

▷ Rényi entropy powers, the entropy approach without rescaling:
(Savaré, Toscani): scalings, nonlinearity and a concavity property
inspired by information theory

The fast diffusion equation in original variables

Consider the nonlinear diffusion equation in \mathbb{R}^d , $d \geq 1$

$$\frac{\partial v}{\partial t} = \Delta v^m$$

with initial datum $v(x, t = 0) = v_0(x) \geq 0$ such that $\int_{\mathbb{R}^d} v_0 \, dx = 1$ and $\int_{\mathbb{R}^d} |x|^2 v_0 \, dx < +\infty$. The large time behavior of the solutions is governed by the source-type Barenblatt solutions

$$\mathcal{U}_\star(t, x) := \frac{1}{(\kappa t^{1/\mu})^d} \mathcal{B}_\star\left(\frac{x}{\kappa t^{1/\mu}}\right)$$

where

$$\mu := 2 + d(m - 1), \quad \kappa := \left| \frac{2\mu m}{m - 1} \right|^{1/\mu}$$

and \mathcal{B}_\star is the Barenblatt profile

$$\mathcal{B}_\star(x) := \begin{cases} (C_\star - |x|^2)_+^{1/(m-1)} & \text{if } m > 1 \\ (C_\star + |x|^2)^{1/(m-1)} & \text{if } m < 1 \end{cases}$$

The Rényi entropy power F

The *entropy* is defined by

$$E := \int_{\mathbb{R}^d} v^m dx$$

and the *Fisher information* by

$$I := \int_{\mathbb{R}^d} v |\nabla p|^2 dx \quad \text{with} \quad p = \frac{m}{m-1} v^{m-1}$$

If v solves the fast diffusion equation, then

$$E' = (1 - m)I$$

To compute I' , we will use the fact that

$$\frac{\partial p}{\partial t} = (m-1)p \Delta p + |\nabla p|^2$$

$$F := E^\sigma \quad \text{with} \quad \sigma = \frac{\mu}{d(1-m)} = 1 + \frac{2}{1-m} \left(\frac{1}{d} + m - 1 \right) = \frac{2}{d} \frac{1}{1-m} - 1$$

has a linear growth asymptotically as $t \rightarrow +\infty$

The variation of the Fisher information

Lemma

If v solves $\frac{\partial v}{\partial t} = \Delta v^m$ with $1 - \frac{1}{d} \leq m < 1$, then

$$I' = \frac{d}{dt} \int_{\mathbb{R}^d} v |\nabla p|^2 dx = -2 \int_{\mathbb{R}^d} v^m \left(\|D^2 p\|^2 + (m-1) (\Delta p)^2 \right) dx$$

Explicit arithmetic geometric inequality

$$\|D^2 p\|^2 - \frac{1}{d} (\Delta p)^2 = \left\| D^2 p - \frac{1}{d} \Delta p \text{Id} \right\|^2$$

.... there are no boundary terms in the integrations by parts ?



The concavity property

Theorem

[Toscani-Savaré] Assume that $m \geq 1 - \frac{1}{d}$ if $d > 1$ and $m > 0$ if $d = 1$. Then $F(t)$ is increasing, $(1 - m) F''(t) \leq 0$ and

$$\lim_{t \rightarrow +\infty} \frac{1}{t} F(t) = (1 - m) \sigma \lim_{t \rightarrow +\infty} E^{\sigma-1} I = (1 - m) \sigma E_{\star}^{\sigma-1} I_{\star}$$

[Dolbeault-Toscani] The inequality

$$E^{\sigma-1} I \geq E_{\star}^{\sigma-1} I_{\star}$$

is equivalent to the Gagliardo-Nirenberg inequality

$$\|\nabla w\|_{L^2(\mathbb{R}^d)}^{\theta} \|w\|_{L^{q+1}(\mathbb{R}^d)}^{1-\theta} \geq C_{\text{GN}} \|w\|_{L^{2q}(\mathbb{R}^d)}$$

if $1 - \frac{1}{d} \leq m < 1$. Hint: $v^{m-1/2} = \frac{w}{\|w\|_{L^{2q}(\mathbb{R}^d)}} , q = \frac{1}{2m-1}$

Euclidean space: self-similar variables and relative entropies

- In the Euclidean space, it is possible to characterize the optimal constants using a spectral gap property

Self-similar variables and relative entropies

The large time behavior of the solution of $\frac{\partial v}{\partial t} = \Delta v^m$ is governed by the source-type *Barenblatt solutions*

$$v_{\star}(t, x) := \frac{1}{\kappa^d (\mu t)^{d/\mu}} \mathcal{B}_{\star} \left(\frac{x}{\kappa (\mu t)^{1/\mu}} \right) \quad \text{where} \quad \mu := 2 + d(m-1)$$

where \mathcal{B}_{\star} is the Barenblatt profile (with appropriate mass)

$$\mathcal{B}_{\star}(x) := (1 + |x|^2)^{1/(m-1)}$$

A time-dependent rescaling: **self-similar variables**

$$v(t, x) = \frac{1}{\kappa^d R^d} u \left(\tau, \frac{x}{\kappa R} \right) \quad \text{where} \quad \frac{dR}{dt} = R^{1-\mu}, \quad \tau(t) := \frac{1}{2} \log \left(\frac{R(t)}{R_0} \right)$$

Then the function u solves **a Fokker-Planck type equation**

$$\frac{\partial u}{\partial \tau} + \nabla \cdot \left[u (\nabla u^{m-1} - 2x) \right] = 0$$

Free energy and Fisher information

- The function u solves a Fokker-Planck type equation

$$\frac{\partial u}{\partial \tau} + \nabla \cdot \left[u \left(\nabla u^{m-1} - 2x \right) \right] = 0$$

- (Ralston, Newman, 1984) Lyapunov functional:

Generalized entropy or *Free energy*

$$\mathcal{E}[u] := \int_{\mathbb{R}^d} \left(-\frac{u^m}{m} + |x|^2 u \right) dx - \mathcal{E}_0$$

- Entropy production is measured by the *Generalized Fisher information*

$$\frac{d}{dt} \mathcal{E}[u] = -\mathcal{I}[u], \quad \mathcal{I}[u] := \int_{\mathbb{R}^d} u \left| \nabla u^{m-1} + 2x \right|^2 dx$$

Without weights: relative entropy, entropy production

🟢 *Stationary solution:* choose C such that $\|u_\infty\|_{L^1} = \|u\|_{L^1} = M > 0$

$$u_\infty(x) := (C + |x|^2)_+^{-1/(1-m)}$$

🟢 *Entropy – entropy production inequality* (del Pino, JD)

Theorem

$d \geq 3$, $m \in [\frac{d-1}{d}, +\infty)$, $m > \frac{1}{2}$, $m \neq 1$

$$\mathcal{I}[u] \geq 4 \mathcal{E}[u]$$

$p = \frac{1}{2m-1}$, $u = w^{2p}$: (GN) $\|\nabla w\|_{L^2(\mathbb{R}^d)}^\theta \|w\|_{L^{q+1}(\mathbb{R}^d)}^{1-\theta} \geq C_{GN} \|w\|_{L^{2q}(\mathbb{R}^d)}$

Corollary

(del Pino, JD) A solution u with initial data $u_0 \in L^1_+(\mathbb{R}^d)$ such that $|x|^2 u_0 \in L^1(\mathbb{R}^d)$, $u_0^m \in L^1(\mathbb{R}^d)$ satisfies $\mathcal{E}[u(t, \cdot)] \leq \mathcal{E}[u_0] e^{-4t}$

A computation on a large ball, with boundary terms

$$\frac{\partial u}{\partial \tau} + \nabla \cdot \left[u \left(\nabla u^{m-1} - 2x \right) \right] = 0 \quad \tau > 0, \quad x \in B_R$$

where B_R is a centered ball in \mathbb{R}^d with radius $R > 0$, and assume that u satisfies zero-flux boundary conditions

$$\left(\nabla u^{m-1} - 2x \right) \cdot \frac{x}{|x|} = 0 \quad \tau > 0, \quad x \in \partial B_R.$$

With $z(\tau, x) := \nabla Q(\tau, x) := \nabla u^{m-1} - 2x$, the *relative Fisher information* is such that

$$\begin{aligned} & \frac{d}{d\tau} \int_{B_R} u |z|^2 dx + 4 \int_{B_R} u |z|^2 dx \\ & + 2 \frac{1-m}{m} \int_{B_R} u^m \left(\|D^2 Q\|^2 - (1-m)(\Delta Q)^2 \right) dx \\ & = \int_{\partial B_R} u^m (\omega \cdot \nabla |z|^2) d\sigma \leq 0 \quad (\text{by Grisvard's lemma}) \end{aligned}$$

Spectral gap: sharp asymptotic rates of convergence

Assumptions on the initial datum v_0

(H1) $V_{D_0} \leq v_0 \leq V_{D_1}$ for some $D_0 > D_1 > 0$

(H2) if $d \geq 3$ and $m \leq m_*$, $(v_0 - V_D)$ is integrable for a suitable $D \in [D_1, D_0]$

Theorem

(Blanchet, Bonforte, JD, Grillo, Vázquez) Under Assumptions (H1)-(H2), if $m < 1$ and $m \neq m_* := \frac{d-4}{d-2}$, the entropy decays according to


$$\mathcal{E}[v(t, \cdot)] \leq C e^{-2(1-m)\Lambda_{\alpha,d} t} \quad \forall t \geq 0$$

where $\Lambda_{\alpha,d} > 0$ is the best constant in the Hardy-Poincaré inequality

$$\Lambda_{\alpha,d} \int_{\mathbb{R}^d} |f|^2 d\mu_{\alpha-1} \leq \int_{\mathbb{R}^d} |\nabla f|^2 d\mu_{\alpha} \quad \forall f \in H^1(d\mu_{\alpha}), \int_{\mathbb{R}^d} f d\mu_{\alpha-1} = 0$$

with $\alpha := 1/(m-1) < 0$, $d\mu_{\alpha} := h_{\alpha} dx$, $h_{\alpha}(x) := (1 + |x|^2)^{\alpha}$

Spectral gap and best constants



Caffarelli-Kohn-Nirenberg, symmetry and symmetry breaking results, and weighted nonlinear flows

▷ The critical Caffarelli-Kohn-Nirenberg inequality
[JD, Esteban, Loss]

[▷ A family of sub-critical Caffarelli-Kohn-Nirenberg inequalities]
[JD, Esteban, Loss, Muratori]

▷ Large time asymptotics and spectral gaps

▷ Optimality cases

Critical Caffarelli-Kohn-Nirenberg inequality

Let $\mathcal{D}_{a,b} := \left\{ v \in L^p(\mathbb{R}^d, |x|^{-b} dx) : |x|^{-a} |\nabla v| \in L^2(\mathbb{R}^d, dx) \right\}$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx \quad \forall v \in \mathcal{D}_{a,b}$$

holds under conditions on a and b

$$p = \frac{2d}{d-2+2(b-a)} \quad (\text{critical case})$$

▷ *An optimal function among radial functions:*


$$v_*(x) = \left(1 + |x|^{(p-2)(a_c-a)} \right)^{-\frac{2}{p-2}} \quad \text{and} \quad C_{a,b}^* = \frac{\| |x|^{-b} v_* \|_p^2}{\| |x|^{-a} \nabla v_* \|_2^2}$$

Question: $C_{a,b} = C_{a,b}^$ (symmetry) or $C_{a,b} > C_{a,b}^*$ (symmetry breaking) ?*

Critical CKN: range of the parameters

Figure: $d = 3$

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx$$



$a \leq b \leq a + 1$ if $d \geq 3$

$a < b \leq a + 1$ if $d = 2$, $a + 1/2 < b \leq a + 1$ if $d = 1$

and $a < a_c := (d - 2)/2$

$$p = \frac{2d}{d - 2 + 2(b - a)}$$

(Glaser, Martin, Grosse, Thirring (1976))


(Caffarelli, Kohn, Nirenberg (1984))

[F. Catrina, Z.-Q. Wang (2001)]

Linear instability of radial minimizers: the Felli-Schneider curve

The Felli & Schneider curve

$$b_{\text{FS}}(a) := \frac{d(a_c - a)}{2\sqrt{(a_c - a)^2 + d - 1}} + a - a_c$$



[Smets], [Smets, Willem], [Catrina, Wang], [Felli, Schneider]


The functional

$$C_{a,b}^* \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx - \left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p}$$

is linearly unstable at $v = v_*$

Symmetry *versus* symmetry breaking: the sharp result in the critical case

[JD, Esteban, Loss (2016)]



Theorem

Let $d \geq 2$ and $p < 2^*$. If either $a \in [0, a_c)$ and $b > 0$, or $a < 0$ and $b \geq b_{\text{FS}}(a)$, then the optimal functions for the critical Caffarelli-Kohn-Nirenberg inequalities are radially symmetric

The Emden-Fowler transformation and the cylinder

▷ *With an Emden-Fowler transformation, critical the Caffarelli-Kohn-Nirenberg inequality on the Euclidean space are equivalent to Gagliardo-Nirenberg inequalities on a cylinder*

$$v(r, \omega) = r^{a-a_c} \varphi(s, \omega) \quad \text{with} \quad r = |x|, \quad s = -\log r \quad \text{and} \quad \omega = \frac{x}{r}$$

With this transformation, the Caffarelli-Kohn-Nirenberg inequalities can be rewritten as the *subcritical* interpolation inequality

$$\|\partial_s \varphi\|_{L^2(\mathcal{C})}^2 + \|\nabla_\omega \varphi\|_{L^2(\mathcal{C})}^2 + \Lambda \|\varphi\|_{L^2(\mathcal{C})}^2 \geq \mu(\Lambda) \|\varphi\|_{L^p(\mathcal{C})}^2 \quad \forall \varphi \in H^1(\mathcal{C})$$

where $\Lambda := (a_c - a)^2$, $\mathcal{C} = \mathbb{R} \times \mathbb{S}^{d-1}$ and the optimal constant $\mu(\Lambda)$ is

$$\mu(\Lambda) = \frac{1}{C_{a,b}} \quad \text{with} \quad a = a_c \pm \sqrt{\Lambda} \quad \text{and} \quad b = \frac{d}{p} \pm \sqrt{\Lambda}$$

Linearization around symmetric critical points

Up to a normalization and a scaling

$$\varphi_\star(s, \omega) = (\cosh s)^{-\frac{1}{p-2}}$$

is a critical point of

$$H^1(\mathcal{C}) \ni \varphi \mapsto \|\partial_s \varphi\|_{L^2(\mathcal{C})}^2 + \|\nabla_\omega \varphi\|_{L^2(\mathcal{C})}^2 + \Lambda \|\varphi\|_{L^2(\mathcal{C})}^2$$

under a constraint on $\|\varphi\|_{L^p(\mathcal{C})}^2$

φ_\star *is not* optimal for (CKN) if the Pöschl-Teller operator

$$-\partial_s^2 - \Delta_\omega + \Lambda - \varphi_\star^{p-2} = -\partial_s^2 - \Delta_\omega + \Lambda - \frac{1}{(\cosh s)^2}$$

has *a negative eigenvalue*, i.e., for $\Lambda > \Lambda_1$ (explicit)

The variational problem on the cylinder

$$\Lambda \mapsto \mu(\Lambda) := \min_{\varphi \in H^1(C)} \frac{\|\partial_s \varphi\|_{L^2(C)}^2 + \|\nabla_\omega \varphi\|_{L^2(C)}^2 + \Lambda \|\varphi\|_{L^2(C)}^2}{\|\varphi\|_{L^p(C)}^2}$$

is a concave increasing function


Restricted to symmetric functions, the variational problem becomes

$$\mu_\star(\Lambda) := \min_{\varphi \in H^1(\mathbb{R})} \frac{\|\partial_s \varphi\|_{L^2(\mathbb{R}^d)}^2 + \Lambda \|\varphi\|_{L^2(\mathbb{R}^d)}^2}{\|\varphi\|_{L^p(\mathbb{R}^d)}^2} = \mu_\star(1) \Lambda^\alpha$$

Symmetry means $\mu(\Lambda) = \mu_\star(\Lambda)$




Symmetry breaking means $\mu(\Lambda) < \mu_\star(\Lambda)$

Numerical results



Parametric plot of the branch of optimal functions for $p = 2.8$, $d = 5$. Non-symmetric solutions bifurcate from symmetric ones at a bifurcation point Λ_1 computed by V. Felli and M. Schneider. The branch behaves for large values of Λ as predicted by F. Catrina and Z.-Q. Wang

what we have to to prove / discard...



When the local criterion (linear stability) differs from global results in a larger family of inequalities (center, right)...

The uniqueness result and the strategy of the proof

The elliptic problem: rigidity

The symmetry issue can be reformulated as a uniqueness (rigidity) issue. An optimal function for the inequality

$$\left(\int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} dx$$

solves the (elliptic) Euler-Lagrange equation

$$-\nabla \cdot (|x|^{-2a} \nabla v) = |x|^{-bp} v^{p-1}$$

(up to a scaling and a multiplication by a constant). Is any nonnegative solution of such an equation equal to

$$v_*(x) = (1 + |x|^{(p-2)(a_c-a)})^{-\frac{2}{p-2}}$$

(up to invariances) ? On the cylinder

$$-\partial_s^2 \varphi - \partial_\omega \varphi + \Lambda \varphi = \varphi^{p-1}$$

Up to a normalization and a scaling

$$\varphi_*(s, \omega) = (\cosh s)^{-\frac{1}{p-2}}$$

Symmetry in one slide: 3 steps

1 A change of variables: $v(|x|^{\alpha-1}x) = w(x)$, $D_\alpha v = (\alpha \frac{\partial v}{\partial s}, \frac{1}{s} \nabla_\omega v)$

$$\|v\|_{L^{2p,d-n}(\mathbb{R}^d)} \leq K_{\alpha,n,p} \|D_\alpha v\|_{L^{2,d-n}(\mathbb{R}^d)}^\vartheta \|v\|_{L^{p+1,d-n}(\mathbb{R}^d)}^{1-\vartheta} \quad \forall v \in H_{d-n,d-n}^p(\mathbb{R}^d)$$

2 Concavity of the Rényi entropy power: with

$$\mathcal{L}_\alpha = -D_\alpha^* D_\alpha = \alpha^2 \left(u'' + \frac{n-1}{s} u' \right) + \frac{1}{s^2} \Delta_\omega u \text{ and } \frac{\partial u}{\partial t} = \mathcal{L}_\alpha u^m$$

$$\begin{aligned} & - \frac{d}{dt} \mathcal{G}[u(t, \cdot)] \left(\int_{\mathbb{R}^d} u^m d\mu \right)^{1-\sigma} \\ & \geq (1-m)(\sigma-1) \int_{\mathbb{R}^d} u^m \left| \mathcal{L}_\alpha P - \frac{\int_{\mathbb{R}^d} u |D_\alpha P|^2 d\mu}{\int_{\mathbb{R}^d} u^m d\mu} \right|^2 d\mu \\ & + 2 \int_{\mathbb{R}^d} \left(\alpha^4 \left(1 - \frac{1}{n} \right) \left| P'' - \frac{P'}{s} - \frac{\Delta_\omega P}{\alpha^2 (n-1) s^2} \right|^2 + \frac{2\alpha^2}{s^2} \left| \nabla_\omega P' - \frac{\nabla_\omega P}{s} \right|^2 \right) u^m d\mu \\ & + 2 \int_{\mathbb{R}^d} \left((n-2) (\alpha_{\text{FS}}^2 - \alpha^2) |\nabla_\omega P|^2 + c(n, m, d) \frac{|\nabla_\omega P|^4}{P^2} \right) u^m d\mu \end{aligned}$$

3 Elliptic regularity and the Emden-Fowler transformation: justifying the integrations by parts



Proof of symmetry (1/3: changing the dimension)

We rephrase our problem in a space of higher, *artificial dimension* $n > d$ (here n is a dimension at least from the point of view of the scaling properties), or to be precise we consider a weight $|x|^{n-d}$ which is the same in all norms. With $\beta = 2a$ and $\gamma = bp$,

$$v(|x|^{\alpha-1}x) = w(x), \quad \alpha = 1 + \frac{\beta - \gamma}{2} \quad \text{and} \quad n = 2 \frac{d - \gamma}{\beta + 2 - \gamma}$$

we claim that Inequality (CKN) can be rewritten for a function $v(|x|^{\alpha-1}x) = w(x)$ as

$$\|v\|_{L^{2p, d-n}(\mathbb{R}^d)} \leq K_{\alpha, n, p} \|D_{\alpha} v\|_{L^{2, d-n}(\mathbb{R}^d)}^{\vartheta} \|v\|_{L^{p+1, d-n}(\mathbb{R}^d)}^{1-\vartheta} \quad \forall v \in H_{d-n, d-n}^p(\mathbb{R}^d)$$

with the notations $s = |x|$, $D_{\alpha} v = (\alpha \frac{\partial v}{\partial s}, \frac{1}{s} \nabla_{\omega} v)$ and

$$d \geq 2, \quad \alpha > 0, \quad n > d \quad \text{and} \quad p \in (1, p_{\star}]$$

By our change of variables, w_{\star} is changed into

$$v_{\star}(x) := (1 + |x|^2)^{-1/(p-1)} \quad \forall x \in \mathbb{R}^d$$

The strategy of the proof (2/3: Rényi entropy)

The derivative of the generalized *Rényi entropy power* functional is

$$\mathcal{G}[u] := \left(\int_{\mathbb{R}^d} u^m d\mu \right)^{\sigma-1} \int_{\mathbb{R}^d} u |D_\alpha P|^2 d\mu$$

where $\sigma = \frac{2}{d} \frac{1}{1-m} - 1$. Here $d\mu = |x|^{n-d} dx$ and the pressure is

$$P := \frac{m}{1-m} u^{m-1}$$

Looking for an optimal function in (CKN) is equivalent to minimize \mathcal{G} under a mass constraint

With $L_\alpha = -D_\alpha^* D_\alpha = \alpha^2 \left(u'' + \frac{n-1}{s} u' \right) + \frac{1}{s^2} \Delta_\omega u$, we consider the fast diffusion equation

$$\frac{\partial u}{\partial t} = L_\alpha u^m$$

critical case $m = 1 - 1/n$; subcritical range $1 - 1/n < m < 1$

The key computation is the proof that

$$\begin{aligned} & - \frac{d}{dt} \mathcal{G}[u(t, \cdot)] \left(\int_{\mathbb{R}^d} u^m d\mu \right)^{1-\sigma} \\ & \geq (1-m)(\sigma-1) \int_{\mathbb{R}^d} u^m \left| L_\alpha P - \frac{\int_{\mathbb{R}^d} u |D_\alpha P|^2 d\mu}{\int_{\mathbb{R}^d} u^m d\mu} \right|^2 d\mu \\ & + 2 \int_{\mathbb{R}^d} \left(\alpha^4 \left(1 - \frac{1}{n} \right) \left| P'' - \frac{P'}{s} - \frac{\Delta_\omega P}{\alpha^2 (n-1) s^2} \right|^2 + \frac{2\alpha^2}{s^2} \left| \nabla_\omega P' - \frac{\nabla_\omega P}{s} \right|^2 \right) u^m d\mu \\ & + 2 \int_{\mathbb{R}^d} \left((n-2) (\alpha_{\text{FS}}^2 - \alpha^2) |\nabla_\omega P|^2 + c(n, m, d) \frac{|\nabla_\omega P|^4}{P^2} \right) u^m d\mu =: \mathcal{H}[u] \end{aligned}$$

for some numerical constant $c(n, m, d) > 0$. Hence if $\alpha \leq \alpha_{\text{FS}}$, the r.h.s. $\mathcal{H}[u]$ vanishes if and only if P is an affine function of $|x|^2$, which proves the symmetry result. *A quantifier elimination problem (Tarski, 1951) ?*

(3/3: elliptic regularity, boundary terms)

This method has a hidden difficulty: integrations by parts ! Hints:

🟢 use elliptic regularity: Moser iteration scheme, Sobolev regularity, local Hölder regularity, Harnack inequality, and get global regularity using scalings

🟢 use the Emden-Fowler transformation, work on a cylinder, truncate, evaluate boundary terms of high order derivatives using Poincaré inequalities on the sphere

Summary: if u solves the Euler-Lagrange equation, we test by $L_\alpha u^m$

$$0 = \int_{\mathbb{R}^d} d\mathcal{G}[u] \cdot L_\alpha u^m d\mu \geq \mathcal{H}[u] \geq 0$$

$\mathcal{H}[u]$ is the integral of a sum of squares (with nonnegative constants in front of each term)... or test by $|x|^\gamma \operatorname{div}(|x|^{-\beta} \nabla w^{1+p})$ the equation

$$\frac{(p-1)^2}{p(p+1)} w^{1-3p} \operatorname{div}(|x|^{-\beta} w^{2p} \nabla w^{1-p}) + |\nabla w^{1-p}|^2 + |x|^{-\gamma} (c_1 w^{1-p} - c_2) = 0$$

Fast diffusion equations with weights: large time asymptotics

- The entropy formulation of the problem
- [Relative uniform convergence]
- Asymptotic rates of convergence
- From asymptotic to global estimates

Here v solves the *Fokker-Planck type equation*

$$v_t + |x|^\gamma \nabla \cdot \left[|x|^{-\beta} v \nabla (v^{m-1} - |x|^{2+\beta-\gamma}) \right] = 0 \quad (\text{WFDE-FP})$$

Joint work with M. Bonforte, M. Muratori and B. Nazaret

CKN and entropy – entropy production inequalities

When symmetry holds, (CKN) can be written as an *entropy – entropy production* inequality

$$\frac{1-m}{m} (2 + \beta - \gamma)^2 \mathcal{E}[v] \leq \mathcal{I}[v]$$

and equality is achieved by $\mathfrak{B}_{\beta,\gamma}(x) := (1 + |x|^{2+\beta-\gamma})^{\frac{1}{m-1}}$

Here the *free energy* and the *relative Fisher information* are defined by



$$\mathcal{E}[v] := \frac{1}{m-1} \int_{\mathbb{R}^d} \left(v^m - \mathfrak{B}_{\beta,\gamma}^m - m \mathfrak{B}_{\beta,\gamma}^{m-1} (v - \mathfrak{B}_{\beta,\gamma}) \right) \frac{dx}{|x|^\gamma}$$

$$\mathcal{I}[v] := \int_{\mathbb{R}^d} v \left| \nabla v^{m-1} - \nabla \mathfrak{B}_{\beta,\gamma}^{m-1} \right|^2 \frac{dx}{|x|^\beta}$$

If v solves the *Fokker-Planck type equation*

$$v_t + |x|^\gamma \nabla \cdot \left[|x|^{-\beta} v \nabla (v^{m-1} - |x|^{2+\beta-\gamma}) \right] = 0 \quad (\text{WFDE-FP})$$

then $\frac{d}{dt} \mathcal{E}[v(t, \cdot)] = - \frac{m}{1-m} \mathcal{I}[v(t, \cdot)]$



The spectrum of \mathcal{L} as a function of $\delta = \frac{1}{1-m}$, with $n = 5$. The essential spectrum corresponds to the grey area, and its bottom is determined by the parabola $\delta \mapsto \Lambda_{\text{ess}}(\delta)$. The two eigenvalues $\Lambda_{0,1}$ and $\Lambda_{1,0}$ are given by the plain, half-lines, away from the essential spectrum. The spectral gap determines the asymptotic rate of convergence to the Barenblatt functions



Global vs. asymptotic estimates

🟢 *Estimates on the global rates.* When symmetry holds (CKN) can be written as an *entropy - entropy production* inequality

$$(2 + \beta - \gamma)^2 \mathcal{E}[v] \leq \frac{m}{1 - m} \mathcal{I}[v]$$

so that

$$\mathcal{E}[v(t)] \leq \mathcal{E}[v(0)] e^{-2(1-m)\Lambda_\star t} \quad \forall t \geq 0 \quad \text{with} \quad \Lambda_\star := \frac{(2 + \beta - \gamma)^2}{2(1 - m)}$$

🟢 *Optimal global rates.* Let us consider again the *entropy - entropy production* inequality

$$\mathcal{K}(M) \mathcal{E}[v] \leq \mathcal{I}[v] \quad \forall v \in L^{1,\gamma}(\mathbb{R}^d) \quad \text{such that} \quad \|v\|_{L^{1,\gamma}(\mathbb{R}^d)} = M,$$

where $\mathcal{K}(M)$ is the best constant: with $\Lambda(M) := \frac{m}{2} (1 - m)^{-2} \mathcal{K}(M)$

$$\mathcal{E}[v(t)] \leq \mathcal{E}[v(0)] e^{-2(1-m)\Lambda(M)t} \quad \forall t \geq 0$$

Linearization and optimality

Joint work with M.J. Esteban and M. Loss

Linearization and scalar products

With u_ε such that

$$u_\varepsilon = \mathcal{B}_\star (1 + \varepsilon f \mathcal{B}_\star^{1-m}) \quad \text{and} \quad \int_{\mathbb{R}^d} u_\varepsilon \, dx = M_\star$$

at first order in $\varepsilon \rightarrow 0$ we obtain that f solves

$$\frac{\partial f}{\partial t} = \mathcal{L} f \quad \text{where} \quad \mathcal{L} f := (1 - m) \mathcal{B}_\star^{m-2} |x|^\gamma D_\alpha^* (|x|^{-\beta} \mathcal{B}_\star D_\alpha f)$$

Using the scalar products

$$\langle f_1, f_2 \rangle = \int_{\mathbb{R}^d} f_1 f_2 \mathcal{B}_\star^{2-m} |x|^{-\gamma} \, dx \quad \text{and} \quad \langle\langle f_1, f_2 \rangle\rangle = \int_{\mathbb{R}^d} D_\alpha f_1 \cdot D_\alpha f_2 \mathcal{B}_\star |x|^{-\beta} \, dx$$

we compute

$$\frac{1}{2} \frac{d}{dt} \langle f, f \rangle = \langle f, \mathcal{L} f \rangle = \int_{\mathbb{R}^d} f (\mathcal{L} f) \mathcal{B}_\star^{2-m} |x|^{-\gamma} \, dx = - \int_{\mathbb{R}^d} |D_\alpha f|^2 \mathcal{B}_\star |x|^{-\beta} \, dx$$

for any f smooth enough: with $\langle f, \mathcal{L} f \rangle = - \langle\langle f, f \rangle\rangle$

$$\frac{1}{2} \frac{d}{dt} \langle\langle f, f \rangle\rangle = \int_{\mathbb{R}^d} D_\alpha f \cdot D_\alpha (\mathcal{L} f) \mathcal{B}_\star |x|^{-\beta} \, dx = - \langle\langle f, \mathcal{L} f \rangle\rangle$$

Linearization of the flow, eigenvalues and spectral gap

Now let us consider an eigenfunction associated with the smallest positive eigenvalue λ_1 of \mathcal{L}

$$-\mathcal{L} f_1 = \lambda_1 f_1$$

so that f_1 realizes the equality case in the *Hardy-Poincaré inequality*

$$\langle\langle g, g \rangle\rangle := -\langle g, \mathcal{L} g \rangle \geq \lambda_1 \|g - \bar{g}\|^2, \quad \bar{g} := \langle g, 1 \rangle / \langle 1, 1 \rangle \quad (\text{P1})$$

$$-\langle\langle g, \mathcal{L} g \rangle\rangle \geq \lambda_1 \langle\langle g, g \rangle\rangle \quad (\text{P2})$$

Proof by expansion of the square

$$-\langle\langle (g - \bar{g}), \mathcal{L} (g - \bar{g}) \rangle\rangle = \langle \mathcal{L} (g - \bar{g}), \mathcal{L} (g - \bar{g}) \rangle = \|\mathcal{L} (g - \bar{g})\|^2$$

🟢 (P1) is associated with the symmetry breaking issue

🟢 (P2) is associated with the *carré du champ* method

The optimal constants / eigenvalues are the same

🟢 Key observation: $\lambda_1 \geq 4 \iff \alpha \leq \alpha_{\text{FS}} := \sqrt{\frac{d-1}{n-1}}$

Three references

🟢 Lecture notes on *Symmetry and nonlinear diffusion flows...*
a course on entropy methods (see webpage)

🟢 [JD, Maria J. Esteban, and Michael Loss] *Symmetry and symmetry breaking: rigidity and flows in elliptic PDEs*
... the elliptic point of view: arXiv: 1711.11291, Proc. Int. Cong. of Math., Rio de Janeiro, 3: 2279-2304, 2018.

🟢 [JD, Maria J. Esteban, and Michael Loss] *Interpolation inequalities, nonlinear flows, boundary terms, optimality and linearization...* the parabolic point of view
Journal of elliptic and parabolic equations, 2: 267-295, 2016.

These slides can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Conferences/>
▷ Lectures

The papers can be found at

<http://www.ceremade.dauphine.fr/~dolbeaul/Preprints/list/>
▷ Preprints / papers

For final versions, use **Dolbeault** as login and **Jean** as password

Thank you for your attention !