Symmetry breaking issues in Caffarelli-Kohn-Nirenberg inequalities and related problems

Jean Dolbeault

 $http://www.ceremade.dauphine.fr/{\sim}dolbeaul$ 

Ceremade, Université Paris-Dauphine

November 18, 2015

Torino

<ロト <回ト < 注入 < 注入 = 注

# Outline

- An introduction to symmetry and symmetry breaking results in weighted elliptic PDEs
- Caffarelli-Kohn-Nirenberg inequalities
  - $\triangleright$  The symmetry issue
  - $\triangleright$  The result
- The proof
  - ▷ a change of variables and a Sobolev type inequality
     ▷ the fast diffusion flow and the nonlinear Fisher information
  - $\triangleright$  regularity, decay and integrations by parts
- Concavity of the Rényi entropy powers: role of the nonlinear flow
- The Bakry-Emery method: curvature, linear and nonlinear flows
- Conclusion

### In collaboration with M.J. Esteban and M. Loss

・ 同 ト ・ ヨ ト ・ ヨ ト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# An introduction to symmetry and symmetry breaking results in weighted elliptic PDEs

 $\triangleright$  The typical issue is the competition between a potential or a weight and a nonlinearity

- 1月 ト - ヨ ト - - - ト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# The mexican hat potential

Let us consider a nonlinear Schrödinger equation in presence of a radial external potential with a minimum which is not at the origin

$$-\Delta u + V(x) u - f(u) = 0$$



A one-dimensional potential V(x)

- ∢ ≣ >

#### Symmetry and symmetry breaking in elliptic PDEs

Caffarelli-Kohn-Nirenberg inequalities The proof of the symmetry result in 4 steps Two ingredients for the proof and some remarks The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking



A two-dimensional potential V(x) with mexican hat shape

・ロン ・四と ・ヨン ・ヨン

э

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

Radial solutions to  $-\Delta u + V(x)u - F'(u) = 0$ 



... give rise to a radial density of energy  $x \mapsto V |u|^2 + F(u)$ 

イロト イポト イヨト イヨト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# symmetry breaking

... but in some cases minimal energy solutions



... give rise to a non-radial density of energy  $x \mapsto V |u|^2 + F(u)$ 

イロト イポト イヨト イヨト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# Symmetry and symmetry breaking

イロト イポト イヨト イヨト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# Proving symmetry breaking

The most classical method is by perturbation of a radial solution and energy descent

... but there are other methods, like direct energy estimates

< 回 ト く ヨ ト く ヨ ト

The mexican hat potential in Schrödinger equations Symmetry and symmetry breaking

# Methods for proving symmetry

Classical methods (a non exhaustive list)

• Alexandrov moving planes and the result of [B. Gidas, W. Ni, L. Nirenberg (1979, 1980)]

$$-\Delta u = f(|x|, u)$$
 in  $\mathbb{R}^d d \ge 3$ 

If f is of class  $C^1$ ,  $\frac{\partial f}{\partial r} < 0$ ,  $u \ge 0$  is of class  $C^2$  and sufficiently decaying at infinity, then u is a radial function and  $\frac{\partial u}{\partial r} < 0$ .

- Reflexion with respect planes and unique continuation [O. Lopes]
- Symmetrization methods: Schwarz, Steiner, etc.
- A priori estimates, direct energy estimates
- Uniqueness or rigidity: [B. Gidas, J. Spruck], [M.-F. Bidault-Véron, L. Véron, 1991]
- ... probabilistic methods and *carré du champ* methods [D. Bakry, M. Emery, 1984]

 $\triangleright A new method based on entropy functionals and evolution under the action of a nonlinear flow \\ \hline \Box \rightarrow \langle \overline{\sigma} \rangle \land \overline{z} \rightarrow \langle \overline{z} \rangle \land \overline{z} \rightarrow \overline{z}$ 

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Caffarelli-Kohn-Nirenberg inequalities

 $\triangleright$  Nonlinear flows (fast diffusion equation) can be used as a tool for the investigation of sharp functional inequalities

- 4 回 ト 4 ヨ ト

Caffarelli-Kohn-Nirenberg inequalities The proof of the symmetry result in 4 steps Two ingredients for the proof and some remarks Results on CKN inequalities Symmetry and symmetry breaking The sharp result

# Caffarelli-Kohn-Nirenberg inequalities and the symmetry breaking issue

Let 
$$\mathcal{D}_{a,b} := \left\{ v \in \mathrm{L}^{p}\left(\mathbb{R}^{d}, |x|^{-b} dx\right) : |x|^{-a} |\nabla v| \in \mathrm{L}^{2}\left(\mathbb{R}^{d}, dx\right) \right\}$$
$$\left( \int_{\mathbb{R}^{d}} \frac{|v|^{p}}{|x|^{b\,p}} dx \right)^{2/p} \leq C_{a,b} \int_{\mathbb{R}^{d}} \frac{|\nabla v|^{2}}{|x|^{2\,a}} dx \quad \forall v \in \mathcal{D}_{a,b}$$

hold under the conditions that a < b < a + 1 if d > 3, a < b < a + 1if d = 2,  $a + 1/2 < b \le a + 1$  if d = 1, and  $a < a_c := (d - 2)/2$ 

$$p = \frac{2d}{d-2+2(b-a)}$$

With

$$v_{\star}(x) = \left(1 + |x|^{(p-2)(a_{c}-a)}\right)^{-\frac{2}{p-2}} \quad and \quad C_{a,b}^{\star} = \frac{\||x|^{-b} v_{\star}\|_{p}^{2}}{\||x|^{-a} \nabla v_{\star}\|_{2}^{2}}$$

do we have  $C_{a,b} = C^{\star}_{a,b}$  (symmetry) or  $C_{a,b} > C^{\star}_{a,b}$  (symmetry breaking)?

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# CKN: range of the parameters



Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Symmetry and symmetry breaking

イロト イポト イヨト イヨト

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Proving symmetry breaking

[F. Catrina, Z.-Q. Wang], [V. Felli, M. Schneider (2003)]



[J.D., Esteban, Loss, Tarantello, 2009] There is a curve which separates the symmetry region from the symmetry breaking region, which is parametrized by a function  $p \mapsto a + b$ 

くぼう くちゃ くちゃ

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Moving planes and symmetrization techniques



[Chou, Chu], [Horiuchi]
[Betta, Brock, Marcaldo, Posteraro]
+ Perturbation results: [CS Lin, ZQ Wang], [Smets, Willem], [JD, Esteban, Tarantello 2007], [J.D., Esteban, Loss, Tarantello, 2009]

- 4 回 ト 4 三 ト

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Linear instability of radial minimizers: the Felli-Schneider curve



[Catrina, Wang], [Felli, Schneider] The functional

$$C_{a,b}^{\star} \int_{\mathbb{R}^d} \frac{|\nabla v|^2}{|x|^{2a}} \, dx - \left( \int_{\mathbb{R}^d} \frac{|v|^p}{|x|^{bp}} \, dx \right)^{2/p}$$

is linearly instable at  $v=v_\star$ 

< 回 > < 三 > < 三 >

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Direct spectral estimates



[J.D., Esteban, Loss, 2011]: sharp interpolation on the sphere and a Keller-Lieb-Thirring spectral estimate on the line

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

### Numerical results



Parametric plot of the branch of optimal functions for p = 2.8, d = 5. Non-symmetric solutions bifurcate from symmetric ones at a bifurcation point computed by V. Felli and M. Schneider. The branch behaves for large values of  $\Lambda$  as predicted by F. Catrina and Z.-Q. Wang

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Other evidences

• Further numerical results [J.D., Esteban, 2012] (coarse / refined / self-adaptive grids)



• Formal commutation of the non-symmetric branch near the bifurcation point [J.D., Esteban, 2013]

Results on CKN inequalities Symmetry and symmetry breaking **The sharp result** Generalizations and comments

# Symmetry *versus* symmetry breaking: the sharp result

A result based on entropies and nonlinear flows



[J.D., Esteban, Loss, 2015]: http://arxiv.org/abs/1506.03664

Results on CKN inequalities Symmetry and symmetry breaking **The sharp result** Generalizations and comments

## The symmetry result

The Felli & Schneider curve is defined by

$$b_{\rm FS}(a) := rac{d(a_c - a)}{2\sqrt{(a_c - a)^2 + d - 1}} + a - a_c$$

#### Theorem

Let  $d \ge 2$  and  $p < 2^*$ . If either  $a \in [0, a_c)$  and b > 0, or a < 0 and  $b \ge b_{\rm FS}(a)$ , then the optimal functions for the Caffarelli-Kohn-Nirenberg inequalities are radially symmetric

< 回 > < 三 > < 三 >

Results on CKN inequalities Symmetry and symmetry breaking **The sharp result** Generalizations and comments

The Emden-Fowler transformation and the cylinder

▷ With an Emden-Fowler transformation, Caffarelli-Kohn-Nirenberg inequalities on the Euclidean space are equivalent to Gagliardo-Nirenberg inequalities on a cylinder

$$v(r,\omega) = r^{a-a_c} \varphi(s,\omega)$$
 with  $r = |x|$ ,  $s = -\log r$  and  $\omega = \frac{x}{r}$ 

With this transformation, the Caffarelli-Kohn-Nirenberg inequalities can be rewritten as

$$\|\partial_{s}\varphi\|_{\mathrm{L}^{2}(\mathcal{C})}^{2}+\|\nabla_{\omega}\varphi\|_{\mathrm{L}^{2}(\mathcal{C})}^{2}+\Lambda\|\varphi\|_{\mathrm{L}^{2}(\mathcal{C})}^{2}\geq\mu(\Lambda)\|\varphi\|_{\mathrm{L}^{p}(\mathcal{C})}^{2}\quad\forall\,\varphi\in\mathrm{H}^{1}(\mathcal{C})$$

where  $\Lambda := (a_c - a)^2$ ,  $\mathcal{C} = \mathbb{R} \times \mathbb{S}^{d-1}$  and the optimal constant  $\mu(\Lambda)$  is

$$\mu(\Lambda) = \frac{1}{\mathsf{C}_{a,b}} \quad \text{with} \quad a = a_c \pm \sqrt{\Lambda} \quad \text{and} \quad b = \frac{d}{p} \pm \sqrt{\Lambda}$$

- 4 同 6 4 日 6 4 日 6

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Generalizations and comments

イロト イポト イヨト イヨト

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

Generalized Caffarelli-Kohn-Nirenberg inequalities (CKN)

Let  $2^* = \infty$  if d = 1 or d = 2,  $2^* = 2d/(d-2)$  if  $d \ge 3$  and define

$$\vartheta(p,d):=\frac{d(p-2)}{2p}$$

[Caffarelli-Kohn-Nirenberg-84] Let  $d \ge 1$ . For any  $\theta \in [\vartheta(p, d), 1]$ , with  $p = \frac{2d}{d-2+2(b-a)}$ , there exists a positive constant  $C_{\text{CKN}}(\theta, p, a)$  such that

$$\left(\int_{\mathbb{R}^d} \frac{|u|^p}{|x|^{b\,p}}\,dx\right)^{\frac{2}{p}} \leq \mathsf{C}_{\mathrm{CKN}}(\theta,p,a)\left(\int_{\mathbb{R}^d} \frac{|\nabla u|^2}{|x|^{2\,a}}\,dx\right)^{\theta}\left(\int_{\mathbb{R}^d} \frac{|u|^2}{|x|^{2\,(a+1)}}\,dx\right)^{1-\theta}$$

In the radial case, with  $\Lambda = (a - a_c)^2$ , the best constant when the inequality is restricted to radial functions is  $C^*_{\text{CKN}}(\theta, p, a)$  and

$$\mathsf{C}_{\mathrm{CKN}}(\theta, \boldsymbol{p}, \boldsymbol{a}) \geq \mathsf{C}^*_{\mathrm{CKN}}(\theta, \boldsymbol{p}, \boldsymbol{a}) = \mathsf{C}^*_{\mathrm{CKN}}(\theta, \boldsymbol{p}) \, \Lambda^{\frac{\boldsymbol{p}-2}{2\boldsymbol{p}}-\theta}$$

$$\mathsf{C}^{*}_{\mathrm{CKN}}(\theta, p) = \left[\frac{2\pi^{d/2}}{\Gamma(d/2)}\right]^{2\frac{p-1}{p}} \left[\frac{(p-2)^{2}}{2+(2\theta-1)p}\right]^{\frac{p-2}{2p}} \left[\frac{2+(2\theta-1)p}{2p\theta}\right]^{\theta} \left[\frac{4}{p+2}\right]^{\frac{6-p}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{2}{p-2}+\frac{1}{2}\right)}\right]^{\frac{p-2}{2p}} \left[\frac{\Gamma\left(\frac{2$$

Caffarelli-Kohn-Nirenberg inequalities The proof of the symmetry result in 4 steps Two ingredients for the proof and some remarks Symmetry and symmetry breaking The sharp result Generalizations and comments

# Implementing the method of Catrina-Wang / Felli-Schneider

Among functions  $w \in H^1(\mathcal{C})$  which depend only on s, the minimum of

$$\mathcal{J}[w] := \int_{\mathcal{C}} \left( |\nabla w|^2 + \frac{1}{4} \left( d - 2 - 2 \, a \right)^2 |w|^2 \right) \, dx - \left[ \mathsf{C}^*(\theta, p, a) \right]^{-\frac{1}{\theta}} \, \frac{\left( \int_{\mathcal{C}} |w|^p \, dx \right)^{\frac{2}{p \, \theta}}}{\left( \int_{\mathcal{C}} |w|^2 \, dx \right)^{\frac{1-\theta}{\theta}}}$$

is achieved by 
$$\overline{w}(y) := \left[\cosh(\lambda s)\right]^{-\frac{2}{p-2}}, y = (s, \omega) \in \mathbb{R} \times \mathbb{S} = \mathcal{C}$$
 with  $\lambda := \frac{1}{4} \left(d - 2 - 2a\right) \left(p - 2\right) \sqrt{\frac{p+2}{2p\theta - (p-2)}}$  as a solution of  $\lambda^2 \left(p - 2\right)^2 w'' - 4w + 2p |w|^{p-2} w = 0$ 

Spectrum of  $\mathcal{L} := -\Delta + \kappa \overline{w}^{p-2} + \mu$  is given for  $\sqrt{1 + 4\kappa/\lambda^2} \ge 2j + 1$ by

$$\lambda_{i,j} = \mu + i \left( d + i - 2 \right) - \frac{\lambda^2}{4} \left( \sqrt{1 + \frac{4\kappa}{\lambda^2}} - (1 + 2j) \right)^2 \quad \forall i, j \in \mathbb{N}$$

• The eigenspace of  $\mathcal{L}$  corresponding to  $\lambda_{0,0}$  is generated by  $\overline{W}$ **Q** The eigenfunction  $\phi_{(1,0)}$  associated to  $\lambda_{1,0}$  is not radially symmetric and such that  $\int_{\mathcal{C}} \overline{w} \phi_{(1,0)} dx = 0$  and  $\int_{\mathcal{C}} \overline{w}^{p-1} \phi_{(1,0)} dx = 0$ **Q** If  $\lambda_{1,0} < 0$ , optimal functions for (CKN) cannot be radially =Symmetry breaking and sharp functional inequalities J. Dolbeault

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Parametric plot of $\mu \mapsto (\Lambda^{\theta}(\mu), J^{\theta}(\mu))$ for $p = 2.8, d = 5, \theta = 1$



J. Dolbeault Symmetry breaking and sharp functional inequalities

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Parametric plot of $\mu \mapsto (\Lambda^{\theta}(\mu), J^{\theta}(\mu))$ for $p = 2.8, d = 5, \theta = 0.8$



J. Dolbeault Symmetry breaking and sharp functional inequalities

Symmetry and symmetry breaking The sharp result Generalizations and comments

# Parametric plot of $\mu \mapsto (\Lambda^{\theta}(\mu), J^{\theta}(\mu))$ for p = 2.8, d = 5, $\theta = 0.72$



Symmetry breaking and sharp functional inequalities

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

## Enlargement for p = 2.8, d = 5, $\theta = 0.95$



Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Enlargement for p = 2.8, d = 5, $\theta = 0.72$



J. Dolbeault Symmetry breaking and sharp functional inequalities

э

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Critical case $\theta = \vartheta(p, d)$



<ロ> <同> <同> < 回> < 回>

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Parametric plot of $\mu \mapsto (\Lambda^{\theta}(\mu), J^{\theta}(\mu))$ for p = 3.15, d = 5, $\theta = 1$



J. Dolbeault Symmetry breaking and sharp functional inequalities

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Parametric plot of $\mu \mapsto (\Lambda^{\theta}(\mu), J^{\theta}(\mu))$ for p = 3.15, d = 5, $\theta = 0.95$



J. Dolbeault Symmetry breaking and sharp functional inequalities

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# Case p = 3.15, d = 5, $\theta = \vartheta(3.15, 5) \approx 0.9127$



イロト イポト イヨト イヨト

Results on CKN inequalities Symmetry and symmetry breaking The sharp result Generalizations and comments

# local and asymptotic criteria



A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# The main steps of the proof

- A change of variables: an equivalent inequality of Sbolev type
- The fast diffusion flow and the nonlinear Fisher information
- Proving the decay along the flow
- The justification of the integration by parts: decay estimates on the cylinder

- 4 同 6 4 日 6 4 日 6

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

## A change of variables

With  $(r = |x|, \omega = x/r) \in \mathbb{R}^+ \times \mathbb{S}^{d-1}$ , the Caffarelli-Kohn-Nirenberg inequality is

$$\left(\int_0^\infty \int_{\mathbb{S}^{d-1}} |v|^p r^{d-bp} \frac{dr}{r} d\omega\right)^{\frac{2}{p}} \leq \mathsf{C}_{a,b} \int_0^\infty \int_{\mathbb{S}^{d-1}} |\nabla v|^2 r^{d-2a} \frac{dr}{r} d\omega$$

Change of variables  $r \mapsto r^{\alpha}$ ,  $v(r, \omega) = w(r^{\alpha}, \omega)$ 

$$\begin{split} \alpha^{1-\frac{2}{p}} \left( \int_0^\infty \int_{\mathbb{S}^{d-1}} |w|^p r^{\frac{d-bp}{\alpha}} \frac{dr}{r} d\omega \right)^{\frac{2}{p}} \\ &\leq \mathsf{C}_{a,b} \int_0^\infty \int_{\mathbb{S}^{d-1}} \left( \alpha^2 \left| \frac{\partial w}{\partial r} \right|^2 + \frac{1}{r^2} \left| \nabla_\omega w \right|^2 \right) r^{\frac{d-2s-2}{\alpha}+2} \frac{dr}{r} d\omega \end{split}$$

Choice of  $\alpha$ 

$$n = \frac{d - b p}{\alpha} = \frac{d - 2 a - 2}{\alpha} + 2$$

Then  $p = \frac{2n}{n-2}$  is the critical Sobolev exponent associated with n

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# A Sobolev type inequality

The parameters  $\alpha$  and n vary in the ranges  $0 < \alpha < \infty$  and  $d < n < \infty$ and the *Felli-Schneider curve* in the  $(\alpha, n)$  variables is given by

$$\alpha = \sqrt{\frac{d-1}{n-1}} =: \alpha_{\rm FS}$$

With

$$\mathsf{D}w = \left( lpha \, \frac{\partial w}{\partial r}, \frac{1}{r} \, \nabla_{\omega} w \right) \,, \quad d\mu := r^{n-1} \, dr \, d\omega$$

the inequality becomes

$$\alpha^{1-\frac{2}{p}} \left( \int_{\mathbb{R}^d} |w|^p \, d\mu \right)^{\frac{2}{p}} \leq \mathsf{C}_{\mathsf{a},\mathsf{b}} \int_{\mathbb{R}^d} |\mathsf{D}w|^2 \, d\mu$$

#### Proposition

Let  $d\geq 2.$  Optimality is achieved by radial functions and  $C_{a,b}=C^{\star}_{a,b}$  if  $\alpha\leq \alpha_{\rm FS}$ 

Gagliardo-Nirenberg inequalities on general cylinders; similar

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

## Notations

When there is no ambiguity, we will omit the index  $\omega$  and from now on write that  $\nabla = \nabla_{\omega}$  denotes the gradient with respect to the angular variable  $\omega \in \mathbb{S}^{d-1}$  and that  $\Delta$  is the Laplace-Beltrami operator on  $\mathbb{S}^{d-1}$ . We define the self-adjoint operator  $\mathcal{L}$  by

$$\mathcal{L} w := -\mathsf{D}^* \mathsf{D} w = \alpha^2 w'' + \alpha^2 \frac{n-1}{r} w' + \frac{\Delta w}{r^2}$$

The fundamental property of  $\mathcal{L}$  is the fact that

$$\int_{\mathbb{R}^d} w_1 \mathcal{L} w_2 \, d\mu = - \int_{\mathbb{R}^d} \mathsf{D} w_1 \cdot \mathsf{D} w_2 \, d\mu \quad \forall w_1, w_2 \in \mathcal{D}(\mathbb{R}^d)$$

 $\triangleright$  Heuristics: we look for a monotonicity formula along a well chosen nonlinear flow, based on the analogy with the decay of the Fisher information along the fast diffusion flow in  $\mathbb{R}^d$ 

イロト イポト イラト イラト

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# Fisher information

Let 
$$u^{\frac{1}{2}-\frac{1}{n}} = |w| \iff u = |w|^p$$
,  $p = \frac{2n}{n-2}$ 

$$\mathcal{I}[u] := \int_{\mathbb{R}^d} u \left| \mathsf{Dp} \right|^2 d\mu, \quad \mathsf{p} = \frac{m}{1-m} u^{m-1} \quad \text{and} \quad m = 1 - \frac{1}{n}$$

Here  $\mathcal{I}$  is the *Fisher information* and p is the *pressure function* 

#### Proposition

With  $\Lambda = 4 \alpha^2 / (p-2)^2$  and for some explicit numerical constant  $\kappa$ , we have

$$\kappa \mu(\Lambda) = \inf \left\{ \mathcal{I}[u] \, : \, \|u\|_{\mathrm{L}^{1}(\mathbb{R}^{d}, d\mu)} = 1 \right\}$$

 $\rhd$  Optimal solutions solutions of the elliptic PDE) are (constrained) critical point of  $\mathcal I$ 

- 4 回 ト - 4 回 ト

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# The fast diffusion equation

$$\frac{\partial u}{\partial t} = \mathcal{L} u^m, \quad m = 1 - \frac{1}{n}$$

Barenblatt self-similar solutions

$$u_{\star}(t,r,\omega) = t^{-n} \left( c_{\star} + \frac{r^2}{2(n-1)\alpha^2 t^2} \right)^{-n}$$

#### Lemma

Barenblatt solutions realize the minimum of  $\mathcal{I}$  among radial functions:

$$\kappa \, \mu_{\star}(\Lambda) = \mathcal{I}[u_{\star}(t, \cdot)] \quad \forall \, t > 0$$

▷ Strategy: 1) prove that  $\frac{d}{dt}\mathcal{I}[u(t,\cdot)] \leq 0$ , 2) prove that  $\frac{d}{dt}\mathcal{I}[u(t,\cdot)] = 0$  means that  $u = u_*$  up to a time shift

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# Decay of the Fisher information along the flow ?

The pressure function 
$$\mathbf{p} = \frac{m}{1-m} u^{m-1}$$
 satisfies  
 $\frac{\partial \mathbf{p}}{\partial t} = \frac{1}{n} \mathbf{p} \mathcal{L} \mathbf{p} - |\mathsf{D}\mathbf{p}|^2$   
 $\mathcal{Q}[\mathbf{p}] := \frac{1}{2} \mathcal{L} |\mathsf{D}\mathbf{p}|^2 - \mathsf{D}\mathbf{p} \cdot \mathsf{D}\mathcal{L} \mathbf{p}$   
 $\mathcal{K}[\mathbf{p}] := \int_{\mathbb{R}^d} \left( \mathcal{Q}[\mathbf{p}] - \frac{1}{n} (\mathcal{L} \mathbf{p})^2 \right) \mathbf{p}^{1-n} d\mu$ 

#### Lemma

If u solves the weighted fast diffusion equation, then

$$\frac{d}{dt}\mathcal{I}[u(t,\cdot)] = -2(n-1)^{n-1}\mathcal{K}[p]$$

If u is a critical point, then  $\mathcal{K}[\mathbf{p}] = \mathbf{0}$  $\triangleright$  Boundary terms ! Regularity !

- 4 同 ト 4 ヨ ト 4 ヨ ト

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# Proving decay (1/2)

$$k[\mathbf{p}] := \mathcal{Q}(\mathbf{p}) - \frac{1}{n} (\mathcal{L} \mathbf{p})^2 = \frac{1}{2} \mathcal{L} |\mathsf{D}\mathbf{p}|^2 - \mathsf{D}\mathbf{p} \cdot \mathsf{D} \mathcal{L} \mathbf{p} - \frac{1}{n} (\mathcal{L} \mathbf{p})^2$$
$$k_{\mathfrak{M}}[\mathbf{p}] := \frac{1}{2} \Delta |\nabla \mathbf{p}|^2 - \nabla \mathbf{p} \cdot \nabla \Delta \mathbf{p} - \frac{1}{n-1} (\Delta \mathbf{p})^2 - (n-2) \alpha^2 |\nabla \mathbf{p}|^2$$

#### Lemma

Let  $n \neq 1$  be any real number,  $d \in \mathbb{N}$ ,  $d \geq 2$ , and consider a function  $p \in C^3((0,\infty) \times \mathfrak{M})$ , where  $(\mathfrak{M},g)$  is a smooth, compact Riemannian manifold. Then we have

$$k[\mathbf{p}] = \alpha^4 \left(1 - \frac{1}{n}\right) \left[\mathbf{p}'' - \frac{\mathbf{p}'}{r} - \frac{\Delta \mathbf{p}}{\alpha^2 (n-1) r^2}\right]^2 + 2 \alpha^2 \frac{1}{r^2} \left|\nabla \mathbf{p}' - \frac{\nabla \mathbf{p}}{r}\right|^2 + \frac{1}{r^4} k_{\mathfrak{M}}[\mathbf{p}]$$

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# Proving decay (2/2)

#### Lemma

Assume that 
$$d \ge 3$$
,  $n > d$  and  $\mathfrak{M} = \mathbb{S}^{d-1}$ . For some  $\zeta_{\star} > 0$  we have  

$$\int_{\mathbb{S}^{d-1}} k_{\mathfrak{M}}[p] p^{1-n} d\omega \ge (\lambda_{\star} - (n-2)\alpha^2) \int_{\mathbb{S}^{d-1}} |\nabla p|^2 p^{1-n} d\omega$$

$$+ \zeta_{\star} (n-d) \int_{\mathbb{S}^{d-1}} |\nabla p|^4 p^{1-n} d\omega$$

Proof based on the Bochner-Lichnerowicz-Weitzenböck formula

#### Corollary

Let  $d \geq 2$  and assume that  $\alpha \leq \alpha_{FS}$ . Then for any nonnegative function  $u \in L^1(\mathbb{R}^d)$  with  $\mathcal{I}[u] < +\infty$  and  $\int_{\mathbb{R}^d} u \, d\mu = 1$ , we have

 $\mathcal{I}[u] \geq \mathcal{I}_{\star}$ 

When  $\mathfrak{M} = \mathbb{S}^{d-1}$ ,  $\lambda_{\star} = (n-2) \frac{d-1}{n-1}$ 

🗇 🕨 🖉 🖢 🖌 🖉 🕨

A change of variables and a Sobolev type inequality Fisher information and fast diffusion flow Decay along the flow Decay estimates on the cylinder

# A perturbation argument

• If u is a critical point of  $\mathcal{I}$  under the mass constraint  $\int_{\mathbb{R}^d} u \, d\mu = 1$ , then

$$o(\varepsilon) = \mathcal{I}[u + \varepsilon \mathcal{L} u^m] - \mathcal{I}[u] = -2(n-1)^{n-1} \varepsilon \mathcal{K}[p] + o(\varepsilon)$$

because  $\varepsilon \, \mathcal{L} \, u^m$  is an admissible perturbation (formal). Indeed, we know that

$$\int_{\mathbb{R}^d} \left( u + \varepsilon \, \mathcal{L} \, u^m \right) d\mu = \int_{\mathbb{R}^d} u \, d\mu = 1$$

but positivity of  $u + \varepsilon \, \mathcal{L} \, u^m$  is an issue: compute

$$0 = D\mathcal{I}[u] \cdot \mathcal{L} u^m = -\mathcal{K}[p]$$

• Regularity issues (uniform decay of various derivatives up to order 3) and boundary terms

• If  $\alpha \leq \alpha_{\rm FS}$ , then  $\mathcal{K}[\mathbf{p}] = \mathbf{0}$  implies that  $u = u_{\star}$ 

# The justification of the integration by parts: decay estimates on the cylinder

After then Emden-Fowler transformation, a critical point satisfies the Euler-Lagrange equation

$$-\partial_s^2 \varphi - \Delta_\omega \varphi + \Lambda \varphi = \varphi^{p-1} \quad \text{in} \quad \mathcal{C} = \mathbb{R} \times \mathcal{M}$$

(up to a multiplication by a constant;  $\mathcal{M} = \mathbb{S}^{d-1}$  e.g.)

#### Proposition

For all 
$$(s,\omega) \in \mathcal{C}$$
, we have  $C_1 e^{-\sqrt{\Lambda} |s|} \le \varphi(s,\omega) \le C_2 e^{-\sqrt{\Lambda} |s|}$ 

$$|arphi'({f s},\omega)|\,,\;|arphi''({f s},\omega)|\,,\;|
abla arphi({f s},\omega)|\,,\;|\Delta\,arphi({f s},\omega)|\leq C_2\,e^{-\sqrt{\Lambda}\,|{f s}|}$$

and 
$$\begin{split} \int_{\mathfrak{M}} |\mathsf{p}'(r,\omega)|^2 \, d\, \mathsf{v}_g &\leq O(1), \ \int_{\mathfrak{M}} |\nabla \mathsf{p}(r,\omega)|^2 \, d\, \mathsf{v}_g &\leq O(r^2), \\ \int_{\mathfrak{M}} |\mathsf{p}''(r,\omega)|^2 \, d\, \mathsf{v}_g &\leq O(1/r^2) \\ \\ \int_{\mathfrak{M}} |\nabla \mathsf{p}'(r,\omega) - \frac{1}{r} \, \nabla \mathsf{p}(r,\omega)|^2 \, d\, \mathsf{v}_g &\leq O(1), \\ \int_{\mathfrak{M}} |\Delta \mathsf{p}(r,\omega)|^2 \, d\, \mathsf{v}_r &\leq O(1/r^2) \\ \end{bmatrix}$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Two ingredients for the proof

▷ Rényi entropy powers and fast diffusion▷ Flows on the sphere

・ロン ・四と ・ヨン ・ヨン

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Rényi entropy powers and fast diffusion

▷ Rényi entropy powers, the entropy approach without rescaling: [Savaré, Toscani]: scalings, nonlinearity and a concavity property inspired by information theory

▷ faster rates of convergence: [Carrillo, Toscani], [JD, Toscani]

- 4 同 6 4 日 6 4 日 6

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

## The fast diffusion equation in original variables

Consider the nonlinear diffusion equation in  $\mathbb{R}^d,\,d\geq 1$ 

$$\frac{\partial u}{\partial t} = \Delta u^m$$

with initial datum  $u(x, t = 0) = u_0(x) \ge 0$  such that  $\int_{\mathbb{R}^d} u_0 \, dx = 1$  and  $\int_{\mathbb{R}^d} |x|^2 u_0 \, dx < +\infty$ . The large time behavior of the solutions is governed by the source-type Barenblatt solutions

$$\mathcal{U}_{\star}(t,x) \coloneqq rac{1}{ig(\kappa \, t^{1/\mu}ig)^d} \, \mathcal{B}_{\star}ig(rac{x}{\kappa \, t^{1/\mu}}ig)$$

where

$$\mu := 2 + d(m-1), \quad \kappa := \left|\frac{2 \mu m}{m-1}\right|^{1/\mu}$$

and  $\mathcal{B}_{\star}$  is the Barenblatt profile

$$\mathcal{B}_{\star}(x) := egin{cases} \left( C_{\star} - |x|^2 
ight)_+^{1/(m-1)} & ext{if } m > 1 \ \left( C_{\star} + |x|^2 
ight)^{1/(m-1)} & ext{if } m < 1 \end{cases}$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# The Rényi entropy power F

The entropy is defined by

$$\Xi := \int_{\mathbb{R}^d} u^m \, dx$$

and the Fisher information by

$$I := \int_{\mathbb{R}^d} u |\nabla p|^2 dx$$
 with  $p = \frac{m}{m-1} u^{m-1}$ 

If  $\boldsymbol{u}$  solves the fast diffusion equation, then

$$\mathsf{E}' = (1-m)\mathsf{I}$$

To compute  ${\mathsf I}',$  we will use the fact that

$$\frac{\partial p}{\partial t} = (m-1) p \Delta p + |\nabla p|^2$$

$$F := E^{\sigma} \quad \text{with} \quad \sigma = \frac{\mu}{d(1-m)} = 1 + \frac{2}{1-m} \left(\frac{1}{d} + m - 1\right) = \frac{2}{d} \frac{1}{1-m} - 1$$
has a linear growth asymptotically as  $t \to \pm \infty$ 

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# The concavity property

#### Theorem

[Toscani-Savaré] Assume that  $m \ge 1 - \frac{1}{d}$  if d > 1 and m > 0 if d = 1. Then F(t) is increasing,  $(1 - m) F''(t) \le 0$  and

$$\lim_{t \to +\infty} \frac{1}{t} \mathsf{F}(t) = (1 - m) \sigma \lim_{t \to +\infty} \mathsf{E}^{\sigma - 1} \mathsf{I} = (1 - m) \sigma \mathsf{E}_{\star}^{\sigma - 1} \mathsf{I}_{\star}$$

[Dolbeault-Toscani] The inequality

$$\mathsf{E}^{\sigma-1}\,\mathsf{I}\geq\mathsf{E}_\star^{\sigma-1}\,\mathsf{I}_\star$$

is equivalent to the Gagliardo-Nirenberg inequality

$$\|\nabla w\|_{\mathrm{L}^2(\mathbb{R}^d)}^{\theta} \|w\|_{\mathrm{L}^{q+1}(\mathbb{R}^d)}^{1-\theta} \geq \mathsf{C}_{\mathrm{GN}} \|w\|_{\mathrm{L}^{2q}(\mathbb{R}^d)}$$

if  $1 - \frac{1}{d} \le m < 1$ . Hint:  $u^{m-1/2} = \frac{w}{\|w\|_{L^{2q}(\mathbb{R}^d)}}, \ q = \frac{1}{2m-1}$ 

The proof

#### Lemma

If 
$$u$$
 solves  $\frac{\partial u}{\partial t} = \Delta u^m$  with  $\frac{1}{d} \le m < 1$ , then  

$$I' = \frac{d}{dt} \int_{\mathbb{R}^d} u |\nabla p|^2 dx = -2 \int_{\mathbb{R}^d} u^m \left( \|D^2 p\|^2 + (m-1) (\Delta p)^2 \right) dx$$

Rényi entropy powers and fast diffusion

$$\|\mathbf{D}^2 \mathbf{p}\|^2 = \frac{1}{d} (\Delta \mathbf{p})^2 + \left\| \mathbf{D}^2 \mathbf{p} - \frac{1}{d} \Delta \mathbf{p} \operatorname{Id} \right\|^2$$

$$\frac{1}{\sigma(1-m)} \mathsf{E}^{2-\sigma} (\mathsf{E}^{\sigma})'' = (1-m)(\sigma-1) \left( \int_{\mathbb{R}^d} u |\nabla \mathsf{p}|^2 \, dx \right)^2 - 2 \left( \frac{1}{d} + m - 1 \right) \int_{\mathbb{R}^d} u^m \, dx \int_{\mathbb{R}^d} u^m \, (\Delta \mathsf{p})^2 \, dx - 2 \int_{\mathbb{R}^d} u^m \, dx \int_{\mathbb{R}^d} u^m \left\| \mathsf{D}^2 \mathsf{p} - \frac{1}{\sigma} \frac{1}{d} \Delta \mathsf{p} \operatorname{Id} \right\|_{\mathbb{R}^d}^2 \, dx = \mathcal{O}(\mathbf{q})$$
J. Dolbeaut

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Flows on the sphere

 $\triangleright$  The heat flow introduced by D. Bakry and M. Emery (*carré du champ* method) does not cover all exponents up to the critical one

[Bakry, Emery, 1984] [Bidault-Véron, Véron, 1991], [Bakry, Ledoux, 1996] [Demange, 2008][JD, Esteban, Loss, 2014 & 2015]

< 回 ト く ヨ ト く ヨ ト

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# The interpolation inequalities

On the  $d\mbox{-dimensional sphere, let us consider the interpolation inequality}$ 

$$\|\nabla u\|_{\mathrm{L}^2(\mathbb{S}^d)}^2 + \frac{d}{p-2} \|u\|_{\mathrm{L}^2(\mathbb{S}^d)}^2 \geq \frac{d}{p-2} \|u\|_{\mathrm{L}^p(\mathbb{S}^d)}^2 \quad \forall \, u \in \mathrm{H}^1(\mathbb{S}^d, d\mu)$$

where the measure  $d\mu$  is the uniform probability measure on  $\mathbb{S}^d \subset \mathbb{R}^{d+1}$  corresponding to the measure induced by the Lebesgue measure on  $\mathbb{R}^{d+1}$ , and the exposant  $p \geq 1$ ,  $p \neq 2$ , is such that

$$p \leq 2^* := \frac{2d}{d-2}$$

if  $d \ge 3$ . We adopt the convention that  $2^* = \infty$  if d = 1 or d = 2. The case p = 2 corresponds to the logarithmic Sobolev inequality

$$\|\nabla u\|_{\mathrm{L}^{2}(\mathbb{S}^{d})}^{2} \geq \frac{d}{2} \int_{\mathbb{S}^{d}} |u|^{2} \log\left(\frac{|u|^{2}}{\|u\|_{\mathrm{L}^{2}(\mathbb{S}^{d})}^{2}}\right) dv_{g} \quad \forall u \in \mathrm{H}^{1}(\mathbb{S}^{d}, d\mu) \setminus \{0\}$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# The Bakry-Emery method

Entropy functional

1

$$\mathcal{E}_{p}[\rho] := \frac{1}{p-2} \left[ \int_{\mathbb{S}^{d}} \rho^{\frac{2}{p}} dv_{g} - \left( \int_{\mathbb{S}^{d}} \rho dv_{g} \right)^{\frac{2}{p}} \right] \quad \text{if} \quad p \neq 2$$
$$\mathcal{E}_{2}[\rho] := \int_{\mathbb{S}^{d}} \rho \log \left( \frac{\rho}{\|\rho\|_{L^{1}(\mathbb{S}^{d})}} \right) dv_{g}$$

Fisher information functional

$$\mathcal{I}_{
ho}[
ho] := \int_{\mathbb{S}^d} |
abla 
ho^{rac{1}{
ho}}|^2 \; d \, v_g$$

Bakry-Emery (carré du champ): use the heat flow

$$\frac{\partial \rho}{\partial t} = \Delta \rho$$

where  $\Delta$  denotes the Laplace-Beltrami operator on  $\mathbb{S}^d$ , and compute

$$\frac{d}{dt}\mathcal{E}_{p}[\rho] = -\mathcal{I}_{p}[\rho] \quad \text{and} \quad \frac{d}{dt}\mathcal{I}_{p}[\rho] \leq -d\mathcal{I}_{p}[\rho]$$

Rénvi entropy powers and fast diffusion Flows on the sphere

# The evolution under the fast diffusion flow

To overcome the limitation  $p \leq 2^{\#}$ , one can consider a nonlinear diffusion of fast diffusion / porous medium type

$$\frac{\partial \rho}{\partial t} = \Delta \rho^m \,. \tag{1}$$

[Demange], [JD, Esteban, Kowalczyk, Loss]: for any  $p \in [1, 2^*]$ 

$$\mathcal{K}_{\rho}[\rho] := rac{d}{dt} \Big( \mathcal{I}_{\rho}[\rho] - d \, \mathcal{E}_{\rho}[\rho] \Big) \leq 0 \,,$$



Symmetry breaking and sharp functional inequalities

э

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Sobolev's inequality

The stereographic projection of  $\mathbb{S}^d \subset \mathbb{R}^d \times \mathbb{R} \ni (\rho \phi, z)$  onto  $\mathbb{R}^d$ : to  $\rho^2 + z^2 = 1$ ,  $z \in [-1, 1]$ ,  $\rho \ge 0$ ,  $\phi \in \mathbb{S}^{d-1}$  we associate  $x \in \mathbb{R}^d$  such that  $r = |x|, \phi = \frac{x}{|x|}$ 

$$z = \frac{r^2 - 1}{r^2 + 1} = 1 - \frac{2}{r^2 + 1}, \quad \rho = \frac{2r}{r^2 + 1}$$

and transform any function u on  $\mathbb{S}^d$  into a function v on  $\mathbb{R}^d$  using

$$u(y) = \left(\frac{r}{\rho}\right)^{\frac{d-2}{2}} v(x) = \left(\frac{r^2+1}{2}\right)^{\frac{d-2}{2}} v(x) = (1-z)^{-\frac{d-2}{2}} v(x)$$

 $\blacksquare \ p=2^*, \, \mathsf{S}_d=\frac{1}{4}\,d\,(d-2)\,|\mathbb{S}^d|^{2/d}\colon$  Euclidean Sobolev inequality

$$\int_{\mathbb{R}^d} |\nabla v|^2 \, dx \ge \mathsf{S}_d \left[ \int_{\mathbb{R}^d} |v|^{\frac{2d}{d-2}} \, dx \right]^{\frac{d-2}{d}} \quad \forall \, v \in \mathcal{D}^{1,2}(\mathbb{R}^d)$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

## Schwarz symmetrization and the ultraspherical setting

$$(\xi_0, \xi_1, \dots \xi_d) \in \mathbb{S}^d, \, \xi_d = z, \, \sum_{i=0}^d |\xi_i|^2 = 1 \, [\text{Smets-Willem}]$$

#### Lemma

Up to a rotation, any minimizer of  ${\mathcal Q}$  depends only on  $\xi_d=z$ 

• Let 
$$d\sigma(\theta) := \frac{(\sin \theta)^{d-1}}{Z_d} d\theta$$
,  $Z_d := \sqrt{\pi} \frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d+1}{2})}$ :  $\forall v \in \mathrm{H}^1([0,\pi], d\sigma)$ 

$$\frac{p-2}{d}\int_0^\pi |v'(\theta)|^2 \ d\sigma + \int_0^\pi |v(\theta)|^2 \ d\sigma \ge \left(\int_0^\pi |v(\theta)|^p \ d\sigma\right)^{\frac{2}{p}}$$

• Change of variables  $z = \cos \theta$ ,  $v(\theta) = f(z)$ 

$$\frac{p-2}{d} \int_{-1}^{1} |f'|^2 \nu \, d\nu_d + \int_{-1}^{1} |f|^2 \, d\nu_d \ge \left(\int_{-1}^{1} |f|^p \, d\nu_d\right)^{\frac{2}{p}}$$

where  $\nu_d(z) dz = d\nu_d(z) := Z_d^{-1} \nu^{\frac{d}{2}-1} dz, \ \nu(z) := 1 - z^2$ 

Caffarelli-Kohn-Nirenberg inequalities The proof of the symmetry result in 4 steps Two ingredients for the proof and some remarks

Rénvi entropy powers and fast diffusion Flows on the sphere

## The ultraspherical operator

With  $d\nu_d = Z_d^{-1} \nu^{\frac{d}{2}-1} dz$ ,  $\nu(z) := 1 - z^2$ , consider the space  $L^{2}((-1,1), d\nu_{d})$  with scalar product

$$\langle f_1, f_2 \rangle = \int_{-1}^1 f_1 f_2 d\nu_d, \quad \|f\|_{\mathrm{L}^p(\mathbb{S}^d)} = \left(\int_{-1}^1 f^p d\nu_d\right)^{\frac{1}{p}}$$

The self-adjoint *ultraspherical* operator is

$$\mathcal{L} \, f := (1 - z^2) \, f'' - d \, z \, f' = 
u \, f'' + rac{d}{2} \, 
u' \, f'$$

which satisfies  $\langle f_1, \mathcal{L} f_2 \rangle = - \int_{-1}^{1} f'_1 f'_2 \nu d\nu_d$ 

#### Proposition

Let  $p \in [1, 2) \cup (2, 2^*]$ ,  $d \ge 1$ 

$$-\langle f, \mathcal{L} f \rangle = \int_{-1}^{1} |f'|^2 \ \nu \ d\nu_d \ge d \ \frac{\|f\|_{\mathrm{L}^p(\mathbb{S}^d)}^2 - \|f\|_{\mathrm{L}^2(\mathbb{S}^d)}^2}{p-2} \quad \forall f \in \mathrm{H}^1([-1,1], d\nu_d)$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

## Heat flow and the Bakry-Emery method

With 
$$g = f^{p}$$
, *i.e.*  $f = g^{\alpha}$  with  $\alpha = 1/p$ 

(Ineq.) 
$$-\langle f, \mathcal{L} f \rangle = -\langle g^{\alpha}, \mathcal{L} g^{\alpha} \rangle =: \mathcal{I}[g] \ge d \frac{\|g\|_{\mathrm{L}^{1}(\mathbb{S}^{d})}^{2\alpha} - \|g^{2\alpha}\|_{\mathrm{L}^{1}(\mathbb{S}^{d})}}{p-2} =: \mathcal{J}$$

Heat flow

$$rac{\partial g}{\partial t} = \mathcal{L} g$$

~

$$\frac{d}{dt} \|g\|_{\mathrm{L}^{1}(\mathbb{S}^{d})} = 0, \quad \frac{d}{dt} \|g^{2\alpha}\|_{\mathrm{L}^{1}(\mathbb{S}^{d})} = -2(p-2)\langle f, \mathcal{L}f \rangle = 2(p-2) \int_{-1}^{1} |f'|^{2} \nu$$

which finally gives

$$\frac{d}{dt}\mathcal{F}[g(t,\cdot)] = -\frac{d}{p-2}\frac{d}{dt}\|g^{2\alpha}\|_{\mathrm{L}^{1}(\mathbb{S}^{d})} = -2\,d\,\mathcal{I}[g(t,\cdot)]$$

Ineq.  $\iff \frac{d}{dt} \mathcal{F}[g(t,\cdot)] \leq -2 d \mathcal{F}[g(t,\cdot)] \iff \frac{d}{dt} \mathcal{I}[g(t,\cdot)] \leq -2 d \mathcal{I}[g(t,\cdot)]$ 

< 回 > < 三 > < 三 >

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

The equation for  $g = f^p$  can be rewritten in terms of f as

$$rac{\partial f}{\partial t} = \mathcal{L} f + (p-1) rac{|f'|^2}{f} 
u$$

$$-\frac{1}{2}\frac{d}{dt}\int_{-1}^{1}|f'|^{2}\nu d\nu_{d} = \frac{1}{2}\frac{d}{dt}\langle f,\mathcal{L}f\rangle = \langle \mathcal{L}f,\mathcal{L}f\rangle + (p-1)\langle \frac{|f'|^{2}}{f}\nu,\mathcal{L}f\rangle$$

$$\frac{d}{dt}\mathcal{I}[g(t,\cdot)] + 2 d\mathcal{I}[g(t,\cdot)] = \frac{d}{dt} \int_{-1}^{1} |f'|^2 \nu \, d\nu_d + 2 d \int_{-1}^{1} |f'|^2 \nu \, d\nu_d$$
$$= -2 \int_{-1}^{1} \left( |f''|^2 + (p-1) \frac{d}{d+2} \frac{|f'|^4}{f^2} - 2(p-1) \frac{d-1}{d+2} \frac{|f'|^2 f''}{f} \right) \nu^2 \, d\nu_d$$

is nonpositive if

$$|f''|^2 + (p-1)\frac{d}{d+2}\frac{|f'|^4}{f^2} - 2(p-1)\frac{d-1}{d+2}\frac{|f'|^2f''}{f}$$

is pointwise nonnegative, which is granted if

$$\left[ (p-1)\frac{d-1}{d+2} \right]^2 \le (p-1)\frac{d}{d+2} \iff p \le \frac{2d^2+1}{(d-1)^2} = 2^{\#} < \frac{2d}{d-2} = 2^{*}$$

J. Dolbeault Symmetry b

Symmetry breaking and sharp functional inequalities

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

### ... up to the critical exponent: a proof in two slides

$$\left[\frac{d}{dz},\mathcal{L}\right] u = (\mathcal{L} u)' - \mathcal{L} u' = -2 z u'' - d u'$$

$$\int_{-1}^{1} (\mathcal{L} u)^{2} d\nu_{d} = \int_{-1}^{1} |u''|^{2} \nu^{2} d\nu_{d} + d \int_{-1}^{1} |u'|^{2} \nu d\nu_{d}$$
$$\int_{-1}^{1} (\mathcal{L} u) \frac{|u'|^{2}}{u} \nu d\nu_{d} = \frac{d}{d+2} \int_{-1}^{1} \frac{|u'|^{4}}{u^{2}} \nu^{2} d\nu_{d} - 2 \frac{d-1}{d+2} \int_{-1}^{1} \frac{|u'|^{2} u''}{u} \nu^{2} d\nu_{d}$$

On (-1, 1), let us consider the *porous medium (fast diffusion)* flow

$$u_t = u^{2-2\beta} \left( \mathcal{L} \, u + \kappa \, \frac{|u'|^2}{u} \, \nu \right)$$

If  $\kappa = \beta (p-2) + 1$ , the L<sup>p</sup> norm is conserved

$$\frac{d}{dt} \int_{-1}^{1} u^{\beta p} \, d\nu_d = \beta \, p \, (\kappa - \beta \, (p - 2) - 1) \int_{-1}^{1} u^{\beta (p - 2)} \, |u'|^2 \, \nu \, d\nu_d = 0$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

$$f = u^{\beta}, \, \|f'\|_{\mathrm{L}^{2}(\mathbb{S}^{d})}^{2} + \frac{d}{p-2} \, \left(\|f\|_{\mathrm{L}^{2}(\mathbb{S}^{d})}^{2} - \|f\|_{\mathrm{L}^{p}(\mathbb{S}^{d})}^{2}\right) \geq 0 \, ?$$

$$egin{aligned} \mathcal{A} &:= \int_{-1}^1 |u''|^2 \, 
u^2 \, d
u_d - 2 \, rac{d-1}{d+2} \, (\kappa+eta-1) \int_{-1}^1 u'' \, rac{|u'|^2}{u} \, 
u^2 \, d
u_d \ &+ \left[\kappa \, (eta-1) + \, rac{d}{d+2} \, (\kappa+eta-1) 
ight] \int_{-1}^1 rac{|u'|^4}{u^2} \, 
u^2 \, d
u_d \end{aligned}$$

 $\mathcal{A}$  is nonnegative for some  $\beta$  if

$$\frac{8\,d^2}{(d+2)^2}\,(p-1)\,(2^*-p)\geq 0$$

 $\mathcal{A}$  is a sum of squares if  $p \in (2, 2^*)$  for an arbitrary choice of  $\beta$  in a certain interval (depending on p and d)

$$\mathcal{A} = \int_{-1}^{1} \left| u'' - \frac{p+2}{6-p} \frac{|u'|^2}{u} \right|^2 \nu^2 \ d\nu_d \ge 0 \quad \text{if } p = 2^* \text{ and } \beta = \frac{4}{6-p}$$

イロト イポト イヨト イヨト

-

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# The rigidity point of view

Which computation have we done ?  $u_t = u^{2-2\beta} \left( \mathcal{L} u + \kappa \frac{|u'|^2}{u} \nu \right)$ 

$$-\mathcal{L} u - (\beta - 1) \frac{|u'|^2}{u} \nu + \frac{\lambda}{p - 2} u = \frac{\lambda}{p - 2} u^{\kappa}$$

Multiply by  $\mathcal{L}\, u$  and integrate

$$\dots \int_{-1}^{1} \mathcal{L} u u^{\kappa} d\nu_{d} = -\kappa \int_{-1}^{1} u^{\kappa} \frac{|u'|^2}{u} d\nu_{d}$$

Multiply by  $\kappa \frac{|u'|^2}{u}$  and integrate

$$\dots = +\kappa \int_{-1}^{1} u^{\kappa} \frac{|u'|^2}{u} d\nu_d$$

The two terms cancel and we are left only with the two-homogenous terms

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Constraints and improvements

イロト イポト イヨト イヨト

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Integral constraints

#### Proposition

For any  $p \in (2, 2^{\#})$ , the inequality

$$\begin{split} \int_{-1}^{1} |f'|^2 \ \nu \ d\nu_d + \frac{\lambda}{p-2} \, \|f\|_2^2 &\geq \frac{\lambda}{p-2} \, \|f\|_p^2 \\ &\forall f \in \mathrm{H}^1((-1,1), d\nu_d) \ \text{s.t.} \ \int_{-1}^{1} z \, |f|^p \ d\nu_d = 0 \end{split}$$

holds with

$$\lambda \geq d + rac{(d-1)^2}{d(d+2)} \left(2^\# - p
ight) \left(\lambda^\star - d
ight)$$

・ロン ・四と ・ヨン ・ヨン

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

# Antipodal symmetry

With the additional restriction of antipodal symmetry, that is

$$u(-x) = u(x) \quad \forall x \in \mathbb{S}^d$$

#### Theorem

If  $p \in (1,2) \cup (2,2^*)$ , we have

$$\int_{\mathbb{S}^d} |\nabla u|^2 \, d\, \mathsf{v}_g \geq \frac{d}{p-2} \left[ 1 + \frac{(d^2-4)(2^*-p)}{d(d+2)+p-1} \right] \left( \|u\|_{\mathrm{L}^p(\mathbb{S}^d)}^2 - \|u\|_{\mathrm{L}^2(\mathbb{S}^d)}^2 \right)$$

for any  $u \in H^1(\mathbb{S}^d, d\mu)$  with antipodal symmetry. The limit case p = 2 corresponds to the improved logarithmic Sobolev inequality

$$\int_{\mathbb{S}^d} |\nabla u|^2 \,\, d\, v_g \geq \frac{d}{2} \frac{(d+3)^2}{(d+1)^2} \int_{\mathbb{S}^d} |u|^2 \,\, \log\left(\frac{|u|^2}{\|u\|_{\mathrm{L}^2(\mathbb{S}^d)}^2}\right) \, d\, v_g$$

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

## The larger picture: branches of antipodal solutions



伺 ト イヨト イヨト

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

## The optimal constant in the antipodal framework



Numerical computation of the optimal constant when d = 5 and  $1 \le p \le 10/3 \approx 3.33$ . The limiting value of the constant is numerically found to be equal to  $\lambda_{\star} = 2^{1-2/p} d \approx 6.59754$  with d = 5 and p = 10/3

Rényi entropy powers and fast diffusion Flows on the sphere Constraints and improvements

These slides can be found at

# $\label{eq:http://www.ceremade.dauphine.fr/~dolbeaul/Conferences/ $$ $$ $$ $$ $$ Lectures $$$

# Thank you for your attention !

-