

INEQUALITIES INVOLVING AHARONOV-BOHM MAGNETIC POTENTIALS IN DIMENSIONS 2 AND 3

DENIS BONHEURE

*Département de mathématique, Faculté des sciences, Université Libre de Bruxelles,
Campus Plaine CP 213, Bld du Triomphe, B-1050 Brussels, Belgium*

E-mail: denis.bonheure@ulb.be

JEAN DOLBEAULT

*CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine,
Place de Lattre de Tassigny, 75775 Paris 16, France*

E-mail: dolbeaul@ceremade.dauphine.fr

MARIA J. ESTEBAN

*CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine,
Place de Lattre de Tassigny, 75775 Paris 16, France*

E-mail: esteban@ceremade.dauphine.fr

ARI LAPTEV

*Department of Mathematics, Imperial College London,
Huxley Building, 180 Queen's Gate, London SW7 2AZ, UK*

E-mail: a.laptev@imperial.ac.uk

MICHAEL LOSS

*School of Mathematics, Skiles Building, Georgia Institute of Technology,
Atlanta GA 30332-0160, USA*

E-mail: loss@math.gatech.edu

This paper is devoted to a collection of results on nonlinear interpolation inequalities associated with Schrödinger operators involving Aharonov-Bohm magnetic potentials, and to some consequences. As symmetry plays an important role for establishing optimality results, we shall consider various cases corresponding to a circle, a two-dimensional sphere or a two-dimensional torus, and also the Euclidean spaces of dimensions two and three. Most of the results are new and we put the emphasis on the methods, as very little is known on symmetry, rigidity and optimality in presence of a magnetic field. The most spectacular applications are new magnetic Hardy inequalities in dimensions 2 and 3.

Keywords: Aharonov-Bohm magnetic potential; radial symmetry; cylindrical symmetry; symmetry breaking; magnetic Hardy inequality; magnetic interpolation inequality; optimal constants; magnetic Schrödinger operator; magnetic Keller-Lieb-Thirring inequality; magnetic rings.

Mathematics Subject Classification 2010: Primary: 81V10, 81Q10, 35Q60.
Secondary: 35Q40, 49K30, 35P30, 35J10, 35Q55, 46N50, 35J20.

1. Introduction

Variational problems involving magnetic fields play a peculiar role in the calculus of variations. It is fair to say that there are no simple physical intuitions that may serve as a guide and one has to extract information through exact computations, such as in the case of the Landau Hamiltonian. A case in point are the *symmetry properties of optimizers* where in the absence of magnetic fields one has fairly robust methods to prove this. The isoperimetric inequality is one of the main sources of intuition; rearrangements are essentially different versions of the isoperimetric problem. This is certainly not the case in the presence of magnetic fields and there are very few results in this direction.

An example is the work of Avron, Herbst and Simon [2] who proved that the ground state of the hydrogenic atom in a constant magnetic field has cylindrical symmetry. The proof is quite involved. Another result that comes to mind is by Erdős [20] who proved the equivalent of the Faber-Krahn inequality for the Schrödinger operator with a constant magnetic field and with a Dirichlet boundary condition in a domain. The disk yields the smallest ground state energy among domains of equal area. Again, the proof is quite involved and some arguments are tailored for a linear setting. In this connection one should mention the recent result of Bonheure, Nys and van Schaftingen [8] who showed perturbatively that in some nonlinear variational problem involving a small constant magnetic field the minimizers inherit the symmetry of the problem. Besides the constant magnetic field case another class of physically relevant variational problems involve Aharonov-Bohm magnetic fields and the purpose of this article is to give an up-to-date account of our knowledge.

The *Aharonov-Bohm effect* states that the wave function of a charged quantum particle passing by a thin magnetic solenoid experiences a phase shift. This, despite that there is no apparent interaction with the solenoid except through the interaction of the particle with the ‘unphysical’ vector potential. This prediction was made originally by Ehrenberg and Siday in 1949 (see [18]) and then again in 1959 by Aharonov-Bohm (see [1]) and we will stay with the custom of calling it the Aharonov-Bohm effect. It cannot be explained in terms of classical mechanics, but was nevertheless experimentally verified (see [4]). It counts as one of the important quantum mechanical effects.

Another question one may pose is the influence of the Aharonov-Bohm potential on the energies of systems, say of a particle in a potential interacting with the solenoid. It is relatively straightforward to write the Hamiltonian for this situation and one may ask for the effect of the Aharonov-Bohm field on the *spectrum* of the Hamiltonian. One fruitful approach is to relate the ground state energy of a quantum mechanical particle in an external potential to the minimization of a nonlinear dual variational problem. This also works in the presence of magnetic fields and in particular with the Aharonov-Bohm field. In this context this leads to nonlinear versions of Hardy type inequalities, or simply *nonlinear interpolation inequalities*,

which are the dual versions of the *Keller-Lieb-Thirring spectral estimates*. In various symmetry settings, we are interested in getting as much insight as possible about the best constants in the inequalities and also about the qualitative properties of their extremal functions. Indeed, in many cases studying the symmetry properties of those extremal functions allows to get very accurate, and sometimes even sharp, estimates for the best constants in the inequalities (see for instance [7]).

On the Euclidean space \mathbb{R}^d , the *magnetic Laplacian* is defined via a *magnetic potential* \mathbf{A} by

$$-\Delta_{\mathbf{A}} \psi = -\Delta \psi - 2i \mathbf{A} \cdot \nabla \psi + |\mathbf{A}|^2 \psi - i (\operatorname{div} \mathbf{A}) \psi.$$

We consider the case of dimensions $d = 2$ and $d = 3$. The *magnetic field* is $\mathbf{B} = \operatorname{curl} \mathbf{A}$. The quadratic form associated with $-\Delta_{\mathbf{A}}$ is given by $\int_{\mathbb{R}^d} |\nabla_{\mathbf{A}} \psi|^2 dx$ and well defined for all functions in the space

$$H_{\mathbf{A}}^1(\mathbb{R}^d) := \left\{ \psi \in L^2(\mathbb{R}^d) : \nabla_{\mathbf{A}} \psi \in L^2(\mathbb{R}^d) \right\}$$

where the *magnetic gradient* takes the form

$$\nabla_{\mathbf{A}} := \nabla + i \mathbf{A}.$$

The *Aharonov-Bohm magnetic field* can be considered as a singular measure supported in the set $x_1 = x_2 = 0$, where $(x_i)_{i=1}^d$ is a system of cartesian coordinates. The magnetic potential is defined as follows.

- On \mathbb{R}^2 , let us consider *polar coordinates* (r, θ) such that

$$r = |x| = \sqrt{x_1^2 + x_2^2} \quad \text{and} \quad r e^{i\theta} = x_1 + i x_2$$

and the *Aharonov-Bohm magnetic potential*

$$\mathbf{A} = \frac{a}{r^2} (x_2, -x_1) = \frac{a}{r} \mathbf{e}_{\theta} \quad (1.1)$$

where a is a real constant and $\{\mathbf{e}_r, \mathbf{e}_{\theta}\}$, with $\mathbf{e}_r = \frac{x}{r}$, denotes the orthogonal basis associated with our polar coordinates. The magnetic gradient and the magnetic Laplacian are explicitly given by

$$\nabla_{\mathbf{A}} = \left(\frac{\partial}{\partial r}, \frac{1}{r} \left(\frac{\partial}{\partial \theta} - i a \right) \right), \quad -\Delta_{\mathbf{A}} = -\frac{\partial^2}{\partial r^2} - \frac{1}{r} \frac{\partial}{\partial r} - \frac{1}{r^2} \left(\frac{\partial}{\partial \theta} - i a \right)^2.$$

- On \mathbb{R}^3 , let us consider *cylindrical coordinates* (ρ, θ, z) where

$$\rho = \sqrt{x_1^2 + x_2^2}, \quad \rho e^{i\theta} = x_1 + i x_2 \quad \text{and} \quad z = x_3$$

and the *Aharonov-Bohm magnetic potential*

$$\mathbf{A} = \frac{a}{\rho^2} (x_2, -x_1, 0). \quad (1.2)$$

The magnetic gradient and the magnetic Laplacian are explicitly given by

$$\nabla_{\mathbf{A}} = \left(\frac{\partial}{\partial \rho}, \frac{1}{\rho} \left(\frac{\partial}{\partial \theta} - i a \right), \frac{\partial}{\partial z} \right), \quad -\Delta_{\mathbf{A}} = -\frac{\partial^2}{\partial \rho^2} - \frac{1}{\rho} \frac{\partial}{\partial \rho} - \frac{1}{\rho^2} \left(\frac{\partial}{\partial \theta} - i a \right)^2 - \frac{\partial^2}{\partial z^2}.$$

We shall also consider Aharonov-Bohm type magnetic potentials on compact manifolds, namely on the circle, the sphere and the torus. The expression of the magnetic potential will be given case by case.

This paper is intended to provide a general overview of mathematical results and methods concerning various functional inequalities involving Aharonov-Bohm magnetic fields:

- Magnetic ground state energy estimates.
- Nonlinear magnetic interpolation inequalities.
- Rigidity results for optimal functions.
- Magnetic Keller-Lieb-Thirring inequalities.
- Magnetic Hardy inequalities.

Except Hardy inequalities, all above inequalities will be considered on the circle \mathbb{S}^1 , on the two-dimensional sphere \mathbb{S}^2 , on the two-dimensional flat torus \mathbb{T}^2 , on \mathbb{R}^2 and on \mathbb{R}^3 , with consequences on Hardy inequalities on the Euclidean spaces \mathbb{R}^2 and \mathbb{R}^3 . It is crucial to consider precise geometric settings as we are interested in optimal inequalities, which rely on non-trivial symmetry results. A typical nonlinear interpolation inequality is

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathcal{X})}^2 + \lambda \|u\|_{L^2(\mathcal{X})}^2 \geq \mu_{\mathbf{A}}(\lambda) \|u\|_{L^p(\mathcal{X})}^2 \quad (1.3)$$

for any function u in the appropriate $H_{\mathbf{A}}^1$ space, for any $\lambda > 0$, and for any $p > 2$, say on the compact manifold \mathcal{X} in order to fix ideas. Assuming that $\text{vol}(\mathcal{X}) = 1$, the issue of *the optimal inequality* is to determine the largest value of $\lambda > 0$ for which we have $\mu_{\mathbf{A}}(\lambda) = \lambda + C$ for some constant C which is computed in terms of $|\mathbf{A}|$ and depends on \mathcal{X} . Equality is then realized by the constants. It is usually not difficult to prove that the equality is achieved in the inequality if $\mu_{\mathbf{A}}(\lambda)$ denotes the optimal constant, for any $\lambda > 0$. If we consider the Euler-Lagrange equation, this can be reformulated as the slightly more general *rigidity* question. For which values of λ do we know that any solution is actually a constant? To obtain rigidity, it is essential to establish symmetry properties, which is usually the most difficult step of the proof. In the non-compact case, optimal functions are not constants, which is an additional difficulty, but the problem can also be reduced to a symmetry issue.

There is another interpretation of the rigidity issue in terms of a *phase transition*. In the compact manifold case, the optimal function for (1.3) is always a constant if $\lambda > 0$ is taken small enough. For large values of λ , a simple perturbation argument shows that no minimizer can be a constant, which results in a *symmetry breaking* phenomenon, and one can prove in many cases that there is a bifurcation from a symmetric phase (solutions are constant) to a non-symmetric phase for a threshold value of λ corresponding to the optimal inequality.

Keller-Lieb-Thirring (KLT) inequalities are estimates of the ground state energy $\lambda_1[\mathbf{A}, V]$ for the magnetic Schrödinger operator $-\Delta_{\mathbf{A}} - V$ in terms of $\|V\|_{L^q(\mathcal{X})}$ and are obtained by duality from (1.3) with $q = p/(p-2)$. KLT inequalities are in fact

completely equivalent to (1.3), including for optimality issues and related rigidity questions, and essential for proving various *magnetic Hardy inequalities*, which are one of the highlights and the main motivations of this paper. However, we emphasize the fact that the accurate spectral information is carried by the KLT inequalities.

We shall actually consider not only the *superquadratic* case $p > 2$, but also the *subquadratic* case $p < 2$ in which the role of the L^2 and L^p norms are exchanged. The corresponding nonlinear interpolation inequality is

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathcal{X})}^2 + \mu \|u\|_{L^p(\mathcal{X})}^2 \geq \lambda_{\mathbf{A}}(\mu) \|u\|_{L^2(\mathcal{X})}^2 \quad (1.4)$$

for any $p \in [1, 2)$ and we can also draw a whole series of consequences (rigidity, KLT, Hardy) as in the superquadratic case. In particular, we are able to prove several new and interesting results on optimality and rigidity.

Our paper collects many results on functional inequalities with magnetic fields in different geometric settings. Therefore it is difficult to pick particularly significant results. Moreover we believe that the interest of the paper lies as much in the various methods as in the results because very little is known on optimal inequalities in presence of Aharonov-Bohm magnetic fields and on the symmetry properties of the corresponding optimal functions. The most visible outcome of our work is on Hardy inequalities, which are important tools in functional analysis. The presence of a magnetic field is a key feature, for instance in dimension $d = 2$. Let us anyway draw the attention of the reader to some results that are prominent in this paper:

- Theorem 3.1 deals with nonlinear magnetic interpolation inequalities, optimal constants and rigidity results on \mathbb{S}^1 in the subquadratic case. This is a new set of inequalities which complements the theory on *magnetic rings* in the superquadratic case studied in [14]. It was natural to study it in view of the *carré du champ* technique by Bakry and Emery in [3], but as far as we know, it is an entirely new result in presence of a magnetic potential when $p < 2$.
- Theorem 3.2 is the counterpart of Theorem 3.1 in the case of the torus $\mathbb{T}^2 \approx \mathbb{S}^1 \times \mathbb{S}^1$. It is remarkable that we achieve an optimality result here as symmetry results on products of manifolds are known to be difficult.
- Using subquadratic interpolation inequalities for proving KLT and then Hardy type inequalities is, as far as we know, an entirely new strategy: see Theorem 3.3.
- It is natural to consider also what happens on \mathbb{S}^2 : see Proposition 2.2 and Corollary 2.2 for results in the superquadratic case.
- Results on magnetic Hardy inequalities of Theorem 4.2 are a new and striking application of the nonlinear Hardy-Sobolev interpolation inequalities of [7] on the Euclidean space \mathbb{R}^2 .
- Theorems 5.1 and 5.2 are two examples of application of the nonlinear magnetic interpolation inequalities to magnetic Hardy inequalities on \mathbb{R}^3 , which significantly improve upon the results in [19, 22].

Let us conclude this introduction by some mathematical observations and some additional references. The overall question is to determine the functional spaces which are adapted to magnetic Schrödinger operators in the spirit of [13]. Magnetic interpolation inequalities (without optimal constants) are usually not an issue as they can be deduced from the non-magnetic interpolation inequalities by the *diamagnetic inequality*: see for instance [27]. However we are interested in retaining information about the magnetic field and characterizing optimality cases, which is by far a more difficult target. As a convention, we shall speak of *Hardy-Sobolev* inequalities when a term $\int |x|^{-2} |u|^2 dx$ is subtracted from the kinetic energy and of *Caffarelli-Kohn-Nirenberg* inequalities when various pure power weights are taken into account. On \mathbb{R}^d , $\Delta_{\mathbf{A}}$ has the same scaling properties as the non-magnetic Laplacian if \mathbf{A} is a Aharonov-Bohm magnetic potential and its spectrum is explicit. KLT inequalities are spectral estimates but it has to be emphasized that they differ from semi-classical estimates as they inherit nonlinear properties of the interpolation inequalities. Finally, it is a remarkable fact that, in presence of a magnetic potential, a Hardy inequality can be established in the two-dimensional case (see [14,22,24,26] for related papers).

This paper is organized as follows. Section 2 is devoted to some preliminary results on the circle \mathbb{S}^1 and on the two-dimensional sphere \mathbb{S}^2 . New interpolation inequalities are established on the sphere \mathbb{S}^2 , with an optimality result. Section 3 is devoted to the study of a class of subquadratic magnetic interpolation inequalities on \mathbb{S}^1 and on the flat torus \mathbb{T}^2 . We are able to identify a sharp condition of optimality and deduce several Hardy inequalities in dimensions $d = 2$ and $d = 3$. In Section 4 we start by recalling earlier, non-optimal but numerically almost sharp, results on interpolation inequalities on \mathbb{R}^2 in the presence of an Aharonov-Bohm magnetic field in order to establish some Hardy inequalities on \mathbb{R}^2 . Section 5 is devoted to Hardy inequalities on \mathbb{R}^3 with singularities which are either spherically or cylindrically symmetric.

2. General set-up and preliminary results

This section is devoted to various results on the sphere \mathbb{S}^d without magnetic field (Section 2.1), in the superquadratic case when $d = 1$ (Section 2.2) and when $d = 2$ (Section 2.3). Sections 2.1 and 2.2 are devoted to a survey of previous results and are given not only for completeness but also as introductory material for Section 2.3 and Section 3.

2.1. Non-magnetic interpolation inequalities on \mathbb{S}^d

On the sphere \mathbb{S}^d , we consider the uniform probability measure $d\sigma$, which is the measure induced by the Lebesgue measure in \mathbb{R}^{d+1} , duly normalized and denote by $\|\cdot\|_{L^q(\mathbb{S}^d)}$ the corresponding L^q norm. Here we state known results for later use.

2.1.1. *Interpolation inequalities without weights*

The interpolation inequalities

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 \geq \frac{d}{p-2} \left(\|u\|_{L^p(\mathbb{S}^d)}^2 - \|u\|_{L^2(\mathbb{S}^d)}^2 \right) \quad (2.1)$$

hold for any $p \in [1, 2) \cup (2, +\infty)$ if $d = 1$ and $d = 2$, and for any $p \in [1, 2) \cup (2, 2^*)$ if $d \geq 3$, where $2^* := 2d/(d-2)$ is the Sobolev critical exponent. See [5,6] for $p > 2$ and [3] if $d = 1$ or $d \geq 2$ and $p \leq (2d^2 + 1)/(d-1)^2$.

If $p > 2$, we know from [12] that there exists a concave monotone increasing function $\lambda \mapsto \mu_{0,p}(\lambda)$ on $(0, +\infty)$ such that $\mu_{0,p}(\lambda)$ is the optimal constant in the inequality

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu_{0,p}(\lambda) \|u\|_{L^p(\mathbb{S}^d)}^2 \quad \forall u \in H^1(\mathbb{S}^d) \quad (2.2)$$

and that $\mu_{0,p}(\lambda) = \lambda$ if and only if $\lambda \leq d/(p-2)$. In this range, equality is achieved if and only if u is a constant function: this is a *symmetry* range. On the opposite, if $\lambda > d/(p-2)$, the optimal function is not constant and we shall say that there is *symmetry breaking*.

The case $1 \leq p < 2$ is similar: there exists a concave monotone increasing function $\mu \mapsto \lambda_{0,p}(\mu)$ on $(0, +\infty)$ such that $\lambda_{0,p}(\mu)$ is the optimal constant in the inequality

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \mu \|u\|_{L^p(\mathbb{S}^d)}^2 \geq \lambda_{0,p}(\mu) \|u\|_{L^2(\mathbb{S}^d)}^2 \quad \forall u \in H^1(\mathbb{S}^d) \quad (2.3)$$

and that $\lambda_{0,p}(\mu) = \mu$ if and only if $\mu \leq d/(2-p)$. In this symmetry range, constants are the optimal functions, while there is symmetry breaking if $\mu > d/(2-p)$: optimal functions are non-constant.

In the symmetry range, positive constants are actually the only *positive* solutions of the Euler-Lagrange equation

$$-\varepsilon \Delta u + \lambda u = u^{p-1}$$

where $\varepsilon = \pm 1$ is the sign of $(p-2)$, while there are multiple solutions in the symmetry breaking range. The limit case $p = 2$ can be obtained by taking the limit as $p \rightarrow 2$ and the corresponding inequality is the logarithmic Sobolev inequality. Much more is known and we refer to [12] for further details.

 2.1.2. *A weighted Poincaré inequality for the ultra-spherical operator*

Using cylindrical coordinates $(z, \omega) \in [-1, 1] \times \mathbb{S}^{d-1}$, we can rewrite the Laplace-Beltrami operator on \mathbb{S}^d as

$$\Delta = \mathcal{L}_d + \frac{1}{1-z^2} \Delta_\omega \quad \text{with} \quad \mathcal{L}_d u := (1-z^2) u'' - dz u'$$

where Δ_ω denotes the Laplace-Beltrami operator on \mathbb{S}^{d-1} and \mathcal{L}_d is the *ultra-spherical operator*. In other words, \mathcal{L}_d is the Laplace-Beltrami operator on \mathbb{S}^d restricted to functions which depend only on z . The operator \mathcal{L}_d has a basis of

eigenfunctions $G_{\ell,d}$, the Gegenbauer polynomials, associated with the eigenvalues $\ell(\ell + d - 1)$ for any $\ell \in \mathbb{N}$ (see [28]). Here d is not necessarily an integer.

Let us consider the eigenvalue problem

$$-\mathcal{L}_2 f + \frac{4\mathbf{a}^2}{1-z^2} f = \lambda f. \quad (2.4)$$

By changing the unknown function according to $f(z) = (1-z^2)^{\mathbf{a}} g(z)$, we obtain that g solves

$$-\mathcal{L}_{2(2\mathbf{a}+1)} g + 2\mathbf{a}(1+2\mathbf{a})g = \lambda g$$

which determines the eigenvalues $\lambda = \lambda_{\ell,\mathbf{a}}$ given by

$$\lambda_{\ell,\mathbf{a}} = \ell(\ell + 2(2\mathbf{a} + 1) - 1) + 2\mathbf{a}(1 + 2\mathbf{a}) = (\ell + 2\mathbf{a})(\ell + 2\mathbf{a} + 1), \quad \ell \in \mathbb{N}. \quad (2.5)$$

We shall denote by $g_{\ell,\mathbf{a}}(z) = G_{\ell,2(2\mathbf{a}+1)}(z)$ the associated eigenfunctions and define $f_{\ell,\mathbf{a}}(z) := (1-z^2)^{\mathbf{a}} g_{\ell,\mathbf{a}}(z)$. By considering the lowest positive eigenvalue, we obtain a *weighted Poincaré inequality*.

Lemma 2.1. *For any $\mathbf{a} \in \mathbb{R}$ and any function $f \in \mathbf{H}_0^1[-1, 1]$, we have*

$$\int_{-1}^1 \left((1-z^2) |f'(z)|^2 + \frac{4\mathbf{a}^2}{1-z^2} |f(z)|^2 \right) dz \geq \lambda_{1,\mathbf{a}} \int_{-1}^1 |f(z) - \bar{f}(z)|^2 dz,$$

where

$$\bar{f}(z) = (1-z^2)^{\mathbf{a}} \frac{\int_{-1}^1 f(z) (1-z^2)^{\mathbf{a}} dz}{\int_{-1}^1 (1-z^2)^{2\mathbf{a}} dz}.$$

Equality is achieved by a function f if and only if f is proportional to $f_{1,\mathbf{a}}(z) = z(1-z^2)^{\mathbf{a}}$.

Notice for consistency that, if $f(z) = (1-z^2)^{\mathbf{a}} g(z)$, then

$$\begin{aligned} & \int_{-1}^1 \left((1-z^2) |f'(z)|^2 + \frac{4\mathbf{a}^2}{1-z^2} |f(z)|^2 \right) dz \\ &= \int_{-1}^1 \left((1-z^2) |g'(z)|^2 + 2\mathbf{a}(1+2\mathbf{a}) |g(z)|^2 \right) (1-z^2)^{2\mathbf{a}} dz, \end{aligned}$$

where the right-hand side is the Dirichlet form associated with the operator

$$-\mathcal{L}_{2(2\mathbf{a}+1)} + 2\mathbf{a}(1+2\mathbf{a}).$$

2.2. Magnetic rings: superquadratic inequalities on \mathbb{S}^1

In this section, we review a series of results which have been obtained in [14] in the superquadratic case $p > 2$, in preparation for an extension to the subquadratic case $p \in [1, 2)$ that will be studied in Section 3.

2.2.1. *Magnetic interpolation inequalities and consequences*

Let us consider the superquadratic case $p > 2$ in dimension $d = 1$. We recall that $d\sigma = (2\pi)^{-1} d\theta$ where $\theta \in [0, 2\pi) \approx \mathbb{S}^1$. As in [14] we consider the space $H^1(\mathbb{S}^1)$ of the 2π -periodic functions $u \in C^{0,1/2}(\mathbb{S}^1)$, such that $u' \in L^2(\mathbb{S}^1)$. Inequality (2.2) can be rewritten as

$$\|u'\|_{L^2(\mathbb{S}^1)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^1)}^2 \geq \lambda \|u\|_{L^p(\mathbb{S}^1)}^2 \quad \forall u \in H^1(\mathbb{S}^1) \quad (2.6)$$

for any $\lambda \in (0, 1/(p-2)]$. We also have the inequality

$$\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{1}{4} \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} \geq \frac{1}{4} \|u\|_{L^2(\mathbb{S}^1)}^2 \quad \forall u \in H^1(\mathbb{S}^1), \quad (2.7)$$

according to [21], with the convention that $\|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} = 0$ if u^{-2} is not integrable and, as a special case, if u changes sign. Notice that inequality (2.7) is formally the case $p = 2d/(d-2)$ and $\lambda = d/(p-2)$ of (2.2) when $d = 1$ (see [14, Appendix A]).

In [14], it was shown that the inequality (for complex valued functions)

$$\|\psi' - ia\psi\|_{L^2(\mathbb{S}^1)}^2 + \lambda \|\psi\|_{L^2(\mathbb{S}^1)}^2 \geq \mu_{a,p}(\lambda) \|\psi\|_{L^p(\mathbb{S}^1)}^2 \quad \forall \psi \in H^1(\mathbb{S}^1, \mathbb{C}) \quad (2.8)$$

is equivalent, after eliminating the phase, to the inequality

$$\|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \lambda \|u\|_{L^2(\mathbb{S}^1)}^2 \geq \mu_{a,p}(\lambda) \|u\|_{L^p(\mathbb{S}^1)}^2 \quad \forall u \in H^1(\mathbb{S}^1).$$

The equivalence is relatively easy to prove if ψ does not vanish, but some care is required otherwise: see [14] for details. Here we denote by $\mu_{a,p}(\lambda)$ the optimal constant in (2.8). Using (2.8) and then (2.6), we obtain that

$$\begin{aligned} & \|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \lambda \|u\|_{L^2(\mathbb{S}^1)}^2 \\ &= (1 - 4a^2) \|u'\|_{L^2(\mathbb{S}^1)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^1)}^2 + 4a^2 (\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{1}{4} \|u^{-1}\|_{L^2(\mathbb{S}^1)}^2) \\ &\geq (1 - 4a^2) (\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{a^2 + \lambda}{1 - 4a^2} \|u\|_{L^2(\mathbb{S}^1)}^2) \geq (a^2 + \lambda) \|u\|_{L^p(\mathbb{S}^1)}^2 \end{aligned}$$

under the condition $(a^2 + \lambda)/(1 - 4a^2) \leq 1/(p-2)$, which provides an estimate of $\mu_{a,p}(\lambda)$. This estimate turns out to be optimal.

Proposition 2.1 ([14]). *Let $p > 2$, $a \in [0, 1/2]$, and $\lambda > -a^2$.*

- (i) *If $a^2(p+2) + \lambda(p-2) \leq 1$, then $\mu_{a,p}(\lambda) = a^2 + \lambda$ and equality in (2.8) is achieved only by the constants.*
- (ii) *If $a^2(p+2) + \lambda(p-2) > 1$, then $\mu_{a,p}(\lambda) < a^2 + \lambda$ and equality in (2.8) is not achieved by the constants.*

The condition $a \in [0, 1/2]$ is not a restriction. First, replacing ψ by $e^{iks} \psi(s)$ for any $k \in \mathbb{Z}$ shows that $\mu_{a+k,p}(\mu) = \mu_{a,p}(\mu)$ so that we can assume that $a \in [0, 1]$. Then by considering $\chi(s) = e^{-is} \bar{\psi}(s)$, we find that

$$|\psi' + ia\psi|^2 = |\chi' + i(1-a)\chi|^2,$$

hence $\mu_{a,p}(\mu) = \mu_{1-a,p}(\mu)$.

2.2.2. Magnetic Hardy inequalities on \mathbb{S}^1 and \mathbb{R}^2

As in [14], we can draw an easy consequence of Proposition 2.1 on a Keller-Lieb-Thirring type inequality. By Hölder's inequality applied with $q = p/(p-2)$, we have

$$\|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 - \mu^{-1} \int_{\mathbb{S}^1} \phi |\psi|^2 d\theta \geq \|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 - \mu^{-1} \|\phi\|_{L^q(\mathbb{S}^1)} \|\psi\|_{L^p(\mathbb{S}^1)}^2.$$

Using (2.8) with $\lambda = 0$ and μ such that $\mu^{-1} \|\phi\|_{L^q(\mathbb{S}^1)} = \mu_{a,p}(0)$, we know that the right-hand side is nonnegative. See [14] for more details. Altogether we obtain the following *magnetic Hardy inequality on \mathbb{S}^1* : for any $a \in \mathbb{R}$, any $p > 2$ and $q = p/(p-2)$, if ϕ is a non-trivial potential in $L^q(\mathbb{S}^1)$, then

$$\|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 \geq \frac{\mu_{a,p}(0)}{\|\phi\|_{L^q(\mathbb{S}^1)}} \int_{\mathbb{S}^1} \phi |\psi|^2 d\sigma \quad \forall \psi \in H_{\mathbf{A}}^1(\mathbb{S}^1). \quad (2.9)$$

This is a special case of the more general interpolation inequality

$$\begin{aligned} \|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 - \int_{\mathbb{S}^1} \phi |\psi|^2 d\theta &\geq \|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 - \mu \|\psi\|_{L^p(\mathbb{S}^1)}^2 \\ &\geq -\lambda_{a,p}(\mu) \|\psi\|_{L^2(\mathbb{S}^1)}^2 \end{aligned} \quad (2.10)$$

with $\mu = \|\phi\|_{L^q(\mathbb{S}^1)}$, where we denote by $\lambda_{a,p}(\mu)$ the inverse function of $\lambda \mapsto \mu_{a,p}(\lambda)$, as defined in Proposition 2.1. See [13] for details.

The standard non-magnetic Hardy inequality on \mathbb{R}^d , *i.e.*,

$$\int_{\mathbb{R}^d} |\nabla \psi|^2 dx \geq \frac{1}{4} (d-2)^2 \int_{\mathbb{R}^d} \frac{|\psi|^2}{|x|^2} dx \quad \forall \psi \in H^1(\mathbb{R}^d),$$

degenerates if $d = 2$, but this degeneracy is lifted in the presence of a Aharonov-Bohm magnetic field. According to [26], we have

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \min_{k \in \mathbb{Z}} (a-k)^2 \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx \quad \forall \psi \in H^1(\mathbb{R}^d).$$

It is natural to ask whether an improvement can be obtained if the singularity $|x|^{-2}$ is replaced by a weight which has an angular dependence. Using polar coordinates $x \approx (r, \theta)$ and interpolation inequalities of [12], the inequality

$$\int_{\mathbb{R}^d} |\nabla \psi|^2 dx \geq \frac{(d-2)^2}{4 \|\varphi\|_{L^q(\mathbb{S}^{d-1})}} \int_{\mathbb{R}^d} \frac{\varphi(\theta)}{|x|^2} |\psi|^2 dx \quad \forall \psi \in H^1(\mathbb{R}^d)$$

was proved in [24], under the condition that $q \geq 1 + \frac{1}{2} (d-2)^2/(d-1)$, again with normalized measure on \mathbb{S}^{d-1} . Magnetic and non-radial improvements have been combined in [14]. Let us give a statement in preparation for similar extensions to the case of dimension $d = 3$.

Corollary 2.1 ([14]). *Let \mathbf{A} as in (1.1), $a \in [0, 1/2]$, $p > 2$, $q = p/(p-2)$ and assume that φ is a non-negative function in $L^q(\mathbb{S}^1)$. With the above notations, the*

inequality

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \tau \int_{\mathbb{R}^2} \frac{\varphi(\theta)}{|x|^2} |\psi|^2 dx \quad \forall \psi \in H^1(\mathbb{R}^d)$$

holds with a constant $\tau > 0$ which is the unique solution of the equation

$$\lambda_{a,p}(\tau \|\varphi\|_{L^q(\mathbb{S}^1)}) = 0.$$

Moreover, $\tau = a^2 / \|\varphi\|_{L^q(\mathbb{S}^1)}$ if $a^2 \leq 1/(p+2)$.

2.3. Magnetic interpolation inequalities on \mathbb{S}^2

In the spirit of Sections 2.1 and 2.2, we state some new results concerning the two-dimensional sphere with main results in Proposition 2.2 and Corollary 2.2.

2.3.1. A magnetic ground state estimate

Let us consider the magnetic Laplacian on \mathbb{S}^2 and the associated Dirichlet form $\int_{\mathbb{S}^2} |\nabla_{\mathbf{A}} u|^2 d\sigma$ where $d\sigma$ is the uniform probability measure on \mathbb{S}^2 . Using cylindrical coordinates $(\theta, z) \in [0, 2\pi) \times [-1, 1]$, we can write that $d\sigma = \frac{1}{4\pi} dz d\theta$ and assume that the magnetic gradient takes the form

$$\nabla_{\mathbf{A}} u = \left(\begin{array}{c} \sqrt{1-z^2} \frac{\partial u}{\partial z} \\ \frac{1}{\sqrt{1-z^2}} \left(\frac{\partial u}{\partial \theta} - i a u \right) \end{array} \right) \quad (2.11)$$

where $a > 0$ is a magnetic flux, so that

$$|\nabla_{\mathbf{A}} u|^2 = (1-z^2) \left| \frac{\partial u}{\partial z} \right|^2 + \frac{1}{1-z^2} \left| \frac{\partial u}{\partial \theta} - i a u \right|^2.$$

Lemma 2.2. Assume that $a \in \mathbb{R}$. With the notation (2.11), we have

$$\int_{\mathbb{S}^2} |\nabla_{\mathbf{A}} u|^2 d\sigma \geq \Lambda_a \int_{\mathbb{S}^2} |u|^2 d\sigma \quad \forall u \in H_{\mathbf{A}}^1(\mathbb{S}^2)$$

with optimal constant

$$\Lambda_a = \min_{k \in \mathbb{Z}} |k - a| (|k - a| + 1). \quad (2.12)$$

Notice that $\Lambda_a \leq \Lambda_{1/2} = 3/4$.

Proof. We can write u using a Fourier decomposition

$$u(z, \theta) = \sum_{\ell \in \mathbb{N}} \sum_{k \in \mathbb{Z}} u_{k,\ell}(z) e^{i k \theta}$$

and observe that

$$|\nabla_{\mathbf{A}} u|^2 = \sum_{\ell \in \mathbb{N}} \sum_{k \in \mathbb{Z}} \left((1-z^2) |u'_{k,\ell}(z)|^2 + \frac{(k-a)^2}{1-z^2} |u_{k,\ell}(z)|^2 \right)$$

where

$$u_{k,\ell}(z) = f_{\ell,|k-a|/2}(z) \int_{\mathbb{S}^2} u(z,\theta) \frac{2 f_{\ell,|k-a|/2}(z) e^{-ik\theta}}{\int_{-1}^1 (1-z^2)^{|k-a|} dz} d\sigma$$

and $f_{\ell,|k-a|/2}$ is an eigenfunction of (2.4) with eigenvalue $\lambda = \lambda_{\ell,\mathbf{a}}$ such that $2\mathbf{a} = |k-a|$. Using (2.5), we conclude that the spectrum of $-\Delta_{\mathbf{A}}$ is given by

$$(\ell + |k-a|)(\ell + |k-a| + 1), \quad k \in \mathbb{Z}, \ell \in \mathbb{N}. \quad \square$$

2.3.2. Superquadratic interpolation inequalities and consequences

Proposition 2.2. *Let $a \in \mathbb{R}$ and $p > 2$. With the notation (2.11), there exists a concave monotone increasing function $\lambda \mapsto \mu_{a,p}(\lambda)$ on $(-\Lambda_a, +\infty)$ such that $\mu_{a,p}(\lambda)$ is the optimal constant in the inequality*

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{S}^2)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^2)}^2 \geq \mu_{a,p}(\lambda) \|u\|_{L^p(\mathbb{S}^2)}^2 \quad \forall u \in H_{\mathbf{A}}^1(\mathbb{S}^2).$$

Furthermore, $\mu_{a,p}(\lambda) \geq 2(\lambda + \Lambda_a)/(2 + (p-2)\Lambda_a)$ and $\lim_{\lambda \rightarrow -\Lambda_a} \mu_{a,p}(\lambda) = 0$, with Λ_a given by (2.12).

Proof. The proof is adapted from [13, Proposition 3.1]. For an arbitrary $t \in (0, 1)$, we can write that

$$\begin{aligned} \|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{S}^2)}^2 + \lambda \|u\|_{L^2(\mathbb{S}^2)}^2 &\geq t \left(\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{S}^2)}^2 - \Lambda_a \|u\|_{L^2(\mathbb{S}^2)}^2 \right) \\ &\quad + (1-t) \left(\|\nabla |u|\|_{L^2(\mathbb{S}^2)}^2 + \frac{\lambda + t\Lambda_a}{1-t} \|u\|_{L^2(\mathbb{S}^2)}^2 \right) \\ &\geq (1-t) \mu_{0,p} \left(\frac{\lambda + t\Lambda_a}{1-t} \right) \|u\|_{L^p(\mathbb{S}^2)}^2, \end{aligned}$$

as a consequence of Lemma 2.2 and of the *diamagnetic inequality* (see e.g. [27, Theorem 7.21])

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{S}^2)}^2 \geq \|\nabla |u|\|_{L^2(\mathbb{S}^2)}^2.$$

If $\lambda < 2/(p-2)$, the estimate is obtained by choosing t such that

$$\frac{\lambda + t\Lambda_a}{1-t} = \frac{2}{p-2}$$

and recalling that $\mu_{0,p}(2/(p-2)) = 2/(p-2)$. The limit as $\lambda \rightarrow -\Lambda_a$ is obtained by taking the ground state of $-\Delta_{\mathbf{A}}$ on $H^1(\mathbb{S}^2)$ as test function. \square

With the same method as for the proof of (2.9), we can deduce a Hardy-type inequality.

Corollary 2.2. *Let $a \in \mathbb{R}$, $p > 2$ and $q = p/(p-2)$. With the notation (2.11), if ϕ is a non-trivial potential in $L^q(\mathbb{S}^2)$, then*

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{S}^2)}^2 \geq \frac{\mu_{a,p}(0)}{\|\phi\|_{L^q(\mathbb{S}^2)}} \int_{\mathbb{S}^2} \phi |u|^2 d\sigma \quad \forall u \in H_{\mathbf{A}}^1(\mathbb{S}^2).$$

3. Subquadratic magnetic interpolation inequalities

This section is devoted to new and optimal interpolation inequalities involving L^p norms in the *subquadratic* range $p \in (1, 2)$ on the circle and on the torus (Sections 3.1 and 3.2). Dual Keller-Lieb-Thirring inequalities yield new magnetic Hardy inequalities on \mathbb{R}^2 and \mathbb{R}^3 (Section 3.3).

3.1. Magnetic rings: subquadratic interpolation inequalities on \mathbb{S}^1

We extend the results of Section 2.2.1 to the subquadratic range $1 < p < 2$ using the strategy of [14].

3.1.1. Statement of the inequality

As a special case of (2.1) corresponding to $d = 1$, we have the non-magnetic interpolation inequality

$$(2 - p) \|u'\|_{L^2(\mathbb{S}^1)}^2 + \|u\|_{L^p(\mathbb{S}^1)}^2 \geq \|u\|_{L^2(\mathbb{S}^1)}^2 \quad \forall u \in H^1(\mathbb{S}^1) \quad (3.1)$$

for any $p \in [1, 2)$. Our first result is the magnetic counterpart of this inequality.

Lemma 3.1. *Let $a \in \mathbb{R}$ and $p \in [1, 2)$. Then there exists a concave monotone increasing function $\mu \mapsto \lambda_{a,p}(\mu)$ on \mathbb{R}^+ such that*

$$\|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 + \mu \|\psi\|_{L^p(\mathbb{S}^1)}^2 \geq \lambda_{a,p}(\mu) \|\psi\|_{L^2(\mathbb{S}^1)}^2 \quad \forall \psi \in H^1(\mathbb{S}^1, \mathbb{C}). \quad (3.2)$$

Here we denote by $\lambda_{a,p}(\mu)$ the optimal constant in (3.1).

Proof. The existence of $\lambda_{a,p}(\mu)$ is a consequence of (3.1) and of the *diamagnetic inequality*: let $\rho = |\psi|$ and ϕ be such that $\psi = \rho(\theta) \exp(i\phi(\theta))$. Since

$$|\psi' - i a \psi|^2 = |\rho'|^2 + |\phi' - a|^2 \rho^2 \geq |\rho'|^2,$$

we have that $\|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 \geq \|\rho'\|_{L^2(\mathbb{S}^1)}^2$. The concavity of $\mu \mapsto \lambda_{a,p}(\mu)$ is a consequence of the definition of $\lambda_{a,p}(\mu)$ as the optimal constant, *i.e.*, the infimum on $H^1(\mathbb{S}^1) \ni \psi$ of an affine function of μ . \square

3.1.2. Existence of an optimal function

Lemma 3.2. *For all $a \in [0, 1/2]$, $p \in [1, 2)$ and $\mu \geq -a^2$, equality in (3.2) is achieved by at least one function in $H^1(\mathbb{S}^1)$.*

Proof. We consider a minimizing sequence $\{\psi_n\}$ for

$$\lambda_{a,p}(\mu) = \inf \left\{ \|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2 + \mu \|\psi\|_{L^p(\mathbb{S}^1)}^2 : \psi \in H^1(\mathbb{S}^1), \|\psi\|_{L^2(\mathbb{S}^1)} = 1 \right\}.$$

By the diamagnetic inequality we know that the sequence $(\psi_n)_{n \in \mathbb{N}}$ is bounded in $H^1(\mathbb{S}^1)$. By the compact Sobolev embeddings, this sequence is relatively compact in $L^p(\mathbb{S}^1)$ and in $L^2(\mathbb{S}^1)$. The map $\psi \mapsto \|\psi' - i a \psi\|_{L^2(\mathbb{S}^1)}^2$ is lower semicontinuous by Fatou's lemma, which proves the claim. \square

3.1.3. *A non-vanishing property*

Lemma 3.3. *Assume that $a \in (0, 1/2)$, $p \in [1, 2)$ and $\mu \geq -a^2$. If $\psi \in H^1(\mathbb{S}^1)$ is an optimal function for (3.2) with $\|\psi\|_{L^p(\mathbb{S}^1)} = 1$, then $\psi(s) \neq 0$ for any $s \in \mathbb{S}^1$.*

Proof. The proof goes as in [14]. Let us decompose $v(s) = \psi(s) e^{ias}$ as a real and an imaginary part, respectively v_1 and v_2 , which both solve the same Euler-Lagrange equation

$$-v_j'' - \mu (v_1^2 + v_2^2)^{\frac{p}{2}-1} v_j = \lambda_{a,p}(\mu) v_j, \quad j = 1, 2.$$

Notice that $v \in C^{0,1/2}(\mathbb{S}^1)$ and the nonlinear term is continuous, hence v is smooth. The Wronskian $w = (v_1 v_2' - v_1' v_2)$ is constant. If both v_1 and v_2 vanish at the same point, then w vanishes identically, which means that v_1 and v_2 are proportional. With $a \in (0, 1/2)$, ψ is not 2π -periodic, a contradiction. \square

3.1.4. *A reduction to a scalar minimization problem*

We refer to Section 2.1.1 if $a = 0$ and assume in the proofs that $a > 0$. The main steps of the reduction are similar to the case $p > 2$ of [14]. We repeat the key points for completeness. Let us define

$$\mathcal{Q}_{a,p,\mu}[u] := \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \mu \|u\|_{L^p(\mathbb{S}^1)}^2}{\|u\|_{L^2(\mathbb{S}^1)}^2}.$$

Notice that if $u \in H^1(\mathbb{S}^1)$ is such that $u(s_0) = 0$ for some $s_0 \in (-\pi, \pi]$, then

$$|u(s)|^2 = \left(\int_{s_0}^s u' ds \right)^2 \leq \sqrt{2\pi} \|u'\|_{L^2(\mathbb{S}^1)} \sqrt{|s - s_0|}$$

and u^{-2} is not integrable. In this case, as mentioned earlier, we adopt the convention that

$$\mathcal{Q}_{a,p,\mu}[u] := \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + \mu \|u\|_{L^p(\mathbb{S}^1)}^2}{\|u\|_{L^2(\mathbb{S}^1)}^2}. \quad (3.3)$$

Lemma 3.4. *For any $a \in [0, 1/2]$, $p \in [1, 2)$, $\mu > -a^2$,*

$$\lambda_{a,p}(\mu) = \min_{u \in H^1(\mathbb{S}^1) \setminus \{0\}} \mathcal{Q}_{a,p,\mu}[u].$$

Proof. We consider functions on \mathbb{S}^1 as 2π -periodic functions on \mathbb{R} . If $\psi \in H^1(\mathbb{S}^1)$, then $v(s) = \psi(s) e^{ias}$ satisfies the condition

$$v(s + 2\pi) = e^{2i\pi a} v(s) \quad \forall s \in \mathbb{R} \quad (3.4)$$

and

$$\lambda_{a,p}(\mu) = \min \frac{\|v'\|_{L^2(\mathbb{S}^1)}^2 + \mu \|v\|_{L^p(\mathbb{S}^1)}^2}{\|v\|_{L^2(\mathbb{S}^1)}^2}$$

where the minimization is taken on the set of the functions $v \in C^{0,1/2}(\mathbb{R})$ such that $v' \in L^2(-\pi, \pi)$ and (3.4) holds.

With $v = u e^{i\phi}$ written in polar form, the boundary condition becomes

$$u(\pi) = u(-\pi), \quad \phi(\pi) = 2\pi(a + k) + \phi(-\pi) \quad (3.5)$$

for some $k \in \mathbb{Z}$, and $\|v'\|_{L^2(\mathbb{S}^1)}^2 = \|u'\|_{L^2(\mathbb{S}^1)}^2 + \|u\phi'\|_{L^2(\mathbb{S}^1)}^2$ so that

$$\lambda_{a,p}(\mu) = \min \frac{\|u'\|_{L^2(\mathbb{S}^1)}^2 + \|u\phi'\|_{L^2(\mathbb{S}^1)}^2 + \mu \|u\|_{L^p(\mathbb{S}^1)}^2}{\|u\|_{L^2(\mathbb{S}^1)}^2}$$

where the minimization is taken on the set of the functions $(u, \phi) \in C(\mathbb{R})^2$ such that $u', u\phi' \in L^2(\mathbb{S}^1)$ and (3.5) holds.

Up to a multiplication of u by a constant so that $\|u\|_{L^p(\mathbb{S}^1)} = 1$, the Euler-Lagrange equations are

$$-u'' + |\phi'|^2 u + \mu |u|^{p-2} u = \lambda_{a,p}(\mu) u \quad \text{and} \quad (\phi' u^2)' = 0.$$

If $a \in (0, 1/2)$, by integrating the second equation and using Lemma 3.3, we find a constant L such that $\phi' = L/u^2$. Taking (3.5) into account, we deduce from

$$L \int_{-\pi}^{\pi} \frac{ds}{u^2} = \int_{-\pi}^{\pi} \phi' ds = 2\pi(a + k)$$

that

$$\|u\phi'\|_{L^2(\mathbb{S}^1)}^2 = L^2 \int_{-\pi}^{\pi} \frac{ds}{u^2} = \frac{(a + k)^2}{\|u^{-1}\|_{L^2(\mathbb{S}^1)}^2}.$$

This establishes that

$$\lambda_{a,p}(\mu) = \min_{u, k} \mathcal{Q}_{a+k,p,\mu}[u]$$

where the minimization is taken on all $k \in \mathbb{Z}$ and on all functions $u \in H^1(\mathbb{S}^1)$. Because of the restriction $a \in (0, 1/2)$, the minimum is achieved by $k = 0$.

The case $a = 1/2$ is a limit case that can be handled as in [14, Theorem III.7]. In this case the result holds also true, with the minimizer being in $H_0^1(\mathbb{S}^1) \setminus \{0\}$, and with the convention defined in (3.3) for the expression of $\mathcal{Q}_{a,p,\mu}[u]$ when u vanishes in \mathbb{S}^1 . \square

3.1.5. A rigidity result

If $a \in (0, 1/2)$, as in [14], the study of (3.2) is reduced to the study of the inequality

$$\|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \mu \|u\|_{L^p(\mathbb{S}^1)}^2 \geq \lambda_{a,p}(\mu) \|u\|_{L^2(\mathbb{S}^1)}^2 \quad \forall u \in H^1(\mathbb{S}^1) \quad (3.6)$$

where u is now a real valued function. Necessary adaptations to the trivial case $a = 0$ and to the limit case $a = 1/2$ are straightforward and left to the reader. The result below is the analogue of Proposition 2.1 in the subcritical range.

Theorem 3.1. *Let $p \in (1, 2)$, $a \in (0, 1/2)$, and $\mu > 0$.*

- (i) If $\mu(2-p) + 4a^2 \leq 1$, then $\lambda_{a,p}(\mu) = a^2 + \mu$ and equality in (3.6) is achieved only by the constants.
- (ii) If $\mu(2-p) + 4a^2 > 1$, then $\lambda_{a,p}(\mu) < a^2 + \mu$ and equality in (3.6) is not achieved by the constants.

Proof. In case (i) we can write

$$\begin{aligned} & \|u'\|_{L^2(\mathbb{S}^1)}^2 + a^2 \|u^{-1}\|_{L^2(\mathbb{S}^1)}^{-2} + \mu \|u\|_{L^p(\mathbb{S}^1)}^2 \\ &= (1 - 4a^2) \left(\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{\mu}{1 - 4a^2} \|u\|_{L^p(\mathbb{S}^1)}^2 \right) \\ & \quad + 4a^2 \left(\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{1}{4} \|u^{-1}\|_{L^2(\mathbb{S}^1)}^2 \right), \end{aligned}$$

deduce from (3.1) that

$$\|u'\|_{L^2(\mathbb{S}^1)}^2 + \frac{\mu}{1 - 4a^2} \|u\|_{L^p(\mathbb{S}^1)}^2 \geq \frac{\mu}{1 - 4a^2} \|u\|_{L^2(\mathbb{S}^1)}^2$$

if $\mu/(1 - 4a^2) \leq 1/(2-p)$ and conclude using (2.7).

In case (ii), let us consider the test function $u_\varepsilon := 1 + \varepsilon w_1$, where w_1 is the eigenfunction corresponding to the first non-zero eigenvalue of $-d^2/ds^2$ on $H^1(\mathbb{S}^1)$, with periodic boundary conditions, namely, $w_1(s) = \cos s$ and $\lambda_1 = 1$. A Taylor expansion shows that

$$\mathcal{Q}_{a,p,\mu}[u_\varepsilon] = (1 + a^2 - \mu(2-p)) \varepsilon^2 + o(\varepsilon^2),$$

which proves the result. Notice that the Taylor expansion is also valid if $a = 0$, so that $(p-2)$ is the optimal constant in (3.1), and also that a similar Taylor expansion holds in case of (2.7), which formally corresponds to $p = -2$. \square

3.2. Aharonov-Bohm magnetic interpolation inequalities on \mathbb{T}^2

We consider a toy model for Aharonov-Bohm magnetic fields on the flat torus which can be seen as a 2-dimensional cylinder, with periodicity along the axis. It is a 2-dimensional extension of the superquadratic magnetic ring model of Section 3.1.

Let us consider the flat torus $\mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1 \approx [-\pi, \pi) \times [-\pi, \pi) \ni (x, y)$ with periodic boundary conditions in x and y . We denote by $d\sigma$ the uniform probability measure $d\sigma = dx dy/(4\pi^2)$ and consider the magnetic gradient

$$\nabla_{\mathbf{A}} \psi := (\psi_x, \psi_y - i a \psi) \tag{3.7}$$

and the magnetic kinetic energy

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{T}^2)}^2 = \iint_{\mathbb{T}^2} |\nabla_{\mathbf{A}} \psi|^2 d\sigma = \iint_{\mathbb{T}^2} (|\psi_x|^2 + |\psi_y - i a \psi|^2) d\sigma.$$

3.2.1. A magnetic ground state estimate

Lemma 3.5. *Assume that $a \in [0, 1/2]$. With the notation (3.7), we have*

$$\iint_{\mathbb{T}^2} |\nabla_{\mathbf{A}} \psi|^2 d\sigma \geq a^2 \iint_{\mathbb{T}^2} |\psi|^2 d\sigma \quad \forall \psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{T}^2).$$

Proof. We make a Fourier decomposition on the basis $(e^{i\ell x} e^{ik y})_{k,\ell \in \mathbb{Z}}$. We find that the lowest modes are given by

$$\begin{aligned} k = 0, \ell = 0 : \lambda_{00} &= a^2, \\ k = 1, \ell = 0 : \lambda_{10} &= (1-a)^2 \geq a^2 \text{ since } a \in [0, 1/2], \\ k = 0, \ell = 1 : \lambda_{01} &= 1 + a^2. \end{aligned}$$

Therefore, λ_{00} is the lowest mode. \square

3.2.2. The Bakry-Emery method applied to the 2-dimensional torus

We consider the flow given by

$$\frac{\partial u}{\partial t} = \Delta u + (p-1) \frac{|\nabla u|^2}{u}$$

and observe that

$$\frac{d}{dt} \|u(t, \cdot)\|_{L^p(\mathbb{T}^2)}^2 = 0$$

on the one hand, and

$$\begin{aligned} & -\frac{1}{2} \frac{d}{dt} \left(\|\nabla u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2 - \lambda \|u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2 \right) \\ & = \|\Delta u\|_{L^2(\mathbb{T}^2)}^2 + (p-1) \iint_{\mathbb{T}^2} \Delta u \frac{|\nabla u|^2}{u} d\sigma - \lambda (2-p) \|\nabla u\|_{L^2(\mathbb{T}^2)}^2 \end{aligned}$$

on the other hand. Integrations by parts show that

$$\|\Delta u\|_{L^2(\mathbb{T}^2)}^2 = \|\text{Hess } u\|_{L^2(\mathbb{T}^2)}^2$$

and

$$\iint_{\mathbb{T}^2} \Delta u \frac{|\nabla u|^2}{u} d\sigma = -2 \iint_{\mathbb{T}^2} \text{Hess } u : \frac{\nabla u \otimes \nabla u}{u} d\sigma + \iint_{\mathbb{T}^2} \frac{|\nabla u|^4}{u^2} d\sigma.$$

Hence

$$\begin{aligned} & -\frac{1}{2} \frac{d}{dt} \left(\|\nabla u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2 - \lambda \|u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2 \right) \\ & = (2-p) \left(\|\Delta u\|_{L^2(\mathbb{T}^2)}^2 - \lambda \|\nabla u\|_{L^2(\mathbb{T}^2)}^2 \right) + (p-1) \left\| \text{Hess } u - \frac{\nabla u \otimes \nabla u}{u} \right\|_{L^2(\mathbb{T}^2)}^2. \end{aligned}$$

We know from the Poincaré inequality that

$$\|\Delta u\|_{L^2(\mathbb{T}^2)}^2 \geq \|\nabla u\|_{L^2(\mathbb{T}^2)}^2,$$

with optimal constant 1, so we can conclude in the case $1 \leq p < 2$ that $\|\nabla u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2 - \lambda \|u(t, \cdot)\|_{L^2(\mathbb{T}^2)}^2$ is monotone nonincreasing if $0 \leq \lambda \leq 1$. As a consequence, we have the following result.

Proposition 3.1. *For any $p \in [1, 2)$, we have*

$$\|\nabla u\|_{L^2(\mathbb{T}^2)}^2 + \|u\|_{L^p(\mathbb{T}^2)}^2 \geq \|u\|_{L^2(\mathbb{T}^2)}^2 \quad \forall u \in H^1(\mathbb{T}^2).$$

3.2.3. A tensorization result without magnetic potential

A result better than Proposition 3.1 follows from a tensorization argument that can be found in [11,17].

Proposition 3.2. *For any $p \in [1, 2)$, we have*

$$(2-p) \|\nabla u\|_{L^2(\mathbb{T}^2)}^2 + \|u\|_{L^p(\mathbb{T}^2)}^2 \geq \|u\|_{L^2(\mathbb{T}^2)}^2 \quad \forall u \in H^1(\mathbb{T}^2). \quad (3.8)$$

Moreover the factor $(2-p)$ is the optimal constant.

Proof. By taking on \mathbb{T}^2 a function depending only on $x \in \mathbb{S}^1$, it is clear that the constant in (3.8) cannot be improved. The proof of (3.8) can be done with the Bakry-Emery method applied to \mathbb{S}^1 and goes as follows.

Let us consider the flow given by

$$\frac{\partial u}{\partial t} = u'' + (p-1) \frac{|u'|^2}{u}$$

and observe that $\frac{d}{dt} \|u(t, \cdot)\|_{L^p(\mathbb{S}^1)}^2 = 0$ on the one hand, and

$$\begin{aligned} & -\frac{1}{2} \frac{d}{dt} \left(\|u'(t, \cdot)\|_{L^2(\mathbb{S}^1)}^2 - \lambda \|u(t, \cdot)\|_{L^2(\mathbb{S}^1)}^2 \right) \\ &= \|u''\|_{L^2(\mathbb{S}^1)}^2 + (p-1) \int_{\mathbb{S}^1} u'' \frac{|u'|^2}{u} d\sigma - \lambda (2-p) \|u'\|_{L^2(\mathbb{S}^1)}^2 \\ &= \|u''\|_{L^2(\mathbb{S}^1)}^2 + \frac{1}{3} (p-1) \int_{\mathbb{S}^1} \frac{|u'|^4}{u^2} d\sigma - \lambda (2-p) \|u'\|_{L^2(\mathbb{S}^1)}^2 \end{aligned}$$

on the other hand. Hence

$$-\frac{1}{2} \frac{d}{dt} \left(\|u'(t, \cdot)\|_{L^2(\mathbb{S}^1)}^2 - \lambda \|u(t, \cdot)\|_{L^2(\mathbb{S}^1)}^2 \right) \leq 0$$

if $\lambda(2-p) \leq 1$, because of the Poincaré inequality $\|u''\|_{L^2(\mathbb{S}^1)}^2 \geq \|u'\|_{L^2(\mathbb{S}^1)}^2$. Up to a sign change of λ , this computation also holds if $p > 2$ or if $p = -2$, as noticed in [14], and it is straightforward to extend it to the limit case $p = 2$ corresponding to the logarithmic Sobolev inequality.

According to [11, Proposition 3.1] or [17, Theorem 2.1] and up to a straightforward adaptation to the periodic setting, the optimal constant for the inequality on $\mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1$ is the same as for the inequality on \mathbb{S}^1 , provided $1 \leq p < 2$. \square

As a consequence of Proposition 3.2, we have the inequality

$$\|\nabla u\|_{L^2(\mathbb{T}^2)}^2 + \mu \|u\|_{L^p(\mathbb{T}^2)}^2 \geq \Lambda_{0,p}(\mu) \|u\|_{L^2(\mathbb{T}^2)}^2 \quad \forall u \in \mathbf{H}^1(\mathbb{T}^2), \quad (3.9)$$

where $\mu \mapsto \Lambda_{0,p}(\mu)$ is a concave monotone increasing function on $(0, +\infty)$ such that $\Lambda_{0,p}(\mu) = \mu$ for any $\mu \in (0, 1/(2-p))$.

3.2.4. A magnetic interpolation inequality in the flat torus

Now let us consider the generalization of (3.9) to the case $a \neq 0$.

Lemma 3.6. *Assume that $p \in [1, 2)$ and $a \in [0, 1/2]$. With the notation (3.7), there exists a concave monotone increasing function $\mu \mapsto \Lambda_{a,p}(\mu)$ on $(0, +\infty)$ such that $\lim_{\mu \rightarrow 0^+} \Lambda_{a,p}(\mu) = a^2$ where $\Lambda_{a,p}(\mu)$ is the optimal constant in the inequality*

$$\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{T}^2)}^2 + \mu \|u\|_{L^p(\mathbb{T}^2)}^2 \geq \Lambda_{a,p}(\mu) \|u\|_{L^2(\mathbb{T}^2)}^2 \quad \forall u \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{T}^2). \quad (3.10)$$

Moreover, we have that

$$\Lambda_{a,p}(\mu) \geq \mu + (1 - \mu(2-p))a^2 \quad \text{for any } \mu \leq \frac{1}{2-p}.$$

Proof. For an arbitrary $t \in (0, 1)$, we can write that

$$\begin{aligned} & \|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{T}^2)}^2 + \mu \|u\|_{L^p(\mathbb{T}^2)}^2 \\ & \geq t \left(\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{T}^2)}^2 - a^2 \|u\|_{L^2(\mathbb{T}^2)}^2 \right) \\ & \quad + (1-t) \left(\|\nabla |u|\|_{L^2(\mathbb{T}^2)}^2 + \frac{\mu}{1-t} \|u\|_{L^2(\mathbb{T}^2)}^p \right) + t a^2 \|u\|_{L^2(\mathbb{T}^2)}^2 \\ & \geq \left[(1-t) \Lambda_{0,p} \left(\frac{\mu}{1-t} \right) + t a^2 \right] \|u\|_{L^2(\mathbb{T}^2)}^2 \end{aligned}$$

using the diamagnetic inequality $\|\nabla_{\mathbf{A}} u\|_{L^2(\mathbb{T}^2)}^2 \geq \|\nabla |u|\|_{L^2(\mathbb{T}^2)}^2$. Inequality (3.9) applies with $\mu = 1/(2-p)$ and $t = 1 - \mu(2-p)$. \square

3.2.5. A symmetry result in the subquadratic regime

As an application of the results on magnetic rings of Theorem 3.1, we can prove a symmetry result for the optimal functions in (3.10) in the case $p < 2$. Let $\Lambda_{a,p}(\mu)$ be the optimal constant in (3.10).

Theorem 3.2. *Assume that $a \in [0, 1/2]$ and $p \in [1, 2)$. Then*

$$\Lambda_{a,p}(\mu) = \lambda_{a,p}(\mu) \quad \text{if } \mu \leq \frac{1}{p-2}$$

and any optimal function for (3.2) is then constant w.r.t. x . Moreover, $\Lambda_{a,p}(\mu) = a^2 + \mu$ if and only if $\mu(2-p) + 4a^2 \leq 1$ and equality in (3.10) is then achieved only by the constants.

Proof. Let us use the notation $\int f dx := \frac{1}{2\pi} \int_{-\pi}^{\pi} f dx$ in order to denote a normalized integration with respect to the single variable x , where y is considered as a parameter. For almost every $x \in \mathbb{S}^1$ we can apply (3.2) to the function $\psi(x, \cdot)$ and get

$$\begin{aligned} & \|\nabla_{\mathbf{A}} \psi\|_{L^2(\mathbb{T}^2)}^2 + \mu \|\psi\|_{L^p(\mathbb{T}^2)}^2 \\ & \geq \|\partial_x \psi\|_{L^2(\mathbb{T}^2)}^2 + \lambda_{a,p}(\mu) \|\psi\|_{L^2(\mathbb{T}^2)}^2 + \mu \|\psi\|_{L^p(\mathbb{T}^2)}^2 - \mu \int \left(\int |\psi|^p dy \right)^{\frac{2}{p}} dx \end{aligned}$$

Let us define $u := |\psi|$, $v(x) := \left(\int |u(x, y)|^p dy \right)^{1/p}$ and observe that

$$|v_x| = v^{1-p} \int u^{p-1} u_x dy \leq v^{1-p} \left(\int u^p dy \right)^{\frac{p-1}{p}} \left(\int |u_x|^2 dy \right)^{\frac{1}{2}} \left(\int 1 dy \right)^{\frac{1}{2} - \frac{p-1}{p}}$$

by Hölder's inequality, under the condition $p \leq 2$, that is,

$$|v_x|^2 \leq \int |u_x|^2 dy \leq \int |\partial_x \psi|^2 dy.$$

We conclude that if $\mu \leq 1/(2-p)$,

$$\int_{\mathbb{S}^1} |v_x|^2 d\sigma + \mu \left(\int_{\mathbb{S}^1} |v|^p d\sigma \right)^{2/p} - \mu \int_{\mathbb{S}^1} |v|^2 d\sigma + \lambda_{a,p}(\mu) \|\psi\|_{L^2(\mathbb{T}^2)}^2 \geq \lambda_{a,p}(\mu) \|\psi\|_{L^2(\mathbb{T}^2)}^2$$

using (3.1). The equality is achieved by functions v which are constant w.r.t. x and Theorem 3.1 applies. \square

3.3. Magnetic Hardy inequalities in dimensions 2 and 3

In this section, we draw some consequences of our results on magnetic rings of Section 3.1. Here $d\sigma$ denotes the uniform probability measure on \mathbb{S}^1 . The method relies on Keller-Lieb-Thirring dual estimates.

3.3.1. Keller-Lieb-Thirring inequalities on the circle

As in [13], by duality we obtain a spectral estimate.

Proposition 3.3. *Assume that $a \in [0, 1/2]$ and $p \in [1, 2)$. If ϕ is a nonnegative potential such that $\phi^{-1} \in L^q(\mathbb{S}^1)$, then the lowest eigenvalue λ_1 of $-(\partial_y - ia)^2 + \phi$ is bounded from below according to*

$$\lambda_1 \geq \lambda_{a,p} \left(\|\phi^{-1}\|_{L^q(\mathbb{S}^1)}^{-1} \right)$$

and equality is achieved by a constant potential ϕ if $\|\phi^{-1}\|_{L^q(\mathbb{S}^1)}^{-1} (2-p) + 4a^2 \leq 1$.

Proof. Using Hölder's inequality with exponents $2/(2-p)$ and $2/p$, we get that

$$\|\psi\|_{L^p(\mathbb{S}^1)}^2 = \left(\int_{\mathbb{S}^1} \phi^{-\frac{p}{2}} (\phi |\psi|^2)^{\frac{p}{2}} d\sigma \right)^{2/p} \leq \|\phi^{-1}\|_{L^q(\mathbb{S}^1)} \int_{\mathbb{S}^1} \phi |\psi|^2 d\sigma$$

with $q = p/(2 - p)$, and with $\mu = \|\phi^{-1}\|_{L^q(\mathbb{S}^1)}^{-1}$,

$$\begin{aligned} \int_{\mathbb{S}^1} |\psi' - i a \psi|^2 d\sigma + \int_{\mathbb{S}^1} \phi |\psi|^2 d\sigma &\geq \int_{\mathbb{S}^1} |\psi' - i a \psi|^2 d\sigma + \mu \|\psi\|_{L^p(\mathbb{S}^1)}^2 \\ &\geq \lambda_{a,p}(\mu) \int_{\mathbb{S}^1} |\psi|^2 d\sigma. \end{aligned} \quad (3.11)$$

If ϕ is constant, then there is equality in Hölder's inequality. \square

The spectral estimate (3.11) is of a different nature than (2.10) because the potential energy and the magnetic kinetic energy have the same sign. By considering the threshold case $\mu(2 - p) + 4a^2 = 1$, we obtain an interesting estimate.

Corollary 3.1. *Let $a \in [0, 1/2]$, $p \in (1, 2)$ and $q = p/(2 - p)$. If ϕ is a nonnegative potential such that $\phi^{-1} \in L^q(\mathbb{S}^1)$, then*

$$\begin{aligned} \int_{\mathbb{S}^1} |\psi' - i a \psi|^2 d\sigma + \frac{1 - 4a^2}{2 - p} \|\phi^{-1}\|_{L^q(\mathbb{S}^1)} \int_{\mathbb{S}^1} \phi |\psi|^2 d\sigma \\ \geq \left(\frac{1 - 4a^2}{2 - p} + a^2 \right) \|\psi\|_{L^2(\mathbb{S}^1)}^2 \quad \forall \psi \in H^1(\mathbb{S}^1). \end{aligned}$$

3.3.2. Magnetic Hardy-type inequalities in dimensions two and three

Let us denote by $\theta \in [-\pi, \pi)$ the angular coordinate associated with $x \in \mathbb{R}^2$. As in [14], we can deduce a Hardy-type inequality for Aharonov-Bohm magnetic potentials in dimension $d = 2$.

Corollary 3.2. *Let \mathbf{A} as in (1.1), $a \in [0, 1/2]$, $p \in (1, 2)$ and $q = p/(2 - p)$. If ϕ is a nonnegative potential such that $\phi^{-1} \in L^q(\mathbb{S}^1)$ with $\|\phi^{-1}\|_{L^q(\mathbb{S}^1)} = 1$, then for any complex valued function $\psi \in H^1(\mathbb{R}^2)$ we have*

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \frac{1 - 4a^2}{2 - p} \int_{\mathbb{R}^2} \frac{\phi(\theta)}{|x|^2} |\psi(x)|^2 dx \geq \left(\frac{1 - 4a^2}{2 - p} + a^2 \right) \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx.$$

Let us consider cylindrical coordinates $(\rho, \theta, z) \in \mathbb{R}^+ \times [0, 2\pi) \times \mathbb{R}$ such that $|x|^2 = \rho^2 + z^2$. In this system of coordinates the magnetic kinetic energy is

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx = \int_{\mathbb{R}^3} \left(\left| \frac{\partial \psi}{\partial \rho} \right|^2 + \frac{1}{\rho^2} \left| \frac{\partial \psi}{\partial \theta} - i a \psi \right|^2 + \left| \frac{\partial \psi}{\partial z} \right|^2 \right) dx$$

where $d\mu := \rho d\rho d\theta dz$. The following result was proved in [22, Section 2.2].

Lemma 3.7. *For any $\psi \in H^1(\mathbb{R}^3)$, we have*

$$\iiint_{\mathbb{R}^+ \times [0, 2\pi) \times \mathbb{R}} \left(\left| \frac{\partial \psi}{\partial \rho} \right|^2 + \left| \frac{\partial \psi}{\partial z} \right|^2 \right) d\mu \geq \frac{1}{4} \iiint_{\mathbb{R}^+ \times [0, 2\pi) \times \mathbb{R}} \frac{|\psi|^2}{\rho^2 + z^2} d\mu \quad \forall \psi \in H^1(\mathbb{R}^3).$$

Proof. We give an elementary proof. Assume that ψ is smooth and has compact support. The inequality follows from the expansion of the square

$$\iiint_{\mathbb{R}^+ \times [0, 2\pi) \times \mathbb{R}} \left(\left| \frac{\partial \psi}{\partial \rho} + \frac{\rho \psi}{2(\rho^2 + z^2)} \right|^2 + \left| \frac{\partial \psi}{\partial z} + \frac{z \psi}{2(\rho^2 + z^2)} \right|^2 \right) d\mu \geq 0 \quad \forall \psi \in H^1(\mathbb{R}^3)$$

and of an integration by parts of the cross terms. \square

Lemma 3.7 is an improved version of the standard Hardy inequality in the sense that the left-hand side of the inequality does not involve the angular part of the kinetic energy. A consequence of Corollary 3.1 and Lemma 3.7 is a Hardy-like estimate in dimension $d = 3$. For the angular part we argue as in Corollary 3.2. Details of the proof are left to the reader.

Theorem 3.3. *Let \mathbf{A} as in (1.2), $a \in [0, 1/2]$, $p \in (1, 2)$ and $q = p/(2 - p)$. If ϕ is a potential such that $\phi^{-1} \in L^q(\mathbb{S}^1)$ with $\|\phi^{-1}\|_{L^q(\mathbb{S}^1)} = 1$, then for any complex valued function $\psi \in H^1(\mathbb{R}^3)$ we have*

$$\begin{aligned} \int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx + \frac{1 - 4a^2}{2 - p} \int_{\mathbb{R}^3} \frac{\phi(\theta)}{\rho^2} |\psi(x)|^2 dx \\ \geq \frac{1}{4} \int_{\mathbb{R}^3} \frac{|\psi|^2}{|x|^2} dx + \left(\frac{1 - 4a^2}{2 - p} + a^2 \right) \int_{\mathbb{R}^3} \frac{|\psi|^2}{\rho^2} dx. \end{aligned}$$

A simple case is $\phi \equiv 1$, for which we obtain that

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \frac{1}{4} \int_{\mathbb{R}^3} \frac{|\psi|^2}{|x|^2} dx + a^2 \int_{\mathbb{R}^3} \frac{|\psi|^2}{|\rho|^2} dx \quad \forall \psi \in H_{\mathbf{A}}^1(\mathbb{R}^3).$$

4. Aharonov-Bohm magnetic interpolation inequalities in \mathbb{R}^2

Magnetic interpolation inequalities on \mathbb{R}^2 are considered without weights in Section 4.1. Weights are then introduced as in [7] in order to prove the new magnetic Caffarelli-Kohn-Nirenberg inequality of Corollary 4.1 in Section 4.2 and a magnetic Hardy inequality on \mathbb{R}^2 in Theorem 4.2 (Section 4.3).

4.1. Magnetic interpolation inequalities without weights

Let us consider on \mathbb{R}^2 the Aharonov-Bohm magnetic potential \mathbf{A} given by (1.1). Using the *diamagnetic inequality*

$$|\nabla_{\mathbf{A}} \psi|^2 \geq |\nabla |\psi||^2 \quad \text{a.e. in } \mathbb{R}^2$$

and, for any $p \in (2, \infty)$ and $\lambda > 0$, the Gagliardo-Nirenberg inequality

$$\|\nabla \psi\|_{L^2(\mathbb{R}^2)}^2 + \lambda \|\psi\|_{L^p(\mathbb{R}^2)}^2 \geq C_p \lambda^{\frac{p}{2}} \|\psi\|_{L^2(\mathbb{R}^2)}^2 \quad \forall \psi \in H^1(\mathbb{R}^2) \cap L^p(\mathbb{R}^2) \quad (4.1)$$

with optimal constant C_p , we deduce that

$$\|\nabla_{\mathbf{A}} \psi\|_{L^2(\mathbb{R}^2)}^2 + \lambda \|\psi\|_{L^2(\mathbb{R}^2)}^2 \geq \mu_{a,p}(\lambda) \|\psi\|_{L^p(\mathbb{R}^2)}^2 \quad \forall \psi \in H_a^1(\mathbb{R}^2). \quad (4.2)$$

See [13, Section 3] for details. Here $\mu_{a,p}(\lambda)$ is the optimal constant in (4.2) for any given a, p and λ and, as a function of λ , $\mu_{a,p}(\lambda)$ is monotone increasing and concave. Notice that right-hand sides in (4.1) and (4.2) involve norms with respect to Lebesgue's measure. It turns out that $\mu_{a,p}(\lambda)$ is equal to the best constant of the non-magnetic problem.

Proposition 4.1. *Let $a \in \mathbb{R}$ and $p \in (2, \infty)$. The optimal constant in (4.2) is*

$$\mu_{a,p}(\lambda) = C_p \lambda^{\frac{p}{2}} \quad \forall \lambda > 0$$

and equality is not achieved on $H^1(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ if $a \in \mathbb{R} \setminus \mathbb{Z}$.

Proof. By construction we know that $\mu_{a,p}(\lambda) \geq C_p \lambda^{p/2}$. By taking an optimal function ψ for (4.1) and considering $\psi_n(x) = \psi(x + n\mathbf{e})$ with $n \in \mathbb{N}$ and $\mathbf{e} \in \mathbb{S}^1$, we see that there is equality.

Let us prove by contradiction that equality is not achieved. If $\psi \in H^1(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ is optimal, let $\phi = e^{i a \theta} \psi$. Since

$$\|\nabla_{\mathbf{A}} \psi\|_{L^2(\mathbb{R}^2)}^2 = \int_{\mathbb{R}^2} |\partial_r \psi|^2 + \frac{|\partial_\theta \phi|^2}{|x|^2} dx$$

and equality in (4.1) is achieved by functions with a constant phase only, this means that $\partial_\theta \phi = 0$ a.e., a contradiction with the periodicity of ψ with respect to $\theta \in [0, 2\pi)$ if $a \notin \mathbb{Z}$. \square

Proposition 4.1 means that the Aharonov-Bohm magnetic potential plays no role in non-weighted interpolation inequalities. This is why it is natural to introduce weighted norms with adapted scaling properties.

4.2. Magnetic Caffarelli-Kohn-Nirenberg inequalities in \mathbb{R}^2

The *Caffarelli-Kohn-Nirenberg inequality*

$$\int_{\mathbb{R}^2} \frac{|\nabla v|^2}{|x|^{2a}} dx \geq C_a \left(\int_{\mathbb{R}^2} \frac{|v|^p}{|x|^{bp}} dx \right)^{2/p} \quad \forall v \in \mathcal{D}(\mathbb{R}^2) \quad (4.3)$$

has been established in [9] and, earlier, in [25]. The exponent $\mathbf{b} = \mathbf{a} + 2/p$ is determined by the scaling invariance and as p varies in $(2, \infty)$, the parameters \mathbf{a} and \mathbf{b} are such that $\mathbf{a} < \mathbf{b} \leq \mathbf{a} + 1$ and $\mathbf{a} < 0$. The case $\mathbf{a} > 0$ can be considered in an appropriate functional space after a Kelvin-type transformation: see [10,16], but we will not consider this case here. As noticed for instance in [16], by considering $v(x) = |x|^a u(x)$, Ineq. (4.3) is equivalent to the *Hardy-Sobolev inequality*

$$\int_{\mathbb{R}^2} |\nabla u|^2 dx + \mathbf{a}^2 \int_{\mathbb{R}^2} \frac{|u|^2}{|x|^2} dx \geq C_a \left(\int_{\mathbb{R}^2} \frac{|u|^p}{|x|^2} dx \right)^{2/p} \quad \forall u \in \mathcal{D}(\mathbb{R}^2). \quad (4.4)$$

The optimal functions for (4.3) are radially symmetric if and only if

$$\mathbf{b} \geq \mathbf{b}_{\text{FS}}(\mathbf{a}) := \mathbf{a} - \frac{\mathbf{a}}{\sqrt{1 + \mathbf{a}^2}}$$

according to [23,15]. We refer to [7] for more details and for the proof of the following magnetic Hardy-Sobolev inequality.

Theorem 4.1 ([7]). *Let $a \in [0, 1/2]$, \mathbf{A} as in (1.1) and $p > 2$. For any $\lambda > -a^2$, there is an optimal function $\lambda \mapsto \mu(\lambda)$ which is monotone increasing and concave such that*

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + \lambda \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx \geq \mu(\lambda) \left(\int_{\mathbb{R}^2} \frac{|\psi|^p}{|x|^2} dx \right)^{2/p} \quad \forall \psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{R}^2). \quad (4.5)$$

If $a \in [0, 1/2]$, the optimal function in (4.5) is

$$\psi(x) = (|x|^\alpha + |x|^{-\alpha})^{-\frac{2}{p-2}} \quad \forall x \in \mathbb{R}^2, \quad \text{with} \quad \alpha = \frac{p-2}{2} \sqrt{\lambda + a^2},$$

up to a scaling and a multiplication by a constant, if

$$\lambda \leq \lambda_\star := 4 \frac{1 - 4a^2}{p^2 - 4} - a^2.$$

Conversely, if $a \in [0, 1/2]$ and $\lambda > \lambda_\bullet$ with

$$\lambda_\bullet := \frac{8 \left(\sqrt{p^4 - a^2 (p-2)^2 (p+2) (3p-2) + 2} \right) - 4p(p+4)}{(p-2)^3 (p+2)} - a^2,$$

there is symmetry breaking, i.e., the optimal functions are not radially symmetric.

An explicit computation shows that $\lambda_\star < \lambda_\bullet$ for any $a \in (0, 1/2)$, and so there is a zone where we do not know whether the optimal functions in (4.5) are symmetric or not. Nevertheless, as shown in [7], the values of λ_\star and λ_\bullet are numerically very close to each other. If $\lambda \leq \lambda_\star$, the expression of $\mu(\lambda)$ is explicit and given by

$$\mu(\lambda) = \frac{p}{2} (2\pi)^{1-\frac{2}{p}} (\lambda + a^2)^{1+\frac{2}{p}} \left(\frac{2\sqrt{\pi} \Gamma(\frac{p}{p-2})}{(p-2) \Gamma(\frac{p}{p-2} + \frac{1}{2})} \right)^{1-\frac{2}{p}}.$$

See [7, Appendix] for the details of the computation of the constant.

Inspired by the equivalence of (4.3) and (4.4), we prove that the magnetic Hardy-Sobolev inequality (4.5) is equivalent to an interpolation inequality of Caffarelli-Kohn-Nirenberg type in the presence of the Aharonov-Bohm magnetic field.

Corollary 4.1 (Magnetic Caffarelli-Kohn-Nirenberg inequality). *Let $p \in (2, +\infty)$ and \mathbf{A} as in (1.1) for some $a \in [0, 1/2]$ and $\mathbf{a} \leq 0$. With μ as in Theorem 4.1, for any $\gamma < \mathbf{a}^2 + a^2$, we have that*

$$\int_{\mathbb{R}^2} \frac{|\nabla_{\mathbf{A}} \phi|^2}{|x|^{2\mathbf{a}}} dx \geq \gamma \int_{\mathbb{R}^2} \frac{|\phi|^2}{|x|^{2\mathbf{a}+2}} dx + \mu(\mathbf{a}^2 - \gamma) \left(\int_{\mathbb{R}^2} \frac{|\phi|^p}{|x|^{\mathbf{a}p+2}} dx \right)^{2/p} \quad \forall \phi \in \mathcal{D}(\mathbb{R}^2; \mathbb{C})$$

and $\mu(\lambda)$ with $\lambda = \mathbf{a}^2 - \gamma$ is the optimal constant.

The cases of symmetry and symmetry breaking in Theorem 4.1 have their exact counterpart in Corollary 4.1. Details are left to the reader.

Proof. Let us consider the function $\phi(x) = |x|^a \psi(x)$ and observe that

$$\int_{\mathbb{R}^2} \frac{|\nabla_{\mathbf{A}} \phi|^2}{|x|^{2a}} dx = \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx + a^2 \int_{\mathbb{R}^2} \frac{|\psi|^2}{|x|^2} dx$$

and conclude by applying (4.5) to ψ with $\lambda = a^2 - \gamma$. \square

4.3. A magnetic Hardy inequality in \mathbb{R}^2

Another consequence of Theorem 4.1 is the following *magnetic Keller-Lieb-Thirring inequality*, which can be found in [7, Theorem 1]. Let $q = p/(p-2)$. The ground state energy λ_1 of the magnetic Schrödinger operator $-\Delta_{\mathbf{A}} - \phi$ on \mathbb{R}^2 is such that

$$\lambda_1(-\Delta_{\mathbf{A}} - \phi) \geq -\lambda(\mu) \quad \text{where} \quad \mu = \left(\int_{\mathbb{R}^2} |\phi|^q |x|^{2(q-1)} dx \right)^{1/q} \quad (4.6)$$

and $\mu \mapsto \lambda(\mu)$ is a convex monotone increasing function on \mathbb{R}^+ such that $\lim_{\mu \rightarrow 0^+} \lambda(\mu) = -a^2$, defined as the inverse of $\lambda \mapsto \mu(\lambda)$ of Theorem 4.1. Again $\lambda(\mu)$ is optimal in (4.6) and the cases of symmetry and symmetry breaking are in correspondence with the ones of Theorem 4.1.

Alternatively, let us consider a function ϕ on \mathbb{R}^2 . We can estimate an associated magnetic Schrödinger energy from below by

$$\int_{\mathbb{R}^2} \left(|\nabla_{\mathbf{A}} \psi|^2 - \tau \frac{\phi}{|x|^2} |\psi|^2 \right) dx \geq \int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx - \tau \left(\int_{\mathbb{R}^2} \frac{|\phi|^q}{|x|^2} dx \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^2} \frac{|\psi|^p}{|x|^2} dx \right)^{\frac{2}{p}}$$

by Hölder's inequality, with $q = p/(p-2)$, for an arbitrary parameter $\tau > 0$. For an appropriate choice of τ , we obtain the following result.

Theorem 4.2 (A magnetic Hardy inequality). *Let $q \in (1, 2)$ and \mathbf{A} as in (1.1) for some $a \in [0, 1/2]$. Then for any function $\phi \in L^q(\mathbb{R}^2 |x|^{-2} dx)$, we have*

$$\int_{\mathbb{R}^2} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \mu(0) \left(\int_{\mathbb{R}^2} \frac{|\phi|^q}{|x|^2} dx \right)^{-\frac{1}{q}} \int_{\mathbb{R}^2} \frac{\phi}{|x|^2} |\psi|^2 dx \quad \forall \psi \in H_{\mathbf{A}}^1(\mathbb{R}^2),$$

where $\mu(\cdot)$ is the best constant in (4.5). Finally, when $a^2 \leq 4/(12 + p^2)$, we know the value of $\mu(0)$ explicitly:

$$\mu(0) = \frac{p}{2} (2\pi)^{1-\frac{2}{p}} a^{2+\frac{4}{p}} \left(\frac{2\sqrt{\pi} \Gamma(\frac{p}{p-2})}{(p-2) \Gamma(\frac{p}{p-2} + \frac{1}{2})} \right)^{1-\frac{2}{p}}.$$

5. Aharonov-Bohm magnetic Hardy inequalities in \mathbb{R}^3

In this section we address the issue of improved magnetic Hardy inequalities with the Aharonov-Bohm magnetic potential in dimension $d = 3$ as defined by (1.2). Our results improve upon [19, Section V.B], including the case of a constant magnetic field.

5.1. An improved Hardy inequality with radial symmetry

In [19, Section V.B], it is proved that for all $a > 0$, there is a constant $\mathcal{C}(a)$ such that $\mathcal{C}(a) = a^2$ if $a \in [0, 1/2]$ and

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \left(\frac{1}{4} + \mathcal{C}(a)\right) \int_{\mathbb{R}^3} \frac{|\psi|^2}{|x|^2} dx \quad \forall \psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{R}^3). \quad (5.1)$$

If we allow for an angular dependence, we have the following result.

Theorem 5.1. *Let \mathbf{A} as in (1.2), $a \in [0, 1/2]$ and $q \in (1, +\infty)$. Then, for all $\phi \in L^q(\mathbb{S}^2)$,*

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \int_{\mathbb{R}^3} \left(\frac{1}{4} + \frac{\mu_{a,p}(0)}{\|\phi\|_{L^q(\mathbb{S}^2)}} \phi(\omega) \right) \frac{|\psi|^2}{|x|^2} dx \quad \forall \psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{R}^3).$$

Here $\omega = x/|x|$ and $\mu_{a,p}$ is defined as in Proposition 2.2.

In the case $a \in [0, 1/2]$, according to Proposition 2.2, we find in the limit case as $p \rightarrow 2_+$ that $\mu_{a,2}(0) \geq \Lambda_a = a(a+1)$ and improve the estimate (5.1) to $\mathcal{C}(a) = a(a+1)$ if $\phi \equiv 1$.

Proof. Let us use spherical coordinates $(r, \omega) \in [0, +\infty) \times \mathbb{S}^2$. The result follows from an expansion of the square and an integration by parts in

$$0 \leq \int_0^{+\infty} \left| \partial_r \psi + \frac{1}{2r} \psi \right|^2 r^2 dr = \int_0^{+\infty} |\partial_r \psi|^2 r^2 dr - \frac{1}{4} \int_0^{+\infty} |\psi|^2 dr$$

for the radial part of the Dirichlet integral, and from Corollary 2.2 for the angular part. \square

5.2. An improved Hardy inequality with cylindrical symmetry

The improved Hardy inequality (without angular kinetic energy) of Lemma 3.7 and (2.9) can be combined into the following improved Hardy inequality in presence of a magnetic potential.

Theorem 5.2. *Let \mathbf{A} as in (1.2), $a \in [0, 1/2]$, $p > 2$, $q = p/(p-2)$ and $\phi \in L^q(\mathbb{S}^1)$. For any $\psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{R}^3)$, we have*

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \frac{1}{4} \int_{\mathbb{R}^3} \frac{|\psi|^2}{|x|^2} dx + \frac{\mu_{a,p}(0)}{\|\phi\|_{L^q(\mathbb{S}^1)}} \iiint_{\mathbb{R}^+ \times [0, 2\pi) \times \mathbb{R}} \frac{\phi(\theta)}{\rho^2} |\psi(\rho, \theta, z)|^2 d\mu.$$

Notice that the inequality is a strict improvement upon the Hardy inequality without a magnetic potential combined with the diamagnetic inequality. A simple case which is particularly illuminating is $\phi \equiv 1$ with $a^2 \leq 1/(p+2)$ so that $\mu_{a,p}(0) = a^2$ according to Proposition 2.1, in which case we obtain that

$$\int_{\mathbb{R}^3} |\nabla_{\mathbf{A}} \psi|^2 dx \geq \frac{1}{4} \int_{\mathbb{R}^3} \frac{|\psi|^2}{|x|^2} dx + a^2 \int_{\mathbb{R}^3} \frac{|\psi|^2}{|\rho|^2} dx \quad \forall \psi \in \mathbf{H}_{\mathbf{A}}^1(\mathbb{R}^3).$$

Acknowledgments

This research has been partially supported by the project *EFI*, contract ANR-17-CE40-0030 (D.B., J.D.) of the French National Research Agency (ANR), by the PDR (FNRS) grant T.1110.14F and the ERC AdG 2013 339958 “Complex Patterns for Strongly Interacting Dynamical Systems - COMPAT” grant (D.B.), by the RSF grant 18-11-00032 (A.L.) and by the NSF grant DMS-1600560 (M.L.). The authors thank an anonymous referee for constructive remarks which significantly improved the presentation of the results.

© 2020 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] Y. AHARONOV AND D. BOHM, *Significance of electromagnetic potentials in the quantum theory*, Physical Review, 115 (1959), pp. 485–491.
- [2] J. E. AVRON, I. W. HERBST, AND B. SIMON, *Schrödinger operators with magnetic fields: III. Atoms in homogeneous magnetic field*, Communications in Mathematical Physics, 79 (1981), pp. 529–572.
- [3] D. BAKRY AND M. ÉMERY, *Inégalités de Sobolev pour un semi-groupe symétrique*, C. R. Acad. Sci. Paris Sér. I Math., 301 (1985), pp. 411–413.
- [4] H. BATELAAN AND A. TONOMURA, *The Aharonov–Bohm effects: Variations on a subtle theme*, Physics Today, 62 (2009), pp. 38–43.
- [5] W. BECKNER, *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. (2), 138 (1993), pp. 213–242.
- [6] M.-F. BIDAUT-VÉRON AND L. VÉRON, *Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations*, Invent. Math., 106 (1991), pp. 489–539.
- [7] D. BONHEURE, J. DOLBEAULT, M. J. ESTEBAN, A. LAPTEV, AND M. LOSS, *Symmetry results in two-dimensional inequalities for Aharonov-Bohm magnetic fields*, Comm. Math. Phys., 375 (2020), pp. 2071–2087.
- [8] D. BONHEURE, M. NYS, AND J. VAN SCHAFTINGEN, *Properties of ground states of nonlinear Schrödinger equations under a weak constant magnetic field*, J. Math. Pures Appl. (9), 124 (2019), pp. 123–168.
- [9] L. CAFFARELLI, R. KOHN, AND L. NIRENBERG, *First order interpolation inequalities with weights*, Compositio Math., 53 (1984), pp. 259–275.
- [10] F. CATRINA AND Z.-Q. WANG, *On the Caffarelli-Kohn-Nirenberg inequalities: sharp constants, existence (and nonexistence), and symmetry of extremal functions*, Comm. Pure Appl. Math., 54 (2001), pp. 229–258.
- [11] D. CHAFAÏ, *Entropies, convexity, and functional inequalities: on Φ -entropies and Φ -Sobolev inequalities*, J. Math. Kyoto Univ., 44 (2004), pp. 325–363.
- [12] J. DOLBEAULT, M. J. ESTEBAN, AND A. LAPTEV, *Spectral estimates on the sphere*, Anal. PDE, 7 (2014), pp. 435–460.
- [13] J. DOLBEAULT, M. J. ESTEBAN, A. LAPTEV, AND M. LOSS, *Interpolation inequalities and spectral estimates for magnetic operators*, Ann. Henri Poincaré, 19 (2018), pp. 1439–1463.
- [14] J. DOLBEAULT, M. J. ESTEBAN, A. LAPTEV, AND M. LOSS, *Magnetic rings*, Journal of Mathematical Physics, 59 (2018), p. 051504.
- [15] J. DOLBEAULT, M. J. ESTEBAN, AND M. LOSS, *Rigidity versus symmetry breaking via nonlinear flows on cylinders and Euclidean spaces*, Invent. Math., 206 (2016), pp. 397–440.

- [16] J. DOLBEAULT, M. J. ESTEBAN, M. LOSS, AND G. TARANTELO, *On the symmetry of extremals for the Caffarelli-Kohn-Nirenberg inequalities*, Adv. Nonlinear Stud., 9 (2009), pp. 713–726.
- [17] J. DOLBEAULT AND X. LI, *Φ -Entropies: convexity, coercivity and hypocoercivity for Fokker-Planck and kinetic Fokker-Planck equations*, Mathematical Models and Methods in Applied Sciences, 28 (2018), pp. 2637–2666.
- [18] W. EHRENBERG AND R. SIDAY, *The refractive index in electron optics and the principles of dynamics*, Proceedings of the Physical Society. Section B, 62 (1949), p. 8.
- [19] T. EKHOLM AND F. PORTMANN, *A magnetic contribution to the Hardy inequality*, J. Math. Phys., 55 (2014), pp. 022101, 16.
- [20] L. ERDŐS, *Rayleigh-type isoperimetric inequality with a homogeneous magnetic field*, Calculus of Variations and Partial Differential Equations, 4 (1996), pp. 283–292.
- [21] P. EXNER, E. M. HARRELL, AND M. LOSS, *Optimal eigenvalues for some Laplacians and Schrödinger operators depending on curvature*, in Mathematical results in quantum mechanics (Prague, 1998), vol. 108 of Oper. Theory Adv. Appl., Birkhäuser, Basel, 1999, pp. 47–58.
- [22] L. FANELLI, D. KREJČIŘÍK, A. LAPTEV, AND L. VEGA, *On the improvement of the Hardy inequality due to singular magnetic fields*, Comm. Partial Differential Equations, 45 (2020), pp. 1202–1212.
- [23] V. FELLI AND M. SCHNEIDER, *Perturbation results of critical elliptic equations of Caffarelli-Kohn-Nirenberg type*, J. Differential Equations, 191 (2003), pp. 121–142.
- [24] T. HOFFMANN-OSTENHOF AND A. LAPTEV, *Hardy inequalities with homogeneous weights*, J. Funct. Anal., 268 (2015), pp. 3278–3289.
- [25] V. P. IL'IN, *Some integral inequalities and their applications in the theory of differentiable functions of several variables*, Mat. Sb. (N.S.), 54 (96) (1961), pp. 331–380.
- [26] A. LAPTEV AND T. WEIDL, *Hardy inequalities for magnetic Dirichlet forms*, in Mathematical results in quantum mechanics (Prague, 1998), vol. 108 of Oper. Theory Adv. Appl., Birkhäuser, Basel, 1999, pp. 299–305.
- [27] E. H. LIEB AND M. LOSS, *Analysis*, vol. 14 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2001.
- [28] G. SZEGŐ, *Orthogonal polynomials*, American Mathematical Society, Providence, R.I., fourth ed., 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.