

Stability for the logarithmic Sobolev inequality

Giovanni Brigati^a, Jean Dolbeault^{b,*}, Nikita Simonov^c

^a*Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400 Klosterneuburg, Austria*

^b*CEREMADE (CNRS UMR n° 7534), PSL University, Université Paris-Dauphine,
Place de Lattre de Tassigny, 75775 Paris 16, France*

^c*LJLL (CNRS UMR n° 7598), Sorbonne Université, 4 place Jussieu, 75005 Paris, France*

Abstract

This paper is devoted to stability results for the Gaussian logarithmic Sobolev inequality, with explicit stability constants.

Keywords: logarithmic Sobolev inequality, stability, log-concavity, heat flow, entropy, carré du champ, Poincaré inequality.

2020 MSC: Primary: [39B62](#); Secondary: [47J20](#), [49J40](#), [35A23](#), [35K85](#).

1. Introduction

1.1. Main result

Let $d \geq 1$, and let us consider the *Gaussian logarithmic Sobolev inequality*

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log \left(\frac{|u|^2}{\|u\|_{L^2(\mathbb{R}^d, d\gamma)}^2} \right) d\gamma \quad \forall u \in H^1(\mathbb{R}^d, d\gamma) \quad (\text{LSI})$$

where $d\gamma = \gamma(x) dx$ is the normalized Gaussian probability measure with density

$$\gamma(x) = (2\pi)^{-\frac{d}{2}} e^{-\frac{1}{2}|x|^2} \quad \forall x \in \mathbb{R}^d.$$

According to [18, 4], equality in (LSI) is achieved by any function in the manifold

$$\mathcal{M} := \{w_{a,c} : (a, c) \in \mathbb{R}^d \times \mathbb{R}\} \quad \text{where} \quad w_{a,c}(x) = c e^{a \cdot x} \quad \forall x \in \mathbb{R}^d$$

and only by functions in \mathcal{M} .

The issue of *stability* in functional inequalities is either (a) to estimate the distance to \mathcal{M} by the deficit in (LSI), *i.e.*, by the difference of the two sides in the inequality, or (b) to obtain an *improved inequality*, *i.e.*, an improved constant in the

* Corresponding author

Email addresses: giovanni.brigati@ist.ac.at (Giovanni Brigati),
dolbeaul@ceremade.dauphine.fr (Jean Dolbeault), nikita.simonov@sorbonne-universite.fr
(Nikita Simonov)

inequality, under appropriate conditions. In fact, (b) amounts to establish (a) but for a restricted class of functions. More details will be given later. Our main result is in the spirit of (b). By two-homogeneity of (LSI), we may consider functions in

$$\mathcal{H} := \left\{ u \in H^1(\mathbb{R}^d, d\gamma) : \|u\|_{L^2(\mathbb{R}^d, d\gamma)} = 1 \right\}$$

without loss of generality and state an *improved (LSI) inequality* as follows.

Theorem 1. *Let $d \geq 1$. For any $\varepsilon > 0$ and $C > 0$, there is an explicit $\eta \in (0, 1)$ such that*

$$(1 - \eta) \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma$$

for any $u \in \mathcal{H}$ such that

$$\int_{\mathbb{R}^d} x |u|^2 d\gamma = 0 \tag{1}$$

and

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(x)|^2 |u(y)|^2 e^{\varepsilon|x-y|^2} d\gamma(x) d\gamma(y) \leq C. \tag{2}$$

The constant η depends only on C and ε , but not directly on the dimension d . This is consistent with [21]. In the special case of compactly supported functions, η can be estimated explicitly as a function of the size of the support.

To the best of our knowledge, it is the first time such that Condition (2) appears in the study of stability of (LSI). Simpler conditions can also be given: for instance, (2) holds true for any function $u \in \mathcal{H}$ such that

$$\int_{\mathbb{R}^d} e^{2\varepsilon|x|^2} |u|^2 d\gamma \leq \sqrt{C}, \tag{3}$$

as a consequence of the estimate $e^{\varepsilon|x-y|^2} \leq e^{2\varepsilon|x|^2} e^{2\varepsilon|y|^2}$.

Our strategy goes as follows. Under Condition (2), a density $h(t = 0, \cdot) = |u|^2$ evolved by the *Ornstein-Uhlenbeck* flow on \mathbb{R}^d ,

$$\frac{\partial h}{\partial t} = \mathcal{L}h, \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^d, \tag{4}$$

where $\mathcal{L}h := \Delta h - x \cdot \nabla h$ denotes the *Ornstein-Uhlenbeck* operator, is such that the measure $d\mu_T = h(T, \cdot) d\gamma$ satisfies a Poincaré inequality after some finite time $T \geq 0$, as a consequence of the results of H.-B. Chen, S. Chewi, and J. Niles-Weed in [21]. According to M. Fathi, E. Indrei, and M. Ledoux in [33], this measure also satisfies an improved version of (LSI). Using a *backward in time* argument based on the *carré du champ* method, we deduce an improved (LSI) inequality for the initial density $|u|^2$ from the improved (LSI) inequality for the measure $d\mu_T$. This completes the sketch of the proof of Theorem 1.

This paper is organized as follows. The remainder of the introduction is dedicated to a partial review of the literature. In Section 2, we discuss the distinction between improved inequalities and stability results, collect various observations, give

stability results for which we do not claim much originality (but with new proofs and simple, explicit constants) and state the key *backward in time* argument. Section 3 is devoted to the proof of Theorem 1. In Section 4, we state an additional stability result for log-concave functions and establish a dimension-free estimate for compactly supported functions that arises as a consequence of Theorem 1.

1.2. A partial review of the literature

In 1975, the *Gaussian logarithmic Sobolev inequality (LSI)* was shown in [36] by L. Gross to be equivalent to the hypercontractivity of the Ornstein–Uhlenbeck semigroup. A scale invariant (but dimension-dependent) version of the Euclidean form of the inequality appeared in [55, Theorem 2], which was already known from [49, Inequality (2.3)] in dimension $d = 1$. By using a duality argument, (LSI) is also related to a Keller type estimate, see [34]. The reader interested in further historical details is invited to refer to [53, Section 1.3.2] and also to [52, 50, 51] for further background references in information theory. More references can also be found in [29, 33]. The optimality case in the inequality has been characterized in [18], but can also be deduced from [4]. The logarithmic Sobolev inequality can be seen as a limit case of a family of Gagliardo–Nirenberg–Sobolev inequalities, as observed in [24] in the Euclidean setting, or as a large dimension limit of the Sobolev inequality according to [8] (see also [17] for detailed computations and further references). For books on (LSI), we refer to [1, 37, 47, 5].

In a classical result on stability in functional inequalities, G. Bianchi and H. Eggenell proved in [9] that the deficit in the Sobolev inequality measures the $\dot{H}^1(\mathbb{R}^d, dx)$ distance to the manifold of the optimisers. The estimate has been made constructive in [26], where a new $L^2(\mathbb{R}^d, d\gamma)$ stability result for the logarithmic Sobolev inequality is also established (also see [38, 40] for further negative and positive results in other norms, for instance in strong norms like $H^1(\mathbb{R}^d, d\gamma)$). To our knowledge, the first result of stability for the logarithmic Sobolev inequality is a reinforcement of the inequality due to E. Carlen in [18] where he introduces an additional term involving the Wiener transform. An improved (LSI) inequality appeared in [33] based on a Mehler formula for the Ornstein–Uhlenbeck semigroup, which gives deficit estimates in various distances for functions inducing a Poincaré inequality. It is a result in the spirit of (b) and [33] is crucial for our proof of Theorem 1. Under the condition $\|xu\|_{L^2(\mathbb{R}^d, d\gamma)} = \sqrt{d}$, a stability result measured by a relative Fisher information is also given in [29], on the basis of simple scaling properties of the Euclidean form of the logarithmic Sobolev inequality. For sequential stability results in strong norms, we refer to [40] when assuming a bound on u in $L^4(\mathbb{R}^d, d\gamma)$ and to [38] when assuming a bound on $|x|^2 u$ in $L^2(\mathbb{R}^d, d\gamma)$. An earlier sequential stability result in $L^2(\mathbb{R}^d, d\gamma)$ based on a Fourier technique was also given in [40] under Condition (3). Stability according to other notions of distance has been studied in [44, 43, 35]. Stability results where the distance to \mathcal{M} appears with a non-optimal exponent are known for instance from [12, Theorem 1.1] where it is deduced from the HWI inequality due to F. Otto and C. Villani [46]. Such estimates have even been refined in [30]. There are now several other proofs. Various stability results have also been proved in Wasserstein’s distance: we refer to [41, 12, 33, 40, 42, 13, 30, 38] and in particular to [39,

Theorem 1.3] for result with sharp exponent under an exponential moment condition. Stability in logarithmic Sobolev inequality can be related to deficit in Gaussian isoperimetry and we refer to [12] for an introduction to early results in this direction, [6] for a sharp, dimension-free quantitative Gaussian isoperimetric inequality, and [30] for recent results and further references.

In this paper, we carefully distinguish (a) stability results where a distance to \mathcal{M} is controlled by the deficit, and (b) improved inequalities under appropriate constraints. Even if functions are normalized and centered, this is in some cases not enough for obtaining improved inequalities as shown in [38]. In fact, many counter-examples to stability are known, involving Wasserstein's distance for instance in [41, 12, 33, 40, 42], weaker distances like p -Wasserstein, or stronger norms like L^p or H^1 : see for instance [40, 38]. The classical counter-examples that apply to our setting are those of [42, Theorem 1.3] and [30, Theorem 4] but, as already noted in [40], they are based on the fact that the second moment diverges along a sequence of test functions. In case of Theorem 1, this is forbidden by Assumption (2).

A large part of our intuition comes from the fact that the heat flow (or the Ornstein–Uhlenbeck flow) preserves log-concavity: see, e.g., [15, 48]. Log-concavity is a natural property in this framework for several reasons, see for instance [23]. By J. Cheeger's inequality [20], from log-concavity follows an explicit Poincaré inequality that we can use to establish an improved form of (LSI). Also see [19] and references therein. However, what really matters is the Poincaré inequality, as noted by M. Fathi, E. Indrei and M. Ledoux in [33]. Condition (2) comes from the result of [21] obtained by H.-B. Chen, S. Chewi, and J. Nilsson-Weed: such a Poincaré inequality holds under the evolution of the Ornstein–Uhlenbeck flow, after some explicit delay. The study of Poincaré inequalities evolved under (4) (or equivalently under convolutions with Gaussian measures) has many applications, cf. [21]. The study of this question has been initiated in [56, 57] and further investigated in [54, 7, 21].

In [33, Theorem 1], the assumption is that $|u|^2 d\gamma$ satisfies a Poincaré inequality. According to [10, Equation (4.3)], this also implies the *exponential moment* condition $\int_{\mathbb{R}^d} e^{\theta|x|} u^2 d\gamma < \infty$, for some $\theta > 0$. Condition (2) is stronger than the *exponential moment* condition, the classical example being the measure $|u(x)|^2 d\gamma(x) = e^{-|x|} dx$ which satisfies a Poincaré inequality but not (2). On the other hand, (2) is, in some cases, less restrictive than the assumption of [33, Theorem 1], for instance, in the case of a compactly supported function u with several disconnected components. Whether Theorem 1 can be extended, eventually with additional restrictions, to the case of an *exponential moment* condition is, to our knowledge, an open question.

Bounds on the Poincaré constant of a probability measure may depend on the dimension and degenerate for large dimensions according to [19, 22]. As a consequence, the same issue arises for the improvement of (LSI) of [33, Theorem 1]. For strongly log-concave measures $u^2 d\gamma$, one has a Poincaré inequality with a dimension-independent estimate of the constant according to [4], but this class falls into the much wider class considered in Theorem 1.

2. Preliminary observations, an important tool and some simple consequences

2.1. A discussion on stability estimates and improved inequalities

Let us define the *deficit* functional of (LSI) by

$$\delta[u] := \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log \left(\frac{|u|^2}{\|u\|_{L^2(\mathbb{R}^d, d\gamma)}^2} \right) d\gamma.$$

The goal of *stability estimates* is to find a notion of distance d , an explicit constant $\beta > 0$ and an explicit exponent $\alpha > 0$ such that

$$\delta[u] \geq \beta \inf_{w \in \mathcal{M}} d(u, w)^\alpha \quad \forall u \in \mathcal{H}. \quad (5)$$

It is known from [26, Corollary 1.2] that for some explicit, dimension-independent constant $\beta > 0$, one has

$$\delta[u] \geq \beta \inf_{w \in \mathcal{M}} \|u - w\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \quad \forall u \in H^1(\mathbb{R}^d, d\gamma). \quad (6)$$

In this paper we consider *improved inequalities* in the form

$$\delta[u] \geq \beta d(u, 1)^\alpha \quad \forall u \in \mathcal{H}. \quad (7)$$

Any estimate of α and β for (7) is also an estimate for (5), as $\inf_{w \in \mathcal{M}} d(u, w) \leq d(u, 1)$, because $w \equiv 1 \in \mathcal{M}$. When $d(u, w) = \|u - w\|_{L^2(\mathbb{R}^d, d\gamma)}$, $\alpha = 2$, and β as in (6), Inequalities (5) and (7) are in fact equivalent if u is nonnegative, normalized and centred as proven in [38, Lemma 1]. The equivalence between (5) and (7) does not hold for $d(u, w) = \|\nabla u - \nabla w\|_{L^2(\mathbb{R}, d\gamma)}$ because the best possible exponent α in (5) and (7) differ, as the following example shows. Assume that $d = 1$ and consider the functions

$$u_\varepsilon(x) = 1 + \varepsilon x \quad \forall x \in \mathbb{R}^d \quad (8)$$

in the limit as $\varepsilon \rightarrow 0$. With $d(u, w) = \|u' - w'\|_{L^2(\mathbb{R}, d\gamma)}$, we have

$$d(u_\varepsilon, 1)^2 = \|u'_\varepsilon\|_{L^2(\mathbb{R}, d\gamma)}^2 = \varepsilon^2 \quad \text{and} \quad \delta[u_\varepsilon] = \frac{1}{2} \varepsilon^4 + O(\varepsilon^6) \quad \text{as} \quad \varepsilon \rightarrow 0.$$

Hence, the best we can hope for in (7) written with $w_u = 1$ is $\beta = 1/2$ and $\alpha = 4$. On the other hand, using the test function $w_{a_\varepsilon, c_\varepsilon} \in \mathcal{M}$ where $a_\varepsilon = 2\varepsilon$ and $\log c_\varepsilon = -a_\varepsilon^2/4$, we obtain

$$\inf_{w \in \mathcal{M}} d(u_\varepsilon, w)^2 \leq d(u_\varepsilon, w_{a_\varepsilon, c_\varepsilon})^2 = \frac{1}{2} \varepsilon^4 + O(\varepsilon^6) = \delta[u_\varepsilon] + O(\varepsilon^6) \quad \text{as} \quad \varepsilon \rightarrow 0,$$

which would allow for $\beta = 1$ and $\alpha = 2$ in (5). Up to a Gaussian Poincaré inequality, this is compatible with the fact that (6) still yields a stability estimate with $\alpha = 2$, which is the smallest possible exponent: see [38, Theorem 2]. On the other hand, the example of u_ε given by (8) suggests that, for non-centred functions, $\alpha = 4$ is the best possible exponent in (7) for the distance $d(u, w) = \|\nabla u - \nabla w\|_{L^2(\mathbb{R}, d\gamma)}$.

To establish improved inequalities with strong notions of distance d , one needs, unlike in (5), an *additional condition*. In the case of $d(u, w) = \|\nabla u - \nabla w\|_{L^2(\mathbb{R}^d, d\gamma)}$, at least a control on the second-order moment $K = \int_{\mathbb{R}^d} |x|^2 u^2 d\gamma$ is necessary: in [40, 42], for any $K > d$, the authors build sequences $(u_\varepsilon)_{\varepsilon>0}$ of functions in \mathcal{H} which satisfy (1) such that

$$\lim_{\varepsilon \rightarrow 0^+} \delta[u_\varepsilon] = 0 \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0^+} W_2^2(u_\varepsilon^2 \gamma, \gamma) = \frac{1}{2}(K - d)$$

where W_2 denotes the 2-Wasserstein distance. By (LSI) and the Talagrand inequality,

$$2 \|\nabla u_\varepsilon\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \int_{\mathbb{R}^d} |u_\varepsilon|^2 \log |u_\varepsilon|^2 d\gamma \geq \frac{1}{2} W_2^2(u_\varepsilon^2 \gamma, \gamma),$$

so that (7) cannot hold along such a sequence with a distance d that controls W_2 . Moreover, in [38, Theorem 1], E. Indrei proves that, for any sequence $(u_n)_{n \in \mathbb{N}}$ of functions in \mathcal{H} such that (1) holds, one can deduce that $\lim_{n \rightarrow +\infty} \|\nabla u_n\|_{L^2(\mathbb{R}^d, d\gamma)} = 0$ from $\lim_{n \rightarrow +\infty} \delta[u_n] = 0$ if and only if the condition $\lim_{n \rightarrow +\infty} \|x u_n\|_{L^2(\mathbb{R}^d, d\gamma)} = \sqrt{d}$ is satisfied. Nevertheless, several results are available under a second moment condition [12, 29], fourth moment condition [38] and in the class of probability densities which satisfy a Poincaré inequality [33]. See also Corollary 4 for a result under a second moment condition.

Let us comment on our main result Theorem 1. We aim at results in the strongest possible notion of distance, *i.e.*, $d(u, w) = \|u - w\|_{H^1(\mathbb{R}^d, d\gamma)}$ where $w \in \mathcal{M}$. From Theorem 1 and by the Gaussian Poincaré inequality, we find

$$\delta[u] \geq \eta \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \frac{\eta}{2} \|u - 1\|_{H^1(\mathbb{R}^d, d\gamma)}^2$$

for all $u \in H^1(\mathbb{R}^d, d\gamma)$ satisfying (1) and (2): this proves (7) for $\alpha = 2$ and $\beta = \eta/2$. Notice that by two-homogeneity of δ , $\alpha = 2$ is the best possible exponent. Assumption (1) is crucial, as illustrated by the functions u_ε defined by (8). Condition (2) is sharp in the following sense: for any $0 < \varepsilon < 1/2$, there exists a sequence $(u_{\varepsilon, n})_{n \in \mathbb{N}}$ of functions in \mathcal{H} built as in [40, 42] and satisfying (1) such that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} e^{2\varepsilon|x|^2} |u_{\varepsilon, n}|^2 d\gamma = +\infty, \quad \liminf_{n \rightarrow \infty} \|\nabla u_{\varepsilon, n}\|_{L^2(\mathbb{R}^d, d\gamma)}^2 > 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \delta[u_{\varepsilon, n}] = 0,$$

so that $\lim_{n \rightarrow \infty} \delta[u_{\varepsilon, n}] / \|\nabla u_{\varepsilon, n}\|_{L^2(\mathbb{R}^d, d\gamma)}^2 = 0$. How to fill the gap between a control on the second-order moment, which is a necessary for (7), and the much more restrictive condition of Theorem 1, is an open question.

2.2. The Ornstein–Uhlenbeck equation and the carré du champ method

Let us recall some classical results on (4). If $h_0 \in L^1(\mathbb{R}^d, d\gamma)$ is nonnegative, then there exists a unique nonnegative weak solution to (4) (see for instance [32]). The

two key properties of the Ornstein–Uhlenbeck operator $\mathcal{L}h = \Delta h - x \cdot \nabla h$ are

$$\int_{\mathbb{R}^d} h_1 (\mathcal{L}h_2) d\gamma = - \int_{\mathbb{R}^d} \nabla h_1 \cdot \nabla h_2 d\gamma \quad \text{and} \quad [\nabla, \mathcal{L}]h = -\nabla h.$$

As a consequence, we obtain the two identities

$$\int_{\mathbb{R}^d} (\mathcal{L}h)^2 d\gamma = \int_{\mathbb{R}^d} \|\text{Hess } h\|^2 d\gamma + \int_{\mathbb{R}^d} |\nabla h|^2 d\gamma \quad (9)$$

and

$$\int_{\mathbb{R}^d} \mathcal{L}h \frac{|\nabla h|^2}{h} d\gamma = -2 \int_{\mathbb{R}^d} \text{Hess } h : \frac{\nabla h \otimes \nabla h}{h} d\gamma + \int_{\mathbb{R}^d} \frac{|\nabla h|^4}{h^2} d\gamma, \quad (10)$$

where $\text{Hess } h = (\nabla \otimes \nabla)h$ is the *Hessian matrix* of h . Here we use the following notations. If a and b take values in \mathbb{R}^d , $a \otimes b$ denotes the matrix $(a_i b_j)_{1 \leq i, j \leq d}$. With matrix valued $m = (m_{i,j})_{1 \leq i, j \leq d}$ and $n = (n_{i,j})_{1 \leq i, j \leq d}$, we define $m : n = \sum_{i,j=1}^d m_{i,j} n_{i,j}$ and $\|m\|^2 = m : m$. If h is a nonnegative solution of (4), we also notice that $v = \sqrt{h}$ solves

$$\frac{\partial v}{\partial t} = \mathcal{L}v + \frac{|\nabla v|^2}{v}. \quad (11)$$

Let us fix $\|v\|_{L^2(\mathbb{R}^d, d\gamma)} = 1$. The *entropy* and the *Fisher information*, respectively defined by

$$\mathcal{E}[v] := \int_{\mathbb{R}^d} |v|^2 \log |v|^2 d\gamma \quad \text{and} \quad \mathcal{I}[v] := \int_{\mathbb{R}^d} |\nabla v|^2 d\gamma,$$

evolve along the flow (11) according to

$$\frac{d}{dt} \mathcal{E}[v(t, \cdot)] = -4 \mathcal{I}[v(t, \cdot)] \quad \text{and} \quad \frac{d}{dt} \mathcal{I}[v(t, \cdot)] = -2 \int_{\mathbb{R}^d} \left((\mathcal{L}v)^2 + \mathcal{L}v \frac{|\nabla v|^2}{v} \right) d\gamma$$

if v solves (11). Using (9) and (10), we obtain the classical expression of the *carré du champ* method

$$\frac{d}{dt} \mathcal{I}[v(t, \cdot)] + 2 \mathcal{I}[v(t, \cdot)] = -2 \int_{\mathbb{R}^d} \left\| \text{Hess } v - \frac{\nabla v \otimes \nabla v}{v} \right\|^2 d\gamma \quad (12)$$

as for instance in [3, 27, 5]. By writing that

$$\frac{d}{dt} \delta[v(t, \cdot)] \leq 0 \quad \text{and} \quad \lim_{t \rightarrow +\infty} \delta[v(t, \cdot)] = 0,$$

we recover the standard proof of the *entropy – entropy production* inequality (LSI), by the *carré du champ* method of [4].

Several of the above expression can be rephrased in terms of the *pressure variable*

$$P := -\log h = -2 \log v$$

using the following elementary identities

$$\begin{aligned}\nabla v &= -\frac{1}{2}\sqrt{h}\nabla P, & \frac{\nabla v \otimes \nabla v}{v} &= \frac{1}{4}\sqrt{h}\nabla P \otimes \nabla P, \\ \text{Hess } v &= -\frac{1}{2}\sqrt{h}\text{Hess } P + \frac{1}{4}\sqrt{h}\nabla P \otimes \nabla P,\end{aligned}$$

so that, by taking into account $v\nabla P = -2\nabla v$ and $h = v^2$, we have

$$\mathcal{I}[v] = \frac{1}{4} \int_{\mathbb{R}^d} |\nabla P|^2 h d\gamma \quad \text{and} \quad \int_{\mathbb{R}^d} \left\| \text{Hess } v - \frac{\nabla v \otimes \nabla v}{v} \right\|^2 d\gamma = \frac{1}{4} \int_{\mathbb{R}^d} \|\text{Hess } P\|^2 h d\gamma.$$

2.3. A backward in time estimate

Although rather elementary, this key tool of our paper is based on the following observation: if a solution h of (4) is such that (LSI) holds for $v = \sqrt{h}$ with an improved constant at some time $T > 0$, then this is also the case for the initial datum. A similar approach was used in [14, Lemma 2.9] in the case of a different flow.

Lemma 2. *Let $u \in \mathcal{H}$ be such that (1) holds and consider the solution v of (11) with initial datum u . If for some $T > 0$ we have $\delta[w] \geq c \|\nabla w\|_{L^2(\mathbb{R}^d, d\gamma)}^2$ with $w := v(T, \cdot)$ for some $c \in (0, 1)$, then*

$$\delta[u] \geq c e^{-2T} \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2.$$

Notice that $c \geq 1$ is impossible unless $c = 1$ and w is a constant.

Proof. The (LSI) ensures that $\mathcal{Q}(t) := \mathcal{I}[v(t, \cdot)] / \mathcal{E}[v(t, \cdot)] \geq 1/2$ for all $t \geq 0$. By our assumption, $(1 - c)\mathcal{Q}(T) \geq 1/2$ and we learn from (12) that

$$\frac{d\mathcal{Q}}{dt} \leq 2\mathcal{Q}(2\mathcal{Q} - 1).$$

The conclusion follows from an integration on $(0, T)$, which shows that

$$\mathcal{Q}(0) \geq \frac{1}{2} \frac{1}{1 - e^{-2T} c}.$$

2.4. Some simple stability estimates □

In this section, we collect various stability estimates and provide new proofs or explicit estimates which are new. We put the emphasis on the use of the Ornstein–Uhlenbeck equation (4) and on the improvements based on the *carré du champ* method.

2.4.1. Improvements under moment constraints

In standard computations of the *carré du champ* method, one usually drops the Hessian terms, as those in right-hand side of (12). Keeping track of the remainder terms provides us with improvements as shown in [2, 25, 28] in various interpolation inequalities but generically fails in the case of the logarithmic Sobolev inequality. We remedy to this issue by introducing moment constraints.

Lemma 3. *If $\sqrt{h} = v \in \mathcal{H} \cap H^2(\mathbb{R}^d, d\gamma)$ is such that (1) holds and $xv \in L^2(\mathbb{R}^d, d\gamma)$, then*

$$4 \cdot \mathcal{I}[v] \leq \sqrt{d \int_{\mathbb{R}^d} \|\text{Hess } P\|^2 h \, d\gamma} + \int_{\mathbb{R}^d} |x|^2 h \, d\gamma - d.$$

Proof. Using $h \nabla P = -\nabla h$, we obtain

$$4 \cdot \mathcal{I}[v] = \int_{\mathbb{R}^d} |\nabla P|^2 h \, d\gamma = - \int_{\mathbb{R}^d} \nabla P \cdot \nabla h \, d\gamma = \int_{\mathbb{R}^d} h (\mathcal{L}P) \, d\gamma.$$

After recalling that $\mathcal{L}P = \Delta P - x \cdot \nabla P$, we deduce that

$$- \int_{\mathbb{R}^d} h x \cdot \nabla P \, d\gamma = \int_{\mathbb{R}^d} x \cdot \nabla h \, d\gamma = \int_{\mathbb{R}^d} h (|x|^2 - d) \, d\gamma = \int_{\mathbb{R}^d} |v|^2 (|x|^2 - d) \, d\gamma$$

using an integration by parts, which proves

$$4 \cdot \mathcal{I}[v] \leq \int_{\mathbb{R}^d} (\Delta P) h \, d\gamma.$$

The use of the Cauchy-Schwarz inequality and the arithmetic-geometric inequality

$$(\Delta P)^2 \leq d \|\text{Hess } P\|^2$$

completes the proof. \square

With the estimate of Lemma 3 on $\mathcal{I}[v]$, we have the following result.

Corollary 4. *Let $\Psi(s) := s - \frac{d}{4} \log(1 + \frac{4}{d}s)$. For all $u \in \mathcal{H}$ such that (1) holds and $\|xu\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \leq d$, we have the stability estimate*

$$\delta[u] \geq \Psi\left(\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2\right). \quad (13)$$

Notice that Ψ is such that $\Psi(s) = \frac{2}{d}s^2 + o(s^2)$ as $s \rightarrow 0_+$, so that $\alpha = 4$ is the minimal admissible exponent in Inequality (7), at least for results obtained by this method.

Proof of Corollary 4. Let $h = v^2$ be the solution of (4) with initial datum $h_0 = u^2$. Since $x \mapsto (|x|^2 - d)$ is an eigenfunction of \mathcal{L} with corresponding eigenvalue -2 and \mathcal{L} is self-adjoint on $L^2(\mathbb{R}^d, d\gamma)$, we have that $\mathcal{K}(t) := \int_{\mathbb{R}^d} (|x|^2 - d) h \, d\gamma$ satisfies

$$\frac{d\mathcal{K}}{dt} = \int_{\mathbb{R}^d} (|x|^2 - d) (\mathcal{L}h) \, d\gamma = \int_{\mathbb{R}^d} h \mathcal{L}(|x|^2 - d) \, d\gamma = -2\mathcal{K}. \quad (14)$$

The sign of $t \mapsto \mathcal{K}(t)$ is conserved and in particular we have that $\int_{\mathbb{R}^d} |x|^2 |v|^2 \, d\gamma \leq d$ for any $t \geq 0$. For any $i = 1, 2, \dots, d$, we also notice that $x \mapsto x_i$ is also an eigenfunction of \mathcal{L} with corresponding eigenvalue -1 so that

$$\frac{d}{dt} \int_{\mathbb{R}^d} x h \, d\gamma = - \int_{\mathbb{R}^d} x h \, d\gamma$$

and, as a consequence $\int_{\mathbb{R}^d} x h(t, \cdot) d\gamma = 0$ for all $t \geq 0$ because $\int_{\mathbb{R}^d} x h_0 d\gamma = 0$.

For smooth enough solutions, we deduce from Lemma 3, (12) and (LSI) that

$$\frac{d}{dt} \mathcal{J}[v(t, \cdot)] + 2 \mathcal{J}[v(t, \cdot)] \leq -\frac{8}{d} \mathcal{J}^2[v(t, \cdot)] \leq \frac{1}{2d} \frac{d}{dt} (\mathcal{E}[v(t, \cdot)])^2$$

if v solves (11). The fact that $\lim_{t \rightarrow +\infty} \mathcal{J}[v(t, \cdot)] = 0$ follows from a Gronwall estimate relying on $\frac{d}{dt} \mathcal{J}[v(t, \cdot)] \leq -2 \mathcal{J}[v(t, \cdot)]$ and $\lim_{t \rightarrow +\infty} \mathcal{E}[v(t, \cdot)] = 0$ is obtained as a consequence of (LSI). Since $t \mapsto \mathcal{J}[v(t, \cdot)] - \frac{1}{2} \mathcal{E}[v(t, \cdot)] - \frac{1}{2d} (\mathcal{E}[v(t, \cdot)])^2$ is monotone nonincreasing with limit 0 as $t \rightarrow +\infty$, we conclude that it is nonnegative for any $t \geq 0$ and, as a special case, at $t = 0$, thus proving that

$$\delta[u] = \mathcal{J}[u] - \frac{1}{2} \mathcal{E}[u] \geq \frac{1}{2d} (\mathcal{E}[u])^2. \quad (15)$$

A better estimate goes as follows. Let

$$\Phi(s) := \frac{d}{4} \left(e^{\frac{2}{d}s} - 1 \right) \quad \forall s \geq 0.$$

Using $\frac{d}{dt} \mathcal{E}[v(t, \cdot)] = -4 \mathcal{J}[v(t, \cdot)]$, we notice that

$$\frac{d}{dt} \left(\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)]) \right) \leq -\frac{8}{d} \left(\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)]) \right) \mathcal{J}[v(t, \cdot)].$$

As before, we know that

$$\lim_{t \rightarrow +\infty} \left(\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)]) \right) = 0.$$

Moreover, Gronwall estimates show that $\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)])$ cannot change sign and an asymptotic expansion as $t \rightarrow +\infty$ as in [16, Appendix B.4] is enough to obtain that $\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)])$ takes nonnegative values for $t > 0$ large enough. Altogether, we conclude that

$$\mathcal{J}[v(t, \cdot)] - \Phi(\mathcal{E}[v(t, \cdot)]) \geq 0$$

for any $t \geq 0$ and, as a particular case, at $t = 0$ for $v(0, \cdot) = u$. This provides us with the estimate

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \geq \frac{d}{4} \left(e^{\frac{2}{d} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma} - 1 \right).$$

In the general case, one can get rid of the $H^2(\mathbb{R}^d, d\gamma)$ regularity of Lemma 3 by a standard approximation scheme, which is classical and will not be detailed here.

As in [16], an estimate in a stronger norm is achieved as follows. The function Φ is convex increasing and, as such, invertible, so that we can also write

$$\Phi^{-1}(\mathcal{J}[u]) - \mathcal{E}[u] \geq 0.$$

This completes the proof of (13) with the convex monotone increasing function

$$\Psi(s) := s - \frac{1}{2} \Phi^{-1}(s) \quad \forall s \geq 0.$$

□

As far as we know, the above proof of Corollary 4 is new, but the result was known by other methods, as explained in Section 2.1. An inequality in the spirit of (13) is proved in [12, Theorem 1.1 and Inequality (1.8)] using probabilistic tools. Inequality (13) is also established in [29] using the scaling properties of the Euclidean version of (LSI) under the more restrictive second moment condition $\|x u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 = d$. In dimension $d = 1$ with a second moment finite but arbitrarily large, a stability result in a weak norm can be found in [39, Theorem 1.2]. If $\mathcal{K}[u] < 0$, i.e.,

$$\|x u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 < d,$$

further improvements can be achieved. Here is an example. With

$$\mathcal{J}[u] := \mathcal{I}[u] - \frac{1}{4} \mathcal{K}[u],$$

we notice that (LSI) can be recast as

$$\mathcal{J}[u] \geq \frac{1}{2} \left(\mathcal{E}[u] - \frac{1}{2} \mathcal{K}[u] \right).$$

Moreover, if v solves (11), then

$$\frac{d}{dt} \left(\mathcal{E}[v(t, \cdot)] - \frac{1}{2} \mathcal{K}[v(t, \cdot)] \right) = -4 \mathcal{J}[v(t, \cdot)]$$

and using Lemma 3, the estimate in the proof of Corollary 4 becomes

$$\begin{aligned} \frac{d}{dt} \mathcal{J}[v(t, \cdot)] + 2 \mathcal{J}[v(t, \cdot)] &= \frac{d}{dt} \mathcal{I}[v(t, \cdot)] + 2 \mathcal{J}[v(t, \cdot)] \\ &\leq -\frac{8}{d} \mathcal{J}^2[v(t, \cdot)] \leq \frac{1}{2d} \frac{d}{dt} \left(\mathcal{E}[v(t, \cdot)] - \frac{1}{2} \mathcal{K}[v(t, \cdot)] \right)^2. \end{aligned}$$

An integration from $t = 0$ to $+\infty$ shows that

$$\delta[u] \geq \frac{1}{2d} \left(\mathcal{E}[u] - \frac{1}{2} \mathcal{K}[u] \right)^2,$$

which is an improvement upon (15) under the assumption that $\mathcal{K}[u] < 0$, as we know that $\mathcal{E}[u] \geq 0$ by Jensen's inequality.

2.4.2. A result in the framework of log-concave densities

We say that a measure $d\mu$ with density $e^{-\psi}$ with respect to Lebesgue's measure is a *log-concave probability measure* if ψ is a convex function.

Lemma 5. *If $d\mu$ is a log-concave probability measure such that $\int_{\mathbb{R}^d} |x - x_\mu|^2 d\mu \leq K$ where $x_\mu = \int_{\mathbb{R}^d} x d\mu$, then we have the Poincaré inequality*

$$\int_{\mathbb{R}^d} |\nabla \varphi|^2 d\mu \geq \frac{1}{432K} \int_{\mathbb{R}^d} |\varphi|^2 d\mu \quad \forall \varphi \in H^1(\mathbb{R}^d, d\mu) \text{ such that } \int_{\mathbb{R}^d} \varphi d\mu = 0. \quad (16)$$

Proof. Let us denote by $\lambda_1(\mu)$ the first positive eigenvalue of $-\mathcal{L}_\psi$ where \mathcal{L}_ψ is the Ornstein–Uhlenbeck operator $\mathcal{L}_\psi := \Delta - \nabla \psi \cdot \nabla$. We learn from [11, Theorem 1.2] and [11, Ineq. (3.4)] that $432K\lambda_1(\mu) \geq 1$. \square

Lemma 6. *Let us consider a solution h of (4) with initial datum h_0 and assume that $d\mu_0 := h_0 d\gamma$ is a log-concave probability measure. Then $d\mu_t := h(t, \cdot) d\gamma$ is a log-concave probability measure for all $t \geq 0$.*

Proof. The function $g := h\gamma$ solves the Fokker-Planck equation

$$\frac{\partial g}{\partial t} = \Delta g + \nabla \cdot (xg)$$

and the function f such that

$$f(s, x) := (1 + 2s)^{-\frac{d}{2}} g\left(\frac{1}{2} \log(1 + 2s), \frac{x}{\sqrt{1 + 2s}}\right) \quad \forall (s, x) \in \mathbb{R}^+ \times \mathbb{R}^d \quad (17)$$

solves the heat equation

$$\frac{\partial f}{\partial s} = \Delta f \quad \forall (s, x) \in \mathbb{R}^+ \times \mathbb{R}^d. \quad (18)$$

Hence f can be represented using the heat kernel. According for instance to [48, 7], log-concavity is preserved under convolution, which completes the proof. \square

Lemma 7. *If $h \in H^1(\mathbb{R}^d, d\gamma)$ is such that $\int_{\mathbb{R}^d} xh d\gamma = 0$ and $P = -\log h$ is the pressure variable, then*

$$\int_{\mathbb{R}^d} \nabla P h d\gamma = 0.$$

Proof. The result follows from $\int_{\mathbb{R}^d} \nabla P h d\gamma = -\int_{\mathbb{R}^d} \nabla h d\gamma = \int_{\mathbb{R}^d} xh d\gamma = 0$. \square

Let

$$\mathcal{C}_\star(K) = 1 + \frac{1}{432K} \quad (19)$$

where $1/432 \approx 0.00231481$.

Proposition 8. *For all $u \in \mathcal{H}$ such that (1) holds, $\int_{\mathbb{R}^d} |x|^2 |u|^2 d\gamma = K$ and $u^2 \gamma$ is log-concave, with \mathcal{C}_\star defined by (19), we have*

$$\|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{1}{2} \mathcal{C}_\star(\max(K, d)) \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \geq 0.$$

Proof. The solution $h = v^2$ of (4) is such that $\int_{\mathbb{R}^d} x h d\gamma = 0$ and Lemma 7 applies. Since $h(t, \cdot)\gamma$ is log-concave for any $t \geq 0$ by Lemma 6, we can apply (16) with $f = \partial P / \partial x_i$ for any $i = 1, 2, \dots, d$ and obtain

$$\int_{\mathbb{R}^d} \|\text{Hess } P\|^2 h d\gamma \geq \frac{1}{432 \max(K, d)} \int_{\mathbb{R}^d} |\nabla P|^2 h d\gamma$$

because, using (14), $d + \mathcal{K} = d + (K - d)e^{-2t} \leq \max(K, d)$. It follows from (12) that

$$\frac{d}{dt} \mathcal{I}[v(t, \cdot)] + 2\mathcal{C}_\star(K) \mathcal{I}[v(t, \cdot)] \leq 0,$$

and the stability result is obtained as in the standard proof of the *entropy – entropy production inequality* (LSI) by the *carré du champ* method, cf. Section 2.2. \square

2.4.3. From compact support to log-concavity

The log-concavity property becomes true under the action of the flow of (4) after some delay t_\star for large classes of initial data. With the notation of Lemma 6, for any $R > 0$, we read from [45, Theorem 5.1] by K. Lee and J-L. Vázquez that $d\mu_t = |v(t, \cdot)|^2 d\gamma$ is log-concave for any

$$t \geq t_\star(R) := \log\left(\sqrt{R^2 + 1}\right) \quad (20)$$

if v solves (11) with a compactly supported initial datum u that is supported in a ball of radius $R > 0$. The reduction from (11) to the heat flow (18) goes as in the proof of Lemma 6. As a consequence, we know that (16) holds for any $t \geq t_\star(R)$.

Corollary 9. *Let $d \geq 1$ and assume that $u \in \mathcal{H}$ is compactly supported in a ball of radius $R > 0$. Then for all $u \in \mathcal{H}$ such that (1) holds, with \mathcal{C}_\star defined by (19), we have*

$$\int_{\mathbb{R}^d} |\nabla u|^2 d\gamma \geq \frac{\mathcal{C}}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma$$

with

$$\mathcal{C} = 1 + \frac{\mathcal{C}_\star(K_\star) - 1}{1 + R^2 \mathcal{C}_\star(K_\star)} \quad \text{and} \quad K_\star := \max\left(d, \frac{(d+1)R^2}{1+R^2}\right).$$

Proof. Corollary 9 follows from $t_\star(R)$ given by (20) so that $e^{-2t_\star(R)} = 1/(1+R^2)$, Proposition 8 applied at $t = t_\star(R)$ with $K = (d+1)R^2/(1+R^2) \geq \mathcal{K}(t_\star(R)) + d$ by (14), and Lemma 2 applied with $T = t_\star(R)$ and $c = 1 - 1/\mathcal{C}_\star(K_\star)$, so that

$$\delta[u] = \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 - \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 \log |u|^2 d\gamma \geq \frac{\mathcal{C}_\star(K_\star) - 1}{(1+R^2)\mathcal{C}_\star(K_\star)} \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2.$$

\square

3. Proof of Theorem 1

The proof of Theorem 1 is based on three ingredients:

1. A stability result for (LSI) obtained by M. Fathi, E. Indrei, and M. Ledoux in [33] for a special class of initial data satisfying a Poincaré inequality,
2. The result of [21] by H.-B. Chen, S. Chewi, and J. Niles-Weed which states that after a finite time $T \geq 0$, a solution to the Ornstein–Uhlenbeck flow is in the above class under Condition (2),
3. The backward in time argument based on the *carré du champ* method of Section 2.3, as in [14], which is used on the interval $[0, T)$.

3.1. Evolving a Poincaré inequality by the Ornstein–Uhlenbeck flow and application

In this section we collect some information about the Poincaré inequality for a measure $\nu^2 d\gamma$ where the function ν is evolving under the Ornstein–Uhlenbeck equation (11). As in [33], let us define $\mathcal{P}(\lambda)$, for any $\lambda > 0$, as the set of all functions u such that the measure $u^2 d\gamma$ is a probability measure which satisfies a Poincaré inequality

$$\int_{\mathbb{R}^d} |\nabla \varphi|^2 u^2 d\gamma \geq \lambda \int_{\mathbb{R}^d} |\varphi|^2 u^2 d\gamma \quad \forall \varphi \in \mathcal{H}_u$$

where \mathcal{H}_u is the space of the functions $\varphi \in H^1(\mathbb{R}^d, u^2 d\gamma)$ such that $\int_{\mathbb{R}^d} \varphi u^2 d\gamma = 0$. The following Lemma is a key step in obtaining [33, Theorem 1] and its proof can be found in [33, Section 2].

Lemma 10. *If $v(t, \cdot)$ solves (11) with initial datum $v(t=0, \cdot) = u$ such that $u \in \mathcal{P}(\lambda)$ for some $\lambda > 0$, then for any $t \geq 0$ we have the Poincaré inequality*

$$\int_{\mathbb{R}^d} |\nabla \varphi|^2 |v(t, \cdot)|^2 d\gamma \geq \frac{\lambda}{\lambda + (1 - \lambda) e^{-2t}} \int_{\mathbb{R}^d} |\varphi|^2 |v(t, \cdot)|^2 d\gamma \quad \forall \varphi \in \mathcal{H}_{v(t, \cdot)}.$$

Let us consider the function

$$\sigma(\lambda) := \frac{\lambda^2 - \lambda - \lambda \log \lambda}{(\lambda - 1)^2} \quad \forall \lambda \in (0, 1) \cup (1, +\infty)$$

and extend it by $\sigma(1) = 1/2$. We may notice that σ is monotone increasing and concave, with $\lim_{\lambda \rightarrow 0^+} \sigma(\lambda) = 0$ and $\lim_{\lambda \rightarrow +\infty} \sigma(\lambda) = 1$. An interesting consequence of Lemma 10 is the following stability estimate for (LSI).

Corollary 11 ([33]). *Let $\lambda > 0$ and $u \in \mathcal{H}$ satisfying Condition (1) and such that $u \in \mathcal{P}(\lambda)$, then we have that*

$$\delta[u] \geq \sigma(\lambda) \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2.$$

The proof of Corollary 11 relies on the same strategy as in the proof of Proposition 8, except that one has to use the Poincaré inequality of Lemma 10 to write that

$$\int_{\mathbb{R}^d} \|\text{Hess } P\|^2 h d\gamma \geq \frac{\lambda}{\lambda + (1 - \lambda) e^{-2t}} \int_{\mathbb{R}^d} |\nabla P|^2 h d\gamma \quad \forall t \geq 0.$$

The result follows from an integration on $t \in \mathbb{R}^+$ of (12) using the above inequality.

3.2. Gaussian convolutions of measures through the Ornstein-Uhlenbeck flow

For any $a > 1$ and $b > 0$, let us define the function

$$F(a, b) := \frac{1}{a} \left(\frac{a}{a-1} + b^{\frac{1}{a-1}} \right)^{-1}.$$

The following result is a rephrasing of the main result in [21] using the Ornstein-Uhlenbeck flow.

Lemma 12. *Let $v(t, \cdot)$ be a solution to (11) with $v(t=0, \cdot) = u \in \mathcal{H}$. If u satisfies (1) and (2) for some $\varepsilon > 0$ and $C > 0$, then*

$$v(t, \cdot) \in \mathcal{P}(\lambda(t)) \quad \text{with} \quad \lambda(t) := \varepsilon e^{2t} F(\varepsilon(e^{2t} - 1), C) \quad \forall t \geq t_\varepsilon := \frac{1}{2} \log \left(1 + \frac{1}{\varepsilon} \right). \quad (21)$$

Proof. As in the proof of Lemma 6, if the function f solves (18) with initial datum $|u|^2 \gamma$, then

$$\gamma(x) |v(t, x)|^2 = e^{dt} f(s, y) \quad \text{with} \quad s = \frac{1}{2} (e^{2t} - 1) \quad \text{and} \quad y = e^t x$$

by (17). By a direct application of [21, Theorem 2], we learn that $f(s, \cdot)/\sqrt{\gamma}$ is in $\mathcal{P}(\Lambda)$ with $\Lambda = \varepsilon F(2\varepsilon s, C)$ if $2s > 1/\varepsilon$, i.e., if $t = t(s) \geq t_\varepsilon$. Changing variables, if $\psi(y) = \varphi(e^{-t} y)$, we have

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla_x \varphi(x)|^2 |v(t, x)|^2 d\gamma &= e^{2t} \int_{\mathbb{R}^d} |\nabla \psi(y)|^2 f(s, y) dy, \\ \int_{\mathbb{R}^d} |\varphi(x)|^2 |v(t, x)|^2 d\gamma &= \int_{\mathbb{R}^d} |\psi(y)|^2 f(s, y) dy, \end{aligned}$$

for any $\varphi \in H^1(\mathbb{R}^d, v^2 d\gamma)$ with zero average with respect to the measure $v^2 d\gamma$. Since the function ψ has zero average with respect to the measure $f(s, y) dy$, the corresponding Poincaré inequality amounts to (21) written with $\lambda = \lambda(t) = e^{2t} \Lambda$. This concludes the proof. \square

3.3. Conclusion

We can now complete the proof of Theorem 1 as follows. We recall that $v(t, \cdot)$ solves (11) with initial datum $v(t=0, \cdot) = u$ such that (1) and (2) hold. By Corollary 10 and Lemma 12, we know that

$$\delta[v(t, \cdot)] \geq \sigma(\lambda(t)) \|\nabla v(t, \cdot)\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \quad \forall t \geq t_\varepsilon$$

with $\lambda(t)$ and t_ε given by (21). As a consequence of Lemma 2 applied with $T = t_\varepsilon$, we have

$$\delta[u] \geq \eta \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2 \quad \text{with} \quad \eta = \sup_{t > t_\varepsilon} \sigma(\lambda(t)) e^{-2t}. \quad (22)$$

\square

With the admissible choice of t satisfying $\varepsilon(e^{2t} - 1) = 2$, we obtain the simple lower estimate

$$\eta \geq \frac{\varepsilon}{2 + \varepsilon} \sigma \left(\frac{2 + \varepsilon}{2(2 + C)} \right).$$

4. Log-concavity and compactly supported functions

This section collects some additional results about stability for log-concave measures and compactly supported functions.

4.1. Stability results for log-concave measures

Here is a simple improvement of Proposition 8 based on Corollary 11.

Corollary 13. *Assume $\nu := u^2 d\gamma$ is a logarithmically concave probability measure with $u \in \mathcal{H}$, such that (1) is satisfied, and $\|xu\|_{L^2(\mathbb{R}^d, d\gamma)}^2 = K$. Then, the stability estimate*

$$\delta[u] \geq \sigma(\lambda) \int_{\mathbb{R}^d} |\nabla u|^2 d\gamma$$

holds with $1/\lambda = 432K$.

Proof. This result is a simple consequence of Lemma 5 and Corollary 11. \square

4.2. Compactly supported functions

From Corollary 9, we obtain an improved (LSI) with a bound which depends on the dimension d . Dimensional dependence is a huge topic in functional inequalities and we refer to [31] for further considerations in this direction. As a final remark, let us notice that Theorem 1 provides us with a dimension-free result.

Proposition 14. *If $u \in \mathcal{H}$ is supported in $B(0, R)$ and satisfies (1), then*

$$\delta[u] \geq \frac{\alpha}{1 + R^2} \|\nabla u\|_{L^2(\mathbb{R}^d, d\gamma)}^2$$

with $\alpha = ((1 + e) \log(1 + e) - e) / e^2 \approx 0.292973$.

Proof. We may notice that (2) is satisfied for every $\varepsilon > 0$, and $C = \exp(\varepsilon R^2)$. The result follows by taking the limit as $\varepsilon \rightarrow \infty$ in (22) with $e^{2t} = 1 + R^2$ and $\lambda(t) \geq (1 + e)^{-1}$. \square

Acknowledgements

The authors thank Max Fathi and Pierre Cardaliaguet for fruitful discussions and Emanuel Indrei for stimulating interactions. They also thank an anonymous referee for useful comments and suggestions which have led to an improvement of the manuscript. They also want to express their gratitude to the managing editor, L. Gross, for his encouragements and questions. G.B. has been funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 754362. This work has been (partially) supported by the Project Conviviality ANR-23-CE40-0003 of the French National Research Agency

© 2024 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] C. ANÉ, S. BLACHÈRE, D. CHAFAÏ, P. FOUGÈRES, I. GENTIL, F. MALRIEU, C. ROBERTO, AND G. SCHEFFER, *Sur les inégalités de Sobolev logarithmiques*, vol. 10, Société mathématique de France Paris, 2000.
- [2] A. ARNOLD AND J. DOLBEAULT, *Refined convex Sobolev inequalities*, J. Funct. Anal., 225 (2005), pp. 337–351.
- [3] A. ARNOLD, P. MARKOWICH, G. TOSCANI, AND A. UNTERREITER, *On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations*, Comm. Partial Differential Equations, 26 (2001), pp. 43–100.
- [4] D. BAKRY AND M. ÉMERY, *Diffusions hypercontractives*, in Séminaire de Probabilités XIX 1983/84, Springer, 1985, pp. 177–206.
- [5] D. BAKRY, I. GENTIL, AND M. LEDOUX, *Analysis and geometry of Markov diffusion operators*, vol. 348 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer, Cham, 2014.
- [6] M. BARCHIESI, A. BRANCOLINI, AND V. JULIN, *Sharp dimension free quantitative estimates for the Gaussian isoperimetric inequality*, Ann. Probab., 45 (2017), pp. 668–697.
- [7] J.-B. BARDET, N. GOZLAN, F. MALRIEU, AND P.-A. ZITT, *Functional inequalities for Gaussian convolutions of compactly supported measures: Explicit bounds and dimension dependence*, Bernoulli, 24 (2018).
- [8] W. BECKNER, *Sobolev inequalities, the Poisson semigroup, and analysis on the sphere \mathbb{S}^n* , Proc. Nat. Acad. Sci. U.S.A., 89 (1992), pp. 4816–4819.
- [9] G. BIANCHI AND H. EGNELL, *A note on the Sobolev inequality*, J. Funct. Anal., 100 (1991), pp. 18–24.
- [10] S. BOBKOV AND M. LEDOUX, *Poincaré’s inequalities and Talagrand’s concentration phenomenon for the exponential distribution*, Probab. Theory and Related Fields, 107 (1997), pp. 383–400.
- [11] S. G. BOBKOV, *Isoperimetric and analytic inequalities for log-concave probability measures*, Ann. Probab., 27 (1999), pp. 1903–1921.
- [12] S. G. BOBKOV, N. GOZLAN, C. ROBERTO, AND P.-M. SAMSON, *Bounds on the deficit in the logarithmic Sobolev inequality*, J. Funct. Anal., 267 (2014), pp. 4110–4138.
- [13] F. BOLLEY, I. GENTIL, AND A. GUILLIN, *Dimensional improvements of the logarithmic Sobolev, Talagrand and Brascamp–Lieb inequalities*, Ann. Probab., 46 (2018), pp. 261–301.
- [14] M. BONFORTE, J. DOLBEAULT, B. NAZARET, AND N. SIMONOV, *Stability in Gagliardo-Nirenberg-Sobolev inequalities: Flows, regularity and the entropy method*, Preprint arXiv: [2007.03674](https://arxiv.org/abs/2007.03674) and [hal-02887010](https://arxiv.org/abs/2208.02887), to appear in *Memoirs of the AMS*, (2024).
- [15] H. J. BRASCAMP AND E. H. LIEB, *On extensions of the Brunn-Minkowski and Prekopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation*, J. Funct. Anal., 22 (1976), pp. 366–389.
- [16] G. BRIGATI, J. DOLBEAULT, AND N. SIMONOV, *Logarithmic Sobolev and interpolation inequalities on the sphere: Constructive stability results*, Annales de l’Institut Henri Poincaré C, Analyse non linéaire, (2023), pp. 1–33.
- [17] ———, *On Gaussian interpolation inequalities*, C. R. Math. Acad. Sci. Paris, 362 (2024), pp. 21–44.
- [18] E. A. CARLEN, *Superadditivity of Fisher’s information and logarithmic Sobolev inequalities*, J. Funct. Anal., 101 (1991), pp. 194–211.
- [19] P. CATTIAUX AND A. GUILLIN, *Functional inequalities for perturbed measures with applications to log-concave measures and to some Bayesian problems*, Bernoulli, 28 (2022).
- [20] J. CHEEGER, *A lower bound for the smallest eigenvalue of the Laplacian*, in Problems in

- analysis (Sympos. in honor of Salomon Bochner, Princeton Univ., Princeton, N.J., 1969), Princeton Univ. Press, Princeton, N.J., 1970, pp. 195–199.
- [21] H.-B. CHEN, S. CHEWI, AND J. NILES-WEED, *Dimension-free log-Sobolev inequalities for mixture distributions*, J. Funct. Anal., 281 (2021), p. 109236.
 - [22] Y. CHEN, *An almost constant lower bound of the isoperimetric coefficient in the KLS conjecture*, Geom. Funct. Anal., 31 (2021), pp. 34–61.
 - [23] T. A. COURTADE, M. FATHI, AND A. PANANJADY, *Quantitative stability of the entropy power inequality*, IEEE Trans. Inf. Theory, 64 (2018), pp. 5691–5703.
 - [24] M. DEL PINO AND J. DOLBEAULT, *Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions*, J. Math. Pures Appl. (9), 81 (2002), pp. 847–875.
 - [25] J. DEMANGE, *Des équations à diffusion rapide aux inégalités de Sobolev sur les modèles de la géométrie*, PhD thesis, Université Paul Sabatier Toulouse 3, 2005.
 - [26] J. DOLBEAULT, M. J. ESTEBAN, A. FIGALLI, R. L. FRANK, AND M. LOSS, *Sharp stability for Sobolev and log-Sobolev inequalities, with optimal dimensional dependence*, Preprint arXiv: [2209.08651](https://arxiv.org/abs/2209.08651) and [hal-03780031](https://arxiv.org/abs/hal-03780031), (2023).
 - [27] J. DOLBEAULT, B. NAZARET, AND G. SAVARÉ, *On the Bakry-Emery criterion for linear diffusions and weighted porous media equations*, Commun. Math. Sci., 6 (2008), pp. 477–494.
 - [28] J. DOLBEAULT AND G. TOSCANI, *Improved interpolation inequalities, relative entropy and fast diffusion equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 30 (2013), pp. 917–934.
 - [29] ———, *Stability results for logarithmic Sobolev and Gagliardo–Nirenberg inequalities*, Int. Math. Res. Not. IMRN, 2016 (2016), pp. 473–498.
 - [30] R. EL DAN, J. LEHEC, AND Y. SHENFELD, *Stability of the logarithmic Sobolev inequality via the Föllmer process*, Ann. Inst. Henri Poincaré Probab. Stat., 56 (2020), pp. 2253–2269.
 - [31] A. ESKENAZIS AND Y. SHENFELD, *Intrinsic dimensional functional inequalities on model spaces*, J. Funct. Anal., 286 (2024), p. 110338.
 - [32] L. C. EVANS, *Partial differential equations*, vol. 19 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2010.
 - [33] M. FATHI, E. INDREI, AND M. LEDOUX, *Quantitative logarithmic Sobolev inequalities and stability estimates*, Discrete Contin. Dynam. Systems, 36 (2016), pp. 6835–6853.
 - [34] P. FEDERBUSH, *Partially alternate derivation of a result of Nelson*, J. Mathematical Phys., 10 (1969), pp. 50–52.
 - [35] F. FEO, E. INDREI, M. R. POSTERARO, AND C. ROBERTO, *Some remarks on the stability of the log-Sobolev inequality for the Gaussian measure*, Potential Anal., 47 (2017), pp. 37–52.
 - [36] L. GROSS, *Logarithmic Sobolev inequalities*, Amer. J. Math., 97 (1975), pp. 1061–1083.
 - [37] A. GUIONNET AND B. ZEGARLINSKI, *Lectures on logarithmic Sobolev inequalities*, Séminaire de probabilités de Strasbourg, 36 (2002), pp. 1–134.
 - [38] E. INDREI, *Sharp stability for LSI*, Mathematics, 11 (2023), p. 2670.
 - [39] E. INDREI, *$W^{1,1}$ stability for the LSI*, Preprint arXiv: [2007.03674](https://arxiv.org/abs/2007.03674), (2024).
 - [40] E. INDREI AND D. KIM, *Deficit estimates for the logarithmic Sobolev inequality*, Differential Integral Equations, 34 (2021), pp. 437–466.
 - [41] E. INDREI AND D. MARCON, *A quantitative log-Sobolev inequality for a two parameter family of functions*, Int. Math. Res. Not. IMRN, 2014 (2014), pp. 5563–5580.
 - [42] D. KIM, *Instability results for the logarithmic Sobolev inequality and its application to related inequalities*, Discrete Contin. Dyn. Syst., 42 (2022), pp. 4297–4320.
 - [43] M. LEDOUX, I. NOURDIN, AND G. PECCATI, *Stein’s method, logarithmic Sobolev and transport inequalities*, Geom. Funct. Anal., 25 (2015), pp. 256–306.
 - [44] ———, *A Stein deficit for the logarithmic Sobolev inequality*, Sci. China Math., 60 (2017), pp. 1163–1180.

- [45] K.-A. LEE AND J. L. VÁZQUEZ, *Geometrical properties of solutions of the porous medium equation for large times*, Indiana Univ. Math. J., (2003), pp. 991–1016.
- [46] F. OTTO AND C. VILLANI, *Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality*, J. Funct. Anal., 173 (2000), pp. 361–400.
- [47] G. ROYER, *An initiation to logarithmic Sobolev inequalities*, vol. 14 of SMF/AMS Texts and Monographs, American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2007. Translated from the 1999 French original by Donald Babbitt.
- [48] A. SAUMARD AND J. A. WELLNER, *Log-concavity and strong log-concavity: a review*, Stat. Surv., 8 (2014), p. 45.
- [49] A. J. STAM, *Some inequalities satisfied by the quantities of information of Fisher and Shannon*, Information and Control, 2 (1959), pp. 101–112.
- [50] G. TOSCANI, *An information-theoretic proof of Nash's inequality*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., 24 (2013), pp. 83–93.
- [51] ———, *A concavity property for the reciprocal of Fisher information and its consequences on Costa's EPI*, Phys. A, 432 (2015), pp. 35–42.
- [52] C. VILLANI, *A short proof of the “concavity of entropy power”*, IEEE Trans. Inf. Theory, 46 (2000), pp. 1695–1696.
- [53] ———, *Entropy Production and Convergence to Equilibrium*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2008, pp. 1–70.
- [54] F.-Y. WANG AND J. WANG, *Functional inequalities for convolution probability measures*, Ann. Inst. Henri Poincaré, Probab. Stat., 52 (2016), pp. 898–914.
- [55] F. B. WEISSELER, *Logarithmic Sobolev inequalities for the heat-diffusion semigroup*, Trans. Amer. Math. Soc., 237 (1978), pp. 255–269.
- [56] D. ZIMMERMANN, *Logarithmic Sobolev inequalities for mollified compactly supported measures*, J. Funct. Anal., 265 (2013), pp. 1064–1083.
- [57] ———, *Elementary proof of logarithmic Sobolev inequalities for Gaussian convolutions on \mathbb{R}* , Ann. Math. Blaise Pascal, 23 (2016), pp. 129–140.