

Symmetry and non-uniformly elliptic operators

Jean Dolbeault *

Ceremade, UMR no. 7534 CNRS, Université Paris IX-Dauphine,

Place de Lattre de Tassigny, 75775 Paris Cédex 16, France

E-mail: dolbeault@ceremade.dauphine.fr

Internet: <http://www.ceremade.dauphine.fr/~dolbeault>

Patricio Felmer *, †

Departamento de Ingeniería Matemática

and Centro de Modelamiento Matemático, UMR no. 2071 CNRS-UCHile

Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile

E-mail: pfelmer@dim.uchile.cl

Régis Monneau *

CERMICS, Ecole Nationale des Ponts et Chaussées,

6 et 8 avenue Blaise Pascal,

Cité Descartes, Champs-sur-Marne, 77455 Marne-La-Vallée, France

E-mail: monneau@cermics.enpc.fr

Internet: <http://cermics.enpc.fr/~monneau/home.html>

Abstract. The goal of this paper is to study the symmetry properties of nonnegative solutions of elliptic equations involving a non uniformly elliptic operator. We consider on a ball the solutions of

$$\Delta_p u + f(u) = 0$$

with zero Dirichlet boundary conditions, for $p > 2$, where Δ_p is the p -Laplace operator and f a continuous nonlinearity. The main tools are a comparison result for weak solutions and a local moving plane method which has been previously used in the $p = 2$ case. We prove local and global symmetry results when u is of class $C^{1,\gamma}$ for γ large enough, under some additional technical assumptions.

Keywords. Elliptic equations, non uniformly elliptic operators, p -Laplace operator, scalar field equations, monotonicity, symmetry, local symmetry, positivity, non Lipschitz nonlinearities, comparison techniques, weak solutions, maximum principle, Hopf's lemma, local moving plane method, $C^{1,\alpha}$ regularity

AMS classification (2000). 35B50, 35B65, 35B05

*Partially supported by ECOS-Conicyt.

†Partially supported by Fondecyt Grant # 1030929 and FONDAP de Matemáticas Aplicadas.

1 Introduction and main results

The goal of this paper is to prove local and global symmetry results for the nonnegative solutions of

$$\begin{cases} \Delta_p u + f(u) = 0 & \text{in } B \\ u \geq 0 & \text{in } B \\ u = 0 & \text{on } \partial B \end{cases} \quad (1)$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the p -Laplace operator or p -Laplacian, and $B = B(0, 1)$ is the unit ball in \mathbb{R}^d , $d \geq 1$. We will focus on the non uniformly elliptic case: $p > 2$, and consider nonlinearities f which are continuous on \mathbb{R}^+ .

In earlier works of two of the authors [14, 15, 16], it has been noted that for $p = 2$ the C^2 regularity of u is apparently a threshold in order to apply the moving plane technique. Here we will see that the threshold for $p > 2$ is rather $C^{1,1/(p-1)}$. As in [16], additional properties of f are required at least close to its zeros. The main tools of this paper are the maximum principle when the nonlinearity is nonincreasing or, based on a local inversion, when $\nabla u \neq 0$, and a local moving plane technique.

Our results are not entirely satisfactory, since the statements are rather technical, but on the other hand, they are a first achievement in obtaining symmetry results by comparison techniques in the case of non uniformly elliptic operators. Let us start with a simple case, where we assume the following additional conditions on f .

- (H1) *Let f be a real function defined on \mathbb{R}^+ such that:*
- a) $f \in C^\alpha(\mathbb{R}^+)$ for some $\alpha \in (0, 1)$,
 - b) $\exists \theta > 0$ such that $f > 0$ on $(0, \theta)$, $f < 0$ on $(\theta, +\infty)$,
 - c) f is nonincreasing on $(\theta - \eta, \theta + \eta)$ for some $\eta > 0$.

Theorem 1 (Symmetry result for the p -Laplacian with $p > 2$)

Let $p > 2$, $\gamma \in (1/(p-1), 1)$. Consider a function f satisfying (H1) and a solution $u \in C^{1,\gamma}(B)$ of (1). Then u is positive and radially symmetric on B .

Remark 1 *A consequence of the $C^{1,\gamma}$ regularity of the solution on the whole ball is that the maximum of this solution needs to be reached at the level $u = \theta$ (see Lemma 8). Does such a solution exist? Given a function f and using a shooting method, we obtain a solution on a ball of radius large enough. Then a rescaling gives a solution on B for an appropriately rescaled nonlinearity f , which provides an example of existence of a solution.*

A result of symmetry similar to that of Theorem 1 also holds under weaker regularity assumptions on f .

- (H2) $f \in C^0(\mathbb{R}^+)$ and $f > 0$ on $(0, +\infty)$.

Theorem 2 (Symmetry result with f only continuous)

Let $p > 2$, $\gamma \in (1/(p-1), 1)$. Assume that (H2) holds and consider a solution $u \in W^{1,p}(B) \cap L^\infty(B)$ of (1) such that, with $M = \|u\|_{L^\infty(B)}$,

$$u \in C^2(B \setminus (\nabla u)^{-1}(0)) \cap C^{1,\gamma}(u^{-1}([0, M])) .$$

If f is nonincreasing in a neighborhood of M , then u is positive and radially symmetric on B .

Remark 2 *If we assume that u belongs to $C^{1,\gamma}(B)$ with $\gamma > 1/(p-1)$, then $f(M) = 0$. See Lemma 8. Reciprocally if $f(M) \neq 0$, as we shall see in Example 1, a solution like the one of Theorem 2 has a regularity which is not better than $C^{1,1/(p-1)}$ at the points x such that $u(x) = M$. This is the reason why the $C^{1,\gamma}$ regularity cannot be assumed at such points.*

The starting point of the moving plane technique is the celebrated paper by Gidas, Ni and Nirenberg [20] based on [1, 2, 26]. An improvement was made in [19, 5, 27] using the monotonicity properties of the nonlinearity. This led to a local moving plane technique [14, 15, 16] in the case of the Laplacian, which is well suited for nonlinearities with low regularity. The corresponding notion of *local symmetry* also seems appropriate in the case of the p -Laplacian. Further interesting results were obtained in [17, 18].

For nonlinear elliptic operators, the story is shorter [8, 9, 10]. It strongly relies on comparison methods [29, 6, 7] but covers the case of the p -Laplace operator only when $p < 2$. For completeness, let us mention that related results have been obtained by rearrangement techniques, for instance in [23, 21], and that in [3, 4], F. Brock has been able to get symmetry results for the p -Laplace operator by a completely different approach based on a continuous Steiner symmetrization.

Our goal is to get local symmetry results for $p > 2$ using an adapted version of the maximum principle [11, 24, 25] and regularity properties [12, 28, 13, 22], with a *local moving plane* method which is essentially adapted from [15, 16]. The paper is organized as follows. In the next section, we state a local symmetry result (Theorem 3). Theorem 2 is a simple corollary of Theorem 3. In the third section we start with two examples, which motivate the choice of the class of regularity of the solutions and then state three useful lemmas. The fourth section is devoted to the proof of Theorem 3 by the *local moving plane* method. Proofs of Theorems 1 and 2 are contained in the proof of Theorem 3.

2 A general result

In this section we present a local symmetry result, generalizing Theorem 2 to sign-changing functions f .

- (H) *The function $f \in C^0(\mathbb{R}^+)$ has a finite number of zeros and $f(0) \geq 0$. Moreover, if $f(u_0) = 0$ for some $u_0 \geq 0$, then there exists $\eta > 0$ such that:*
- (a) *either f is nonincreasing on $(u_0 - \eta, u_0 + \eta) \cap \mathbb{R}^+$,*
 - (b) *or $f(u) \geq 0$ on $(u_0, u_0 + \eta)$,*

We say that $u_0 \in f^{-1}(0)$ belongs to Z_a in case (a) and to Z_b in case (b). Note that $f^{-1}(0) = Z_a \cup Z_b$.

Theorem 3 (Local symmetry result with f only continuous)

Let $p > 2$, $\gamma \in (1/(p-1), 1)$. Assume that f satisfies (H) and consider a $C^1(\overline{B})$ solution u of (1) such that, with $M = \|u\|_{L^\infty(B)}$,

$$u \in C^2(B \setminus (\nabla u)^{-1}(0)) \cap C^{1,\gamma}(u^{-1}([0, M])) .$$

If f is nonincreasing in a neighborhood of M and if the set $J := u^{-1}(Z_b) \cap (\nabla u)^{-1}(0)$ is empty, then u is locally radially symmetric and the set $u^{-1}([0, M]) \cap (\nabla u)^{-1}(0)$ is contained in Z_a .

Here *locally radially symmetric* [4] means that there exists an at most countable family $(u_i)_{i \in I}$ of radial functions with supports in balls B_i , $i \in I$, such that $u_i|_{B_i}$ is a nonnegative radial nonincreasing solution of $\Delta_p u_i + f(u_i + C_i) = 0$ on B_i , where $u_i|_{(B \setminus B_i)} \equiv 0$ and $C_i \geq 0$ is a constant satisfying $f(C_i) = 0$, and such that

$$u = \sum_{i \in I} u_i .$$

Remark 3 *In the case of Theorem 3, I is finite. Monotonicity holds on balls or annuli. As a consequence, in Theorems 1 and 2, monotonicity holds along any radius. The monotonicity is even strict in case of Theorem 2, while there might be a plateau at level θ in case of Theorem 1.*

The regularity of locally radially symmetric solutions can be studied by elementary methods, which, under an additional condition on f in case (a) of (H), allows to give a sufficient condition under which all sufficiently smooth solutions are globally radially symmetric.

Corollary 4 *Let $\alpha \in (0, (p - 2)/2)$ and consider f such that for any $u_0 \in Z_a$,*

$$0 < \liminf_{u \rightarrow u_0} \frac{|f(u)|}{|u - u_0|^\alpha} < +\infty .$$

Under the same assumptions as in Theorem 3, if $\frac{\alpha+1}{p-\alpha-1} < \gamma < 1$, then u is globally radially symmetric and decreasing along any radius.

This corollary is a consequence of Proposition 5, which will be stated in the next section.

3 Preliminaries

Let us start with two examples and a statement on the regularity of locally radially symmetric solutions of (1). Consider a radial solution in a ball centered at 0 given as a solution, with $r = |x|$, of the ordinary differential equation

$$\begin{cases} \frac{1}{r^{d-1}} (r^{d-1} |u'|^{p-2} u')' + f(u) = 0 , \\ u(0) = u_0 > 0 \quad \text{and} \quad u'(0) = 0 . \end{cases} \quad (2)$$

Example 1 *Let $f(u_0) \neq 0$ and $f(u) \equiv f(u_0)$ on a neighborhood of u_0 . Then $u(r) = u_0 - A \operatorname{sgn}(f(u_0)) r^{\beta+1}$ is the solution of (2) in a neighborhood of $r = 0$, with $\beta = 1/(p - 1)$ and $A = (|f(u_0)|/d)^\beta / (\beta + 1)$. The same formula for u is true up to lower order terms if f is smooth but not constant. In any case, u is exactly in $C^{1,1/(p-1)}$ at $r = 0$, and this is why the level $u = M$ has to be excluded in our results if $f(M) \neq 0$.*

Example 2 *Let $f(u) = |u_0 - u|^{\alpha-1}(u_0 - u)$. Then exactly as above, $u(r) = u_0 \pm A r^{\beta+1}$ is a solution of (2) in a neighborhood of $r = 0$, with $\beta = (\alpha + 1)/(p - \alpha - 1)$ and for some $A > 0$, which depends on d , α and p . In this case, u is exactly in $C^{1,\beta}$ at $r = 0$ with $\beta > 1/(p - 1)$ if $\alpha > 0$. However, there is no uniqueness since $u \equiv u_0$ is also a solution. Actually we may find a continuum of solutions as follows: take $u \equiv u_0$ on $(0, r_0)$ for some $r_0 > 0$, and $u(r) = u_0 \pm A (r - r_0)^{\beta+1}(1 + o(r - r_0))$, for r in a neighborhood of r_0^+ . This solution also has exactly a $C^{1,\beta}$ regularity.*

Remark 4 *If $u_0 \in Z_b$, then there is no radial solution with $u(0) > u_0$ and $\nabla u = 0$ on $\{u = u_0\}$, because of Lemma 9 (strong maximum principle and Hopf's lemma, see below).*

These two examples unveil regularity, not only for radially symmetric solutions of (1) but also for locally radially symmetric solutions, since (1) is invariant under translation. This can be stated as follows (the proof is left to the reader).

Proposition 5 *Assume that f is continuous. Any locally radially symmetric solution of (1) is of class C^2 outside of its critical set. At a critical point $x_0 \in B$, u is exactly in $C^{1,1/(p-1)}$ if $u_0 = u(x_0) \notin f^{-1}(0)$. If we further assume that f satisfies the hypotheses of Corollary 4 then u is either locally constant or at most of class $C^{1,\beta}$ with $\beta = (\alpha + 1)/(p - \alpha - 1)$.*

The proof of Corollary 4 is a straightforward consequence of Theorem 3 and Proposition 5. The next result relates the assumption $u \geq 0$ with the fact that, in the definition of the local radial symmetry, $u_i \geq 0$ on B_i .

Corollary 6 *Assume that f is continuous and let u be a solution of*

$$\begin{cases} \Delta_p u + f(u) = 0 & \text{in } B, \\ u = 0 & \text{on } \partial B, \end{cases}$$

having the following property: there exists an at most countable family $(u_i)_{i \in I}$ of radial monotone functions with supports in balls $B_i \subset B$ such that $u = \sum_{i \in I} u_i$, where u_i satisfies $\Delta_p u_i + f(u_i + C_i) = 0$ on B_i , $u_i|_{\partial B_i} \equiv 0$, the constants C_i are all non-negative and $f(C_i) = 0$ for all $i \in I$.

If $d \geq 2$ and $u \geq 0$, then $u_i > 0$ in B_i for any $i \in I$, and u is locally radially symmetric in the sense of the definition given in Section 2. Moreover, for any $d \geq 1$, if there exists an $i \in I$ such that $\inf_{B_i} u_i < 0$, then $\inf_{B_i} u < 0$.

Proof. Under the assumptions of Theorem 3, this is a straightforward consequence of the proof. The general case is an extension to $p > 2$ of a result which has been proved for $p = 2$ in [14], Proposition 1 (also see [16]). \square

Before proving Theorem 3, we shall state three lemmata. Let us start with a weak comparison result due to Montenegro [24] (also see [11, 25] for related results).

Lemma 7 (Weak comparison principle) *Let u and v be solutions in $W^{1,p} \cap L^\infty(\omega)$ of*

$$\Delta_p u + f(u) = 0 \quad \text{and} \quad \Delta_p v + f(v) = 0,$$

respectively, where ω is a bounded open connected set in \mathbb{R}^d with a C^1 by parts boundary. Assume that f is a nonincreasing function on $u(\omega) \cup v(\omega)$. If $u \geq v$ on $\partial\omega$ a.e., then $u \geq v$ in ω a.e.

Lemma 8 (Characterization of the critical set) *Consider a domain ω in \mathbb{R}^d . Let $u \in C^{1,\gamma}(\omega)$ be a solution of*

$$\Delta_p u + f(u) = 0 \quad \text{in } \omega$$

and consider $x_0 \in \bar{\omega}$ such that $\nabla u(x_0) = 0$. If $x_0 \in \partial\omega$, assume moreover that $u \in C^{1,\gamma}(\bar{\omega})$, $\partial\omega$ is Lipschitz and $A(x_0) = \lim_{\epsilon \rightarrow 0} \epsilon^{-d} \text{Vol}(\{x \in \omega : |x - x_0| < \epsilon\}) > 0$. If $\gamma > 1/(p - 1)$, then $f(u(x_0)) = 0$.

Proof. Assume for simplicity that $x_0 = 0$ and $u(x_0) = 0$. Consider then $u^\epsilon(x) = \epsilon^{-1}u(\epsilon x)$. A straightforward computation gives

$$\epsilon^{-1}\Delta_p u^\epsilon + f(\epsilon u^\epsilon) = 0.$$

Because of the $C^{1,\gamma}$ regularity of u ,

$$|\nabla u^\epsilon(x)|^{p-1} \leq C|x|^{\gamma(p-1)}\epsilon^{\gamma(p-1)}$$

for some nonnegative constant C . Let ϕ be a nonnegative nonzero radial test function, then

$$\begin{aligned} f(0) A(x_0) \int_{\mathbb{R}^d} \phi \, dx &= \lim_{\epsilon \rightarrow 0} \int_{\epsilon^{-1}\omega} f(\epsilon u^\epsilon) \phi \, dx \\ &= \lim_{\epsilon \rightarrow 0} \epsilon^{-1} \left(\int_{\epsilon^{-1}\omega} |\nabla u^\epsilon|^{p-2} \nabla u^\epsilon \cdot \nabla \phi \, dx \right) \\ &\leq C \lim_{\epsilon \rightarrow 0} \epsilon^{\gamma(p-1)-1} \int_{\mathbb{R}^d} |x|^{\gamma(p-1)} |\nabla \phi| \, dx = 0, \end{aligned}$$

which proves that $f(0) = 0$. Here we assume $A(x_0) = |S^{d-1}|/d$ if $x_0 \in \omega$. \square

For the completeness of this paper, we finally recall a result due to Vázquez [29].

Lemma 9 (Strong maximum principle and Hopf's lemma) *Let ω be a domain in \mathbb{R}^d , $d \geq 1$, and let $u \in C^1(\omega)$ be such that $\Delta_p u \in L^2_{\text{loc}}(\omega)$, $u \geq 0$ a.e. in ω . Assume that*

$$-\Delta_p u + \beta(u) \geq 0 \quad \text{a.e. in } \omega$$

with $\beta : [0, +\infty) \rightarrow \mathbb{R}$ continuous, nondecreasing, $\beta(0) = 0$ and such that either there exists an $\epsilon > 0$ for which $\beta \equiv 0$ on $(0, \epsilon)$ or $\beta(s) > 0$ for any $s > 0$ and $\int_0^1 (s\beta(s))^{-1/p} ds = +\infty$. Then if u does not vanish identically on ω , it is positive everywhere in ω . Moreover, if $u \in C^1(\omega \cup \{x_0\})$ for an $x_0 \in \partial\omega$ that satisfies an interior sphere condition and $u(x_0) = 0$, then the derivative along the unit outgoing normal vector at x_0 satisfies: $\frac{\partial u}{\partial \nu}(x_0) > 0$.

4 Proof of Theorem 3

We will split the proof of Theorem 3 in five steps, which are more or less classical in moving plane techniques. In the first one, we prove that one can start the method. Then we characterize the obstructions and get the global symmetry result when there is no obstruction. In the third and fourth steps, we prove that obstructions are due only to balls on which the solution is radially symmetric. Finally in the fifth step we establish the local symmetry result. We also indicate how the special cases of Theorems 1 and 2 can be treated.

Consider $e_1 \in S^{d-1}$ and denote by x_1 the coordinate along the direction given by e_1 , and by x' the coordinate in the orthogonal hyperplane identified with \mathbb{R}^{d-1} . Let $\lambda > 0$ and

$$\begin{aligned} T_\lambda &= \{x \in B : x = (x_1, x') \in \mathbb{R} \times \mathbb{R}^{d-1}, x_1 = \lambda\}, \\ \Sigma_\lambda &= \{x \in B : x = (x_1, x') \in \mathbb{R} \times \mathbb{R}^{d-1}, x_1 > \lambda\}, \\ x_\lambda &= (2\lambda - x_1, x') \quad \text{if } x = (x_1, x'), \quad u_\lambda(x) = u(x_\lambda) \quad \text{and} \quad w_\lambda = u_\lambda - u. \end{aligned}$$

Step 1: Starting the moving plane method

Consider $\lambda_0 = \sup\{x_1 \in (0, 1] : \exists x' \text{ such that } (x_1, x') \in \overline{B} \text{ and } u(x_1, x') > 0\}$. Up to a change of coordinates, we may therefore assume that $\lambda_0 > 0$. We claim that the moving plane method can be started, that is, there is a $\overline{\lambda} < \lambda_0$ such that $w_\lambda \geq 0$ in Σ_λ for any $\lambda \in (\overline{\lambda}, \lambda_0)$.

Let us assume first that $0 \in Z_b$ or $f(0) > 0$. In this case we notice that for some $\eta > 0$,

$$-\Delta_p u \geq 0 \quad \text{on } \{x \in B : 0 < u(x) < \eta\} .$$

We can use Lemma 9 to conclude that $u > 0$ in B , and $\frac{\partial u}{\partial x_1}(1, 0) < 0$. Thus $\lambda_0 = 1$ and the claim is proved.

Next we assume that $0 \in Z_a$, that is, f is nonincreasing on $(0, \eta)$. Assume by contradiction that there exists a sequence $\lambda_n \nearrow 1$, such that for each $n \in \mathbb{N}$, there exists $x_n \in \Sigma_{\lambda_n}$ such that

$$w_{\lambda_n}(x_n) < 0 .$$

For n large, we can certainly have that $\eta > \sup_{x \in \Sigma_{\lambda_n}} u(x)$. Noticing that on $\partial \Sigma_{\lambda_n}$ we have $u_{\lambda_n} \geq u$, we may apply Lemma 7 with $\omega = \Sigma_{\lambda_n}$ and get a contradiction that proves the claim.

Step 2: Characterizing the obstructions

Let $\bar{\lambda} = \inf\{\lambda > 0 : w_\lambda(x) \geq 0 \quad \forall x \in \Sigma_\lambda\}$. We assume that $\bar{\lambda} > 0$: there exists a sequence $\lambda_n \nearrow \bar{\lambda}$ and a sequence $x_n \in \Sigma_{\lambda_n}$ such that $w_{\lambda_n}(x_n) < 0$. We may assume that x_n is a minimum point of w_{λ_n} , so that $\nabla w_{\lambda_n}(x_n) = 0$. Up to the extraction of a subsequence, x_n converges as $n \rightarrow +\infty$ to some $\bar{x} \in \bar{\Sigma}_{\bar{\lambda}}$, which satisfies $w_{\bar{\lambda}}(\bar{x}) = 0$ and $\nabla w_{\bar{\lambda}}(\bar{x}) = 0$. We call this point \bar{x} an *obstruction*.

We observe that in case $0 \in Z_b$ or $f(0) > 0$, u is decreasing in the direction x_1 near $\partial B \cap \bar{\Sigma}_\lambda$ for λ close to $\bar{\lambda}$ and so that $\bar{x} \notin \partial B$. Then one of the following cases holds:

- 1) $\bar{x} \in \Sigma_{\bar{\lambda}}$. At this point we may have $\nabla u(\bar{x}) = 0$, which implies that $f(u(\bar{x})) = 0$ by Lemma 8 if $u(\bar{x}) < M$, or $w_{\bar{\lambda}} \equiv 0$ on $B(\bar{x}, \epsilon)$ for some $\epsilon > 0$. In fact, if $\nabla u(\bar{x}) \neq 0$, we can use a local inversion method and the strong maximum principle as in [15] to conclude that $w_{\bar{\lambda}} \equiv 0$ near \bar{x} , since $w_{\bar{\lambda}}$ has a local minimum at \bar{x} .
- 2) $\bar{x} \in T_{\bar{\lambda}} \cap B$. Since $\bar{x} \in B$ it is standard that $\frac{\partial u}{\partial x_1}(\bar{x}) = -\frac{1}{2} \frac{\partial w_{\bar{\lambda}}}{\partial x_1}(\bar{x}) = 0$, which implies that $\nabla w_{\bar{\lambda}}(\bar{x}) = \nabla u(\bar{x}) = 0$. Actually, if $\nabla u(\bar{x}) \neq 0$, we can use a local inversion method as in [15] and get a contradiction with Hopf's lemma for usual elliptic operators. Now we have two possibilities: either $u(\bar{x}) = M$, or we can use Lemma 8 to find that $f(u(\bar{x})) = 0$. In summary, either $u(\bar{x}) = M$ or $f(u(\bar{x})) = 0$, and $\nabla u(\bar{x}) = 0$.
- 3) In case $0 \in Z_a$ it may occur that $\bar{x} \in \partial B \cap \bar{\Sigma}_{\bar{\lambda}}$. If $\bar{x} \in \partial B \cap \bar{\Sigma}_{\bar{\lambda}}$ then $u_{\bar{\lambda}}(\bar{x}) = u(\bar{x}) = 0$ and so $\nabla u_{\bar{\lambda}}(\bar{x}) = 0$, since $\bar{x}_{\bar{\lambda}} \in B$. Consequently, if $\nabla u(\bar{x}) \neq 0$ we have that $\nabla w_{\bar{\lambda}}(\bar{x}) \cdot e_1 < 0$, which is a contradiction. Thus we must have $\nabla u(\bar{x}) = 0$. If $\bar{x} \in \bar{T}_{\bar{\lambda}} \cap B$ then $\frac{\partial w_{\bar{\lambda}}}{\partial x_1}(\bar{x}) = -2\frac{\partial u}{\partial x_1}(\bar{x}) = 0$ and we also have $\nabla u(\bar{x}) = 0$.

Step 3: The obstruction set and the critical points of u

Let $\bar{\lambda}$ be the first value of λ for which an obstruction point \bar{x} appears in applying the moving plane method. Remark that all possible obstruction points \bar{x} are contained in the set

$$K := \{x \in \partial \bar{\Sigma}_{\bar{\lambda}} : \nabla u(x) = 0\} \cup \{x \in \bar{\Sigma}_{\bar{\lambda}} \setminus \bar{T}_{\bar{\lambda}} : u(x) = u_{\bar{\lambda}}(x)\} .$$

We claim that the set $K_1 = \{x \in K : \nabla u(x) \neq 0\}$ is not empty.

Assume by contradiction that $K_1 = \emptyset$: $\nabla u(x) = 0$ for all $x \in K$. Let us denote by $Q_\varepsilon(x)$ the set $\{y \in \mathbb{R}^d : \max_{i=1,2,\dots,d} |y_i - x_i| < \varepsilon\}$, where $\varepsilon > 0$ is taken small. From the covering $\cup_{x \in K} Q_\varepsilon(x)$ of the compact set K if $0 \in Z_b$ or $f(0) > 0$, and of the compact set

$K \cup (\overline{K} \cap \overline{T_{\overline{\lambda}}} \cap \partial B)$ if $0 \in Z_a$, we can extract a finite subcovering corresponding to points $x_j \in K$, $j = 1, 2, \dots, L$ for some number L , and define the open set

$$\omega_\varepsilon := \bigcup_{j=1}^L Q_\varepsilon(x_j) \supset K.$$

If $0 \in Z_b$ or by Lemma 9 if $f(0) > 0$, we can choose ε small enough so that $\omega_\varepsilon \subset B$ and observe that by construction ω_ε has a boundary of class C^1 by parts. Next we define $\Gamma = \partial\omega_\varepsilon$. We can find $\delta_1 > 0$ and $\delta > 0$ such that

$$u_{\overline{\lambda}} \geq u + \delta \quad \text{on} \quad \overline{\Sigma_{\overline{\lambda} + \delta_1}} \cap \Gamma.$$

Next we claim that

$$\frac{\partial u}{\partial x_1} < -\delta \quad \text{on} \quad T_{\overline{\lambda}} \cap \Gamma,$$

for a possibly smaller $\delta > 0$. If the claim was false, then there would be a point $x_0 \in T_{\overline{\lambda}} \cap \Gamma$ with $\frac{\partial u}{\partial x_1}(x_0) = 0$. But then, since $x_0 \in \Gamma$ and $u_{\overline{\lambda}}(x_0) = u(x_0)$, we have $\nabla u(x_0) \neq 0$. Thus, we can use a local inversion method and the strong maximum principle as in [15] to prove that $u_{\overline{\lambda}} \equiv u$ in a neighborhood of x_0 in $\Sigma_{\overline{\lambda}}$ and then $K_1 \neq \emptyset$, which is a contradiction with our assumption.

If $0 \in Z_a$, we take δ_1 smaller than $\text{dist}(\overline{T_{\overline{\lambda}}} \cap \partial B, \partial\omega_\varepsilon)$, and we replace ω_ε by $\omega_\varepsilon \cap B$ and Γ by $\partial\omega_\varepsilon \cap \overline{B}$. Let us consider $\eta > 0$ small. First we analyze the behavior of u and u_λ in $A_1 = (\overline{B} \setminus B(0, 1 - \eta)) \cap \overline{\Sigma_{\overline{\lambda} + \delta_1}}$. For any $x_0 \in \Gamma \cap \partial B \cap \overline{\Sigma_{\overline{\lambda} + \delta_1}}$, by definition of K , $u_{\overline{\lambda}}(x_0) = u_{\overline{\lambda}}(x_0) - u(x_0) \geq \delta_0 > 0$. By continuity, for x in a neighborhood of x_0 and for $\overline{\lambda} - \lambda > 0$ small enough, have $u_\lambda(x_0) - u(x_0) \geq \delta_0/2 > 0$. Taking η smaller if necessary, this implies that

$$u_\lambda \geq u \quad \text{in} \quad \Gamma \cap A_1,$$

for $\overline{\lambda} - \lambda > 0$ small enough. Next we consider $A_2 = \overline{B}(0, 1 - \eta) \cap \Sigma_{\overline{\lambda} + \delta_1}$. Then that by continuity, the property

$$u_{\overline{\lambda}} \geq u + \delta \quad \text{on} \quad \Gamma \cap A_2$$

can be extended to

$$u_\lambda \geq u + \delta/2 \quad \text{on} \quad \Gamma \cap A_2,$$

for $\overline{\lambda} - \lambda > 0$ small enough. The property

$$\frac{\partial u}{\partial x_1} < -\delta \quad \text{on} \quad T_{\overline{\lambda}} \cap \Gamma$$

holds for the same reasons as in the first case.

Thus, in both cases, using continuity again, for possibly smaller δ and δ_1 , we see that

$$\frac{\partial u}{\partial x_1} < -\delta \quad \text{on} \quad \left(\overline{\Sigma_{\overline{\lambda}}} \setminus \Sigma_{\overline{\lambda} + \delta_1} \right) \cap \Gamma$$

and, for $\eta > 0$ small enough, we also have

$$\begin{aligned} u_\lambda &\geq u + \delta/2 \quad \text{on} \quad \overline{\Sigma_{\overline{\lambda} + \delta_1}} \cap \Gamma, \quad \text{for} \quad \lambda \in (\overline{\lambda} - \eta, \overline{\lambda}), \\ \frac{\partial u}{\partial x_1} &\leq -\delta/2 \quad \text{on} \quad \left(\overline{\Sigma_{\overline{\lambda}}} \setminus \Sigma_{\overline{\lambda} + \delta_1} \right) \cap \Gamma, \quad \text{for} \quad \lambda \in (\overline{\lambda} - \eta, \overline{\lambda}). \end{aligned}$$

From the last inequality, we get

$$u_\lambda \geq u \quad \text{on} \quad \left(\overline{\Sigma}_\lambda \setminus \Sigma_{\overline{\lambda} + \delta_1} \right) \cap \Gamma ,$$

from where it follows that

$$u_\lambda \geq u \quad \text{on} \quad \Gamma \cap \Sigma_\lambda .$$

But we also have $u_\lambda = u$ on $T_\lambda \cap \omega_\varepsilon$ and $u_\lambda \geq u$ in $\partial B \cap \Gamma$. As a consequence, for $\omega_\varepsilon^\lambda = \omega_\varepsilon \cap \Sigma_\lambda$, we get

$$u_\lambda \geq u \quad \text{on} \quad \partial \omega_\varepsilon^\lambda .$$

By Lemma 8 we have $f(u(K) \setminus \{M\}) \subset \{0\}$, and then, by the assumptions of Theorem 3, $u(K) \subset Z_a \cup \{M\}$. We recall that in a neighborhood of every point in $Z_a \cup \{M\}$, f is nonincreasing, by hypothesis. Thus, for ε, η small enough, this guarantees that f is nonincreasing on $u(\omega_\varepsilon^\lambda) \cup u_\lambda(\omega_\varepsilon^\lambda)$. Then Lemma 7 applies and gives that

$$u_\lambda \geq u \quad \text{on} \quad \omega_\varepsilon^\lambda ,$$

for $\lambda \in (\overline{\lambda} - \eta, \overline{\lambda})$. This proves that there is no obstruction in $\omega_\varepsilon^\lambda$, a contradiction: $K_1 \neq \emptyset$.

Step 4: Getting a local symmetry result

Let us notice that $K_1 \subset \Sigma_{\overline{\lambda}}$. Because of the local inversion method and the strong maximum principle as in [15] again, K_1 is an open set. Let C^+ be one of its connected components:

$$\nabla u = 0 \quad \text{on} \quad \Gamma^+ = \partial C^+ \cap (\overline{\Sigma}_{\overline{\lambda}} \setminus T_{\overline{\lambda}}) .$$

By Lemma 8 and the assumptions of Theorem 3, $u(\Gamma^+) \subset Z_a \cup \{M\}$ and u is equal to a constant on each component of $\partial C^+ \cap \Sigma_{\overline{\lambda}}$. This in particular implies that $\overline{C^+} \cap T_{\overline{\lambda}} \neq \emptyset$, otherwise we would have $\overline{C^+} \subset \overline{\Sigma}_{\overline{\lambda}} \setminus T_{\overline{\lambda}}$: the fact that u is constant in each connected component of ∂C^+ would contradict the fact that u is nondecreasing and $\nabla u \neq 0$ in C^+ .

Define

$$C = \overline{C^+ \cup C^-} ,$$

where C^- is the reflexion of C^+ with respect to $T_{\overline{\lambda}}$. The set C is connected and symmetric with respect to $T_{\overline{\lambda}}$, and by construction the function u is symmetric on C with respect to $T_{\overline{\lambda}}$.

Step 5: Getting a global symmetry result

Let us define $u_1 = \inf_C u$ and $u_2 = \sup_C u$. By monotonicity of u in the x_1 direction, we know that u takes either the value $u_1 > 0$ on the boundary of C , where $\nabla u = 0$, or $u_1 = 0$ on $\partial C \cap \partial B$. Consequently, by Lemma 8, we have $f(u_1) = 0$ if $u_1 \neq 0$. Similarly, either u takes the value u_2 on another connected component of the boundary of C and then $f(u_2) = 0$, or u reaches its maximum in an interior point $\overline{x} \in C$ that we may take in $T_{\overline{\lambda}} \cap C$.

Because u_1 and u_2 belong to $\{0\} \cup Z_a \cup \{M\}$, which is a finite set, and $|\nabla u|$ is uniformly bounded in B , we deduce that C contains a ball of some radius $r_0 > 0$, which depends only on f and u . There are therefore only a finite number of such components C . Moreover, if we consider all possible sets C , symmetric with respect to a certain hyperplane, along which u is increasing and where $\nabla u = 0$ on ∂C (except maybe for $u_1 = 0$), we still have a finite number, say \mathcal{N}_0 .

Now we consider $\mathcal{N} = d\mathcal{N}_0 + 1$ directions $\gamma_i \in S^{d-1}$, $i = 1, 2, \dots, \mathcal{N}$, such that the angle (γ_i, γ_j) is 2π -irrational for any (i, j) with $i \neq j$, and such that any family of d such unit vectors generates \mathbb{R}^d . It is not hard to prove that a set which is symmetric with respect to those directions is actually radially symmetric, with respect to some point in it. See [16] for more details. Thus, if we apply the moving plane procedure in all the directions given above, we find a connected component C which is radially symmetric, i.e. an annulus (or a ball) $C = B_{r_1}(x_0) \setminus B_{r_2}(x_0)$ with $r_1 > r_2 \geq 0$, and such that $u = u_1$ on $\partial B_{r_1}(x_0)$ and $u = u_2$ on $\partial B_{r_2}(x_0)$. Then we can consider the following modifications of the function u . First we consider v_1 defined in B as $v_1 \equiv u_1$ in $B_{r_1}(x_0)$ and $v_1 = u$ in $B \setminus B_{r_1}(x_0)$. Second, in case $r_2 > 0$, we define $v_2 = u$ in $B_{r_2}(x_0)$.

At this point, under the hypothesis of Theorem 1, the function u is radially symmetric, and $M = \theta$. We observe that the set $\{x : u(x) = M\}$ is a closed ball of radius $r \geq 0$.

Under the hypothesis of Theorem 2, we also see that the function u is radially symmetric.

In the case of Theorem 3, we reapply Step 1-5 to the function $x \mapsto v_1(x)$ in B , and to the function $x \mapsto v_2(x) - u_2$ in $B_{r_2}(x_0)$. We remark that $u \geq u_2$ on $B_{r_2}(x_0)$ because of the monotonicity of the solution in x_1 . Thus all assumptions of Theorem 3 hold with B replaced by $B_{r_2}(x_0)$. A finite iteration provides the result. \square

References

- [1] A.D. ALEXANDROFF, *Uniqueness theorem for surfaces in the large*, Vestnik Leningrad Univ. Math. **11**, 5-17, (1956).
- [2] A.D. ALEXANDROFF, *A characteristic property of the spheres*, Ann. Mat. Pura Appl. **58**, 303-354, (1962).
- [3] F. BROCK, *Continuous Steiner symmetrization*, Math. Nach. **172**, 25-48, (1995).
- [4] F. BROCK, *Continuous rearrangements and symmetry of solutions of elliptic problems*, Proc. Indian Acad. Sci. Math. Sci. **110**, 157-204, (2000).
- [5] C. CORTÁZAR, M. ELGUETA, P. FELMER, *Symmetry in an elliptic problem and the blow-up set of a quasilinear heat equation*, Comm. Partial Differential Equations **21** no. 3-4, 507-520, (1996).
- [6] L. DAMASCELLI, *Some remarks on the method of moving planes*, Differential Integral Equations **11**, 493-501, (1998).
- [7] L. DAMASCELLI, *Comparison theorems for some quasilinear degenerate elliptic operators and applications to symmetry and monotonicity results*, Ann. I.H.P. Anal. Non Linéaire **15**, 493-516, (1998).
- [8] L. DAMASCELLI, F. PACELLA, *Monotonicity and symmetry of solutions of p -Laplace equation, $1 < p < 2$, via the moving plane method*, Ann. Sc. Norm. Sup. Pisa Cl. Sci. **26**, 689-707, (1998).
- [9] L. DAMASCELLI, F. PACELLA, M. RAMASWAMY, *Symmetry of ground states of p -Laplace equations via the moving plane method*, Arch. Ration. Mech. Anal. **148**, 291-308, (1999).

- [10] L. DAMASCELLI, M. RAMASWAMY, *Symmetry of C^1 solutions of p -Laplace equations in \mathbb{R}^N* , Adv. Nonlinear Stud. **1** no. 1, 40-64, (2001).
- [11] J.I. DIAZ, J.E. SAA, U. THIEL, *On the equation of prescribed mean curvature and other quasilinear elliptic equations with locally vanishing solutions*, Rev. Unión Mat. Argent. **35**, 175-206, (1989).
- [12] E. DI BENEDETTO, *$C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations*, Nonlinear Analysis **7**, 827-850, (1983).
- [13] E. DIBENEDETTO, *Degenerate parabolic equations*, Universitext, Springer-Verlag, New York, 1993.
- [14] J. DOLBEAULT, P. FELMER, *Symétrie des solutions d'équations semi-linéaires elliptiques [Symmetry of the solutions of semilinear elliptic equations]*, C. R. Acad. Sci. Paris, Série I **329**, 677-682, (1999).
- [15] J. DOLBEAULT, P. FELMER, *Symmetry and monotonicity properties for positive solutions of semi-linear elliptic PDE's*, Comm. Partial Differential Equations **25**, 1153-1169, (2000)
- [16] J. DOLBEAULT, P. FELMER, *Monotonicity and symmetry of positive solutions to non-linear elliptic equations : local moving planes and unique continuation*, Preprint Cere-made no. 9950.
- [17] A. FARINA, *Propriétés de monotonie et de symétrie unidimensionnelle pour les solutions de $\Delta u + f(u) = 0$ avec des fonctions f éventuellement discontinues*, C. R. Acad. Sci. Paris, Sér. I, **330**, 973-978, (2000).
- [18] A. FARINA, *Monotonicity and one-dimensional symmetry for the solutions of $\Delta u + f(u) = 0$ in \mathbb{R}^N with possibly discontinuous nonlinearity*, Adv. Math. Sci. Appl. **11** no. 2, 811-834, (2001).
- [19] B. FRANCHI, E. LANCONELLI, J. SERRIN, *Existence and uniqueness of nonnegative solutions of quasilinear equations in \mathbb{R}^N* , Adv. Math. **118**, 177-243, (1996).
- [20] B. GIDAS, W. M. NI, L. NIRENBERG, *Symmetry and related properties via the maximum principle*, Commun. Math. Phys. **68**, 209-243, (1979).
- [21] M. GROSSI, S. KESAVAN, F. PACELLA, M. RAMASWAMY, *Symmetry of positive solutions of some nonlinear equations*, Topol. Methods Nonlinear Anal. **12** no. 1, 47-59, (1998).
- [22] J. HEINONEN, T. KILPELÄINEN, O. MARTIO, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford New York, Tokyo, Clarendon Press, 1993.
- [23] P.-L. LIONS, *Two geometrical properties of solutions of semilinear problems*, Applicable Anal. **12** no. 4, 267-272, (1981).
- [24] M. MONTENEGRO, *Strong maximum principles for supersolutions of quasilinear elliptic equations*, Nonlinear Anal. **37**, 431-448, (1999).

- [25] P. PUCCI, J. SERRIN, H. ZOU, *A strong maximum principle and a compact support principle for singular elliptic inequalities*, J. Math. Pures Appl. (9), **78** no. 8, 769-789, (1999).
- [26] J. SERRIN, *A symmetry problem in potential theory*, Arch. Ration. Mech. Anal. **43**, 304-318, (1971).
- [27] J. SERRIN, H. ZOU, *Symmetry of ground states of quasilinear elliptic equations*, Arch. Ration. Mech. Anal. **148**, 265-290, (1999).
- [28] P. TOLKSDORF, *Everywhere-regularity for some quasilinear systems with lack of ellipticity*, J. Differential Equations **51**, 126-150, (1984).
- [29] J.L. VÁZQUEZ, *A strong maximum principle for some quasilinear elliptic equations*, Appl. Math. Optim. **12**, 191-202, (1984).