

INVARIANT SUBMANIFOLDS OF CONFORMAL SYMPLECTIC DYNAMICS

MARIE-CLAUDE ARNAUD^{†,‡,◦} & JACQUES FEJOZ^{*,**,◦}

ABSTRACT. We study invariant manifolds of conformal symplectic dynamical systems on a symplectic manifold (\mathcal{M}, ω) of dimension ≥ 4 . This class of systems is the 1-dimensional extension of symplectic dynamical systems for which the symplectic form is transformed colinearly to itself.

In this context, we first examine how the ω -isotropy of an invariant manifold \mathcal{N} relates to the entropy of the dynamics it carries. Central to our study is Yomdin's inequality, and a refinement obtained using that the local entropies have no effect transversally to the characteristic foliation of \mathcal{N} .

When (\mathcal{M}, ω) is exact and \mathcal{N} is isotropic, we also show that \mathcal{N} must be exact for some choice of the primitive of ω , under the condition that the dynamics acts trivially on the cohomology of degree 1 of \mathcal{N} . The conclusion partially extends to the case when a one-sided orbit of \mathcal{N} has compact closure.

We eventually prove the uniqueness of invariant submanifolds \mathcal{N} when \mathcal{M} is a cotangent bundle, provided that the dynamics is isotopic to the identity among Hamiltonian diffeomorphisms. In the case of the cotangent bundle of the torus, a theorem of Shelukhin allows us to conclude that \mathcal{N} is unique even among submanifolds with relatively compact orbits.

2020 *Mathematics Subject Classification.* 37C05, 37J39, 38A35 .

Key words and phrases. conformal symplectic dynamics, isotropy, entropy, exactness, Lagrangian submanifold, invariant manifold.

[†] Université Paris Cité and Sorbonne Université, CNRS, IMJ-PRG, F-75006 Paris, France.

[‡] Member of the *Institut universitaire de France*.

[◦] ANR AAPG 2021 PRC CoSyDy: Conformally symplectic dynamics, beyond symplectic dynamics, ANR-CE40-0014.

^{*} Université Paris Dauphine – PSL, Ceremade.

^{**} Observatoire de Paris – PSL, IMCCE.

CONTENTS

1. Introduction	2
2. Isotropy	6
3. Entropy	10
4. Liouville class of invariant submanifolds	14
5. Liouville class of Lagrangian submanifolds of $T^*\mathcal{Q}$ with compact orbits	18
6. Uniqueness	21
7. Examples	25
Appendix A. Yomdin's inequality	27
Appendix B. Conformal Dynamics are exact	28
References	30

1. INTRODUCTION

Let $(\mathcal{M}^{2d}, \omega)$ be a symplectic manifold. Symplectic dynamical systems form a class of infinite codimension. We will study conformal symplectic dynamics, a now classical extension of symplectic dynamics¹ where the symplectic form may change in its own direction:

Definition 1.

- A diffeomorphism $f : \mathcal{M} \hookrightarrow \mathcal{M}$ is *conformal symplectic* if $f^*\omega = a\omega$ for some $a > 0$ (conformality ratio).²
- A complete vector field X on \mathcal{M} is *conformal symplectic* if $L_X\omega = \alpha\omega$, where L_X is the Lie derivative, for some $\alpha \in \mathbb{R}$ (conformality rate).³

Such dynamics encapsulate mechanical systems whose friction force is proportional to velocity, in which case $a < 1$ or $\alpha < 0$.

In this paper we will focus on the non-symplectic case, i.e. $a \neq 1$ and $\alpha \neq 0$. Of course, time reversal changes a in $1/a$ and α in $-\alpha$.

For such a dynamics, the volume form $\omega^{\wedge d}$ is monotonic. So if such a dynamics exists on \mathcal{M} , \mathcal{M} cannot be closed and has infinite volume. Moreover, when the dynamics is given by a vector field X , the symplectic form satisfies $\omega = \frac{1}{\alpha}L_X\omega = d\left(\frac{1}{\alpha}i_X\omega\right)$ and is exact. Hence conformal vector fields exist only on exact symplectic manifolds. Yet this is not the case for conformal diffeomorphisms (see an example in Proposition 2).

¹Vaisman [20] and others have defined local conformal symplectic structures on a manifold \mathcal{M} . There is a corresponding notion of dynamics preserving the structure, thus extending our setting.

²As Libermann noticed [10]: if $f^*\omega = a\omega$ for some smooth function a , $a\omega$ being closed we have $da \wedge \omega = 0$, which implies, if \mathcal{M} has dimension ≥ 4 , that a is constant.

³Then the flow (φ_t) of X is conformal symplectic and $\varphi_t^*\omega = e^{\alpha t}\omega$.

Also, if a vector field X is conformal symplectic of conformality rate α and if Z is the Liouville vector field associated with the 1-form $\lambda = -\frac{1}{\alpha}i_X\omega$ i.e., $i_Z\omega = \lambda$, then $X + \alpha Z$ is symplectic. Thus, when ω is exact, conformal symplectic vector fields form a 1-dimensional extension of the space of symplectic vector fields.

When (\mathcal{M}, ω) is exact, there exists a 1-parameter subgroup \mathcal{C} of the set of conformal symplectic diffeomorphisms such that the group of conformal symplectic diffeomorphisms is $\{f \circ g; (f, g) \in \mathcal{C} \times \mathcal{S}\}$ where \mathcal{S} is the set of symplectic diffeomorphisms. When \mathcal{M} is not exact, let \mathcal{R} be the subgroup of \mathbb{R}_+^* of conformality ratios of conformal symplectic diffeomorphisms of \mathcal{M} . This subgroup can be trivial, e.g. when \mathcal{M} is compact (all conformal symplectic diffeomorphisms are symplectic).

Questions. Can \mathcal{R} be strictly between $\{1\}$ and \mathbb{R}_+^* ? Assuming that $\mathcal{R} = \mathbb{R}_+^*$, does there exist a continuous 1-parameter family of conformal symplectic diffeomorphisms indexed by its conformality ratio in \mathbb{R}_+^* ?

An important case is that of cotangent bundles $(\mathcal{M} = T^*\mathcal{Q}, \omega = -d\lambda)$, where \mathcal{Q} is a manifold and λ is the canonical Liouville 1-form. A continuous-time example is the flow $\exp(tZ_\lambda)(q, p) = (q, e^{-t}p)$ of the Liouville vector field Z_λ defined by $i_{Z_\lambda}(-d\lambda) = \lambda$ and a discrete-time example is $f = \exp Z_\lambda : (q, p) \mapsto (q, ap)$, $a = e^{-1}$. These two examples of conformal symplectic dynamics have a very simple behaviour:

- there is a global attractor \mathcal{A} ;
- the ω -limit set of every orbit is a point of \mathcal{A} .

More generally, consider a discounted Tonelli vector field X on $T^*\mathcal{Q}$ of negative rate α ; by definition it satisfies $i_X\omega = dH + \alpha\lambda$ for some Hamiltonian H which is superlinear in the fiber direction and whose Hessian in the fiber direction is positive definite. It has been shown that the flow of such a vector field has a global attractor [12].

In the general setting, many natural questions are open, for example:

Questions. Which conditions ensure the existence of a global attractor? And provided that the global attractor exists (necessarily having zero volume), what can be said of its size?

As a first step, in this article we focus on the case of invariant submanifolds (with a digression on the case of submanifolds with compact orbit), although the study of dissipative twist maps proves that there can exist invariant subsets that are not submanifolds [9].

First, we explore the isotropy of invariant submanifolds. This question is akin to its analogue in symplectic dynamics, where both negative and

positive results have been proven in particular for invariant tori carrying minimal quasiperiodic flows.

We start by providing an example where an invariant submanifold is a hypersurface and hence non-isotropic (Propositions 1 and 2 in section 2). There exist similar examples due to McDuff, [13] and Geiges [4, 5], but our example is somewhat more explicit. We do not know if there exist examples of invariant non-isotropic submanifolds that are invariant by a conformally symplectic dynamics on a cotangent bundle. An even more difficult question is to determine whether such submanifolds may exist for discounted Tonelli flows on cotangent bundles. In this case and when $\dim \mathcal{M} \geq 4$, the global attractor never separates \mathcal{M} and hence cannot be a hypersurface.

In turn, we show some positive results regarding the isotropy of invariant submanifolds. If the invariant submanifold is a surface, isotropy follows from a simple argument using the growth of the area. In higher dimension, a first result follows from Yomdin's theory [22, 7]. Proposition 4 of section 2 states that if a smooth⁴ conformal diffeomorphism $f : \mathcal{M} \rightarrow \mathcal{M}$ with conformality rate a has an invariant smooth submanifold $\mathcal{N} \subset \mathcal{M}$ such that the topological entropy of $f|_{\mathcal{N}}$ is less than $|\log(a)|$, then \mathcal{N} is isotropic.

But Yomdin's proof can be improved in the setting of diffeomorphisms which are conformal with respect to a presymplectic form. Here, we prove that the so-called local entropies have no effect on the volume growth transversally to the characteristic foliation of \mathcal{N} (section 3). It follows that if a conformal symplectic C^3 -diffeomorphism of conformality ratio a has an invariant C^3 -manifold on which ω has constant rank 2ℓ and such that the entropy of $f|_{\mathcal{N}}$ is smaller than $\ell |\log a|$, \mathcal{N} is isotropic. In particular, if an invariant submanifold carries a minimal dynamics (every orbit is dense) with zero entropy, it is isotropic (corollary 2).

This new result assumes less regularity than the former one (C^3 instead of smooth in Proposition 4) but requires that the symplectic form restricted to the submanifold has constant rank.

A related result is [2, 2.2.1], where the authors prove that if a C^1 conformal dynamics has a C^1 invariant torus on which the dynamics is C^1 conjugate to a rigid rotation, then this torus is isotropic. This result is a direct consequence of Proposition 4. Corollary 2 of section 3 doesn't imply this result because our result requires more regularity, and on the other hand our result applies when a C^3 dynamics is C^0 conjugated to a transitive rotation.

Second, we examine the question of exactness. In this purpose, in section 4 we assume that $(\mathcal{M}, \omega = -d\lambda)$ is exact. Define the Liouville

⁴Smooth means C^∞ .

class of an isotropic embedding in \mathcal{M} as the cohomology class of the form induced by λ . The embedding is called exact when this class vanishes. The action of conformal symplectic diffeomorphisms on Liouville classes depends on a notion of exactness for the diffeomorphisms themselves. Let $f : \mathcal{M} \hookrightarrow \mathcal{M}$ be a conformal symplectic diffeomorphism of conformality ratio a . The form $f^*\lambda - a\lambda$ is closed.

Definition 2. The diffeomorphism f is λ *conformal exact symplectic (CES)* if $f^*\lambda - a\lambda$ is exact.

It is *Hamiltonian* if f is the time-one map of the flow of a non autonomous conformal Hamiltonian vector field X_t (meaning that $i_{X_t}\omega = \alpha_t\lambda + dH_t$ for all t).

These definitions depend of the chosen primitive of the symplectic form. We prove in appendix B that there is always a choice of primitive for which f is exact. Alternatively, we also show that f is symplectically conjugate to a diffeomorphism which is exact with respect to the initial λ (see appendix B). Hence we state our results for exact conformal symplectic dynamics (see section 4 for more comprehensive statements).

Our main result here is that if f is an exact conformal symplectic diffeomorphism and if \mathcal{S} is a strongly f -invariant submanifold (in the sense that $j \circ f(\mathcal{S}) = j(\mathcal{S})$ and f acts trivially on $H^1(j(\mathcal{S}), \mathbb{R})$), j is exact.

When \mathcal{L} is a Lagrangian submanifold that is H-isotopic⁵ to a graph in $\mathcal{M} = T^*\mathcal{Q}$ and f is CS isotopic⁶ to $\text{Id}_{\mathcal{M}}$, we obtain the same conclusion when assuming only that the orbit of \mathcal{L} is bounded. For example, the submanifolds that are H-isotopic to the zero section and contained in an attractor satisfy this hypothesis.

Question. Is it possible to obtain similar results without assuming that the Lagrangian submanifold is H-isotopic to a graph? On other manifolds?

Third, in section 6, we raise the question of the uniqueness of a invariant Lagrangian submanifolds in a cotangent bundle $(T^*\mathcal{Q}, -d\lambda)$. Indeed, let $f : T^*\mathcal{Q} \hookrightarrow T^*\mathcal{Q}$ be a CES diffeomorphism that is CH isotopic⁷ to $\text{Id}_{T^*\mathcal{Q}}$. We show that there exists at most one submanifold of $T^*\mathcal{Q}$ that is H-isotopic to the zero section and invariant by f . Key to the proof is the Viterbo distance of Lagrangian submanifolds which are H-isotopic to the zero section, and the fact that this distance is monotonic with respect to the action of f .

⁵By *H-isotopic*, we mean isotopic among Hamiltonian diffeomorphisms.

⁶By *CS isotopic*, we mean isotopic among conformal symplectic diffeomorphisms.

⁷By *CH-isotopic*, we naturally mean isotopic among conformally Hamiltonian diffeomorphisms

A recent result of Shelukhin even allows us to show the following. Let $f : T^*\mathbb{T}^n \hookrightarrow T^*\mathbb{T}^n$ be a CES diffeomorphism that is CH-isotopic to $\text{Id}_{T^*\mathbb{T}^n}$. Then there exists at most one submanifold \mathcal{L} which is H-isotopic to the zero section and such that

$$\bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L}) \text{ is relatively compact.}$$

Hence when it exists, \mathcal{L} is invariant by f .

For discounted Tonelli flows, it was known that there is at most one invariant exact Lagrangian graph because this corresponds to the unique weak KAM solution [12]. But we give in Section 7 an example of such a dynamics with an invariant H-isotopic to a graph submanifold that is not a graph, hence even in this case our uniqueness result is new.

2. ISOTROPY

The so-called Mañé example [11] (see subsection 7.1) shows that any flow defined on a closed manifold \mathcal{Q} can be achieved as the restriction of a Tonelli conformal Hamiltonian flow to the zero section of $T^*\mathcal{Q}$. In this case, the zero section is an invariant Lagrangian submanifold.

The following example, which is very similar to an example of [5], is key to this section. It shows that a closed submanifold which is invariant by a conformal symplectic dynamics may be non ω -isotropic. In the remaining of the section, we will give some general conditions under which the submanifold must be ω -isotropic.

Proposition 1. *There exists a conformal symplectic vector field X on a 4-dimensional symplectic manifold (\mathcal{M}, ω) , with a 3-dimensional invariant submanifold \mathcal{L} (hence \mathcal{L} is not isotropic).*

Moreover, the submanifold \mathcal{L} is the global attractor for the flow (φ_t) of X , $(\varphi_t|_{\mathcal{L}})$ is conjugate to the suspension of an Anosov automorphism of \mathbb{T}^2 with 2-dimensional stable and unstable foliations, and $(\varphi_t|_{\mathcal{L}})$ is transitive with entropy equal to $|\alpha|$, where α is the conformality rate of X .

- Remarks 3.**
- (1) In our example, \mathcal{L} is coisotropic, but it is easy to extend this example to an invariant submanifold which is neither isotropic nor coisotropic. Indeed, let Y be a conformal symplectic vector field on a symplectic manifold (\mathcal{N}, ω') with a periodic orbit γ . Then the sum $X \oplus Y$ admits $\mathcal{L} \times \gamma$ as an invariant submanifold that is neither isotropic nor coisotropic in $\mathcal{M} \times \mathcal{N}$ if $\dim \mathcal{N} \geq 4$.
 - (2) The submanifold \mathcal{L} is the maximal (among compact subsets) attractor of the dynamics.
 - (3) Replacing the vector field X by bX for $b \in \mathbb{R}$, we can achieve any positive value for the entropy.

Questions. We don't know if it is possible to build a non-isotropic example on a cotangent bundle endowed with its usual symplectic form or, even stronger, if a similar example exists on such a manifold among Tonelli flows.

Proof of Proposition 1. We consider an Anosov automorphism $A : \mathbb{T}^2 \hookrightarrow \mathbb{T}^2$ induced by a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ with eigenvalues $0 < \lambda_- < 1 < \lambda_+ = \frac{1}{\lambda_-}$ and eigenvectors v_{\pm} . An example of such an automorphism is $A(x, y) = (2x + y, x + y)$, with eigenvalues $\lambda_- = \frac{3-\sqrt{5}}{2} < 1$ and $\lambda_+ = \frac{3+\sqrt{5}}{2} > 1$.

Following [1], we define a suspension T of the diffeomorphism by using the following relation on $\mathbb{T}^2 \times \mathbb{R}$ (writing $\xi = (x, y)$):

$$\forall (\xi, z) \in \mathbb{T}^2 \times \mathbb{R}, (\xi, z) \sim F(\xi, z) := (A\xi, z - 1).$$

Denote by α_{\pm} the linear forms on \mathbb{R}^2 such that $\alpha_{\pm}(v_{\pm}) = 1$ and $\alpha_{\pm}(v_{\mp}) = 0$. Observe that $\alpha_{\pm} \circ A = \lambda_{\pm} \alpha_{\pm}$. Rescale the forms α_{\pm} in the z -direction in order to get F -invariant forms on $\mathbb{T}^2 \times \mathbb{R}$: define

$$\beta_{\pm}(\xi, z) = (\lambda_{\pm})^z \alpha_{\pm}(\xi),$$

so that

$$F^* \beta_{\pm} = (\lambda_{\pm})^{z-1} \alpha_{\pm} \circ A = (\lambda_{\pm})^z \alpha_{\pm} = \beta_{\pm}.$$

Hence β_{\pm} is F -invariant and defines a 1-form on the quotient manifold $\mathcal{N} = (\mathbb{T}^2 \times \mathbb{R}) / \sim$. We use the same notation for these 1-forms. Then

$$(1) \quad d\beta_{\pm} = \ln \lambda_{\pm} dz \wedge \beta_{\pm}.$$

We consider the vector field $X = (0, 0, 1)$ on \mathcal{N} . The lift of its flow to $\mathbb{T}^2 \times \mathbb{R}$ is defined by

$$\widetilde{\Phi}_t(\xi, z) = (\xi, z + t)$$

hence the first return map to $\{z = 0\}$ is $\Phi_1(\xi, 0) = (A\xi, 0)$ and is conjugate to A . The flow (Φ_t) is a suspension of A and has the same Lyapunov exponents as A .

We endow the manifold $\mathcal{M} = \mathcal{N} \times \mathbb{R}$ with the 1-form

$$\Lambda = \beta_- + s\beta_+$$

where s is the \mathbb{R} -coordinate. We define $\Omega = d\Lambda$. By (1), we have

$$\Omega = d\beta_- + ds \wedge \beta_+ + s d\beta_+ = dz \wedge (\ln \lambda_- \beta_- + s \ln \lambda_+ \beta_+) + ds \wedge \beta_+.$$

Thus $\Omega^{\wedge 2} = 2 \ln \lambda_- dz \wedge \beta_+ \wedge ds \wedge \beta_+ \neq 0$ and Ω is a symplectic form.

We define on \mathcal{M} the vector field $Y = X + 2 \ln \lambda_- \partial s$. Its flow is

$$\psi_t(\xi, z, s) = (\Phi_t(\xi, z), (\lambda_-)^{2t} s).$$

Hence $\mathcal{N} \times \{0\}$ is the global attractor for (ψ_t) . We have

$$\begin{aligned} \psi_t^* \Omega = & dz \wedge \left(\ln \lambda_- \cdot (\lambda_-)^t \beta_- + ((\lambda_-)^{2t} \cdot s) \ln \lambda_+ \cdot (\lambda_+)^t \beta_+ \right) + \\ & (\lambda_-)^{2t} ds \wedge (\lambda_+)^t \beta_+. \end{aligned}$$

As $\lambda_- \lambda_+ = 1$, we finally obtain

$$\psi_t^* \Omega = \lambda_-^t \Omega.$$

□

There are also examples of conformal symplectic diffeomorphisms on a non-exact symplectic manifold that have a non-isotropic invariant submanifold on which the restricted dynamics is Anosov.

Proposition 2. *There exists a conformal symplectic diffeomorphism f on a 6-dimensional symplectic manifold (\mathcal{M}, ω) , with a 4-dimensional invariant submanifold \mathcal{L} (so \mathcal{L} is not isotropic).*

Moreover, the submanifold \mathcal{L} is the global attractor for f , $f|_{\mathcal{L}}$ is conjugated to a hyperbolic automorphism of \mathbb{T}^4 with 2-dimensional stable and unstable foliations, and $f|_{\mathcal{L}}$ is transitive with entropy equal to $-\log a$, where a is the conformality ratio of f .

Question. In our example we have $a = \left(\frac{3-\sqrt{5}}{2}\right)^2$. In fact we can replace this number by the square of the largest eigenvalue of any Anosov automorphism of \mathbb{T}^2 . We don't know if we can achieve other constants by a conformal symplectic diffeomorphisms of the same symplectic manifold.

Proof. We consider the hyperbolic toral automorphism $T : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ that is defined by $T(\theta_1, \theta_2) = (2\theta_1 + \theta_2, \theta_1 + \theta_2)$. The associated linear map has eigenvalues $\lambda = \frac{3-\sqrt{5}}{2} < 1$ and $\lambda^{-1} = \frac{3+\sqrt{5}}{2} > 1$. Let $p = \frac{\sqrt{5}-1}{2}$. The unstable direction is spanned by $(1, p)$ and the stable one by $(1, -\frac{1}{p})$. The topological entropy is $-\log \lambda$ (see [8]).

Then the product map $F = (T, T) : (\theta_1, \theta_2, \theta_3, \theta_4) \in \mathbb{T}^2 \times \mathbb{T}^2 \mapsto (T(\theta_1, \theta_2), T(\theta_3, \theta_4))$ has topological entropy equal to $-2 \log \lambda$. We endow \mathbb{T}^4 with the closed 2-form Ω that is defined by

$$\Omega = (d\theta_2 - p d\theta_1) \wedge (d\theta_4 - p d\theta_3).$$

Observe that the kernel of Ω is the direction of the unstable foliation. Obviously, $F^* \Omega = \lambda^2 \Omega$. Now, we consider the subbundle

$$\mathcal{M} = \{(\theta, r) \in \mathbb{T}^4 \times \mathbb{R}^4; r_2 = pr_1 \quad \text{and} \quad r_4 = pr_3\}$$

of $\mathbb{T}^4 \times \mathbb{R}^4$. This bundle corresponds to the tangent bundle to the unstable foliation in the identification of $T\mathbb{T}^4$ with $\mathbb{T}^4 \times \mathbb{R}^4$.

We denote by Ω_1 the closed 2-form on \mathcal{M} that is equal to $\pi^*\Omega$ where $\pi : (\theta, r) \in \mathcal{M} \mapsto \theta \in \mathbb{T}^4$ and by Ω_2 the restriction of the usual symplectic form $d\theta \wedge dr$ of $T^*\mathbb{T}^4$ to \mathcal{M} :

$$\begin{cases} \Omega_1 = (d\theta_2 - p d\theta_1) \wedge (d\theta_4 - p d\theta_3) \\ \Omega_2 = \frac{1}{5}(d\theta_2 + \frac{1}{p}d\theta_1) \wedge (dr_2 + \frac{1}{p}dr_1) + \frac{1}{5}(d\theta_4 + \frac{1}{p}d\theta_3) \wedge (dr_4 + \frac{1}{p}dr_3). \end{cases}$$

Let then $\omega = \Omega_1 + \Omega_2$ be the chosen symplectic form on \mathcal{M} .

If we define $f : \mathcal{M} \rightarrow \mathcal{M}$ by $f(\theta, r) = (T(\theta), \left(\frac{3-\sqrt{5}}{2}\right)^3 r)$, then we have

$$\begin{cases} f^*\Omega_1 = \pi^*F^*\Omega = \lambda^2\Omega_1 \\ f^*\Omega_2 = \frac{\lambda^3}{\lambda}\Omega_2 = \lambda^2\Omega_2. \end{cases}$$

So finally $f : \mathcal{M} \rightarrow \mathcal{M}$ is a conformal symplectic diffeomorphism such that $f^*\omega = \lambda^2\omega$ and $f^*(\mathbb{T}^4 \times \{0\}) = \mathbb{T}^4 \times \{0\}$, where $\mathbb{T}^4 \times \{0\}$ is not isotropic and the topological entropy of $f|_{\mathbb{T}^4 \times \{0\}}$ is $-2 \log \lambda$. \square

Let us come back to the general case of a C^1 conformal symplectic diffeomorphism f of a symplectic manifold (\mathcal{M}, ω) , of conformality ratio $a \neq 1$.

Proposition 3. *A C^1 closed submanifold \mathcal{L} of even dimension which is invariant by f is nowhere symplectic. In particular, if \mathcal{L} is a surface, it is isotropic.*

Proof. Assume the conformality ratio a of f is $\neq 1$ and an f -invariant submanifold \mathcal{L} has dimension 2ℓ . Let $\Omega = \omega^\ell$, so that $f^*\Omega = a^\ell \Omega$. Choose a finite atlas $\mathcal{A} = \{(U_i, \Phi_i)\}_{1 \leq i \leq N}$ of \mathcal{L} , endow \mathcal{L} with a Riemannian metric and define

$$\begin{cases} \|\Omega\|_{\mathcal{L}, \infty} = \sup_{x \in \mathcal{L}, u_1, \dots, u_\ell \in T_x \mathcal{L} \setminus \{0\}} \frac{|\Omega(u_1, \dots, u_\ell)|}{\|u_1\| \dots \|u_\ell\|} \\ \|D\Phi_i^{-1}\|_{\mathcal{L}, \infty} = \sup_{u \in T\mathcal{L} \setminus \{0\}} \frac{\|D\Phi_i^{-1}(u)\|}{\|u\|}. \end{cases}$$

Then, $\int_U \Omega$ is bounded over open subsets U of \mathcal{L} :

$$(2) \quad \left| \int_U \omega \right| \leq \sum_{i=1}^N \int_{\Phi_i(U_i)} \|\omega\|_{\mathcal{L}, \infty} \|D\Phi_i^{-1}\|_{\mathcal{L}, \infty}^\ell d\text{Leb}.$$

Now, let U be any open set of \mathcal{L} . For $n \in \mathbb{Z}$, $f^n U$ is an open subset of \mathcal{L} and

$$\int_{f^n U} \Omega = a^n \int_U \Omega.$$

Since a is assumed $\neq 1$, $\int_U \Omega$ must thus be zero. Hence the 2ℓ -form induced by Ω vanishes identically, whence the conclusion. \square

If \mathcal{L} has any dimension, the same conclusion holds provided some constraint on the topological entropy $\text{ent}(f|_{\mathcal{L}})$ of the dynamics carried by \mathcal{L} . Define the spectral radius of a self-map g as

$$\text{rad}(Dg) = \limsup_{n \rightarrow +\infty} \|Dg^n\|_{\infty}^{\frac{1}{n}}.$$

Proposition 4. *Let f be a conformal diffeomorphism of (\mathcal{M}, ω) , i.e. such that $f^*\omega = a\omega$ with $a \in]0, 1[$. Let \mathcal{L} be an invariant closed submanifold. Assume one of the following hypothesis.*

(1) *The diffeomorphism f is smooth, \mathcal{L} is smooth and*

$$\text{ent}(f|_{\mathcal{L}}) < -\log(a);$$

(2) *The diffeomorphism f and \mathcal{L} are C^r for some $r \geq 1$ and*

$$\text{ent}(f|_{\mathcal{L}}) + \log^+ \left(\text{Rad}(Df|_{\mathcal{L}}^{-1})^{2/r} \right) < -\log(a).$$

Then \mathcal{L} is ω -isotropic.

Proof. We assume that \mathcal{L} is invariant and not isotropic. There exists a constant $k > 0$ such that on \mathcal{L} , we have $|\omega| \leq k|\text{vol}|$ where vol is the 2-dimensional volume form induced by the Riemannian metric. We choose in \mathcal{L} a small piece \mathcal{S} of symplectic surface (whose tangent space intersects the characteristic bundle of \mathcal{L} only in 0). Then $\omega(f^{-n}(\mathcal{S})) = a^{-n}\omega(\mathcal{S}) \neq 0$ and then

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |\text{vol}(f^{-n}(\mathcal{S}))| \geq \lim_{n \rightarrow \infty} \frac{1}{n} (\log |\omega(f^{-n}(\mathcal{S}))| - \log k) = -\log(a).$$

The conclusion follows from Yomdin's inequality, which we have recalled in appendix A. \square

Remark 4. This statement implies in particular that if \mathcal{L} is an invariant submanifold by a conformal flow (φ_t) then

- if \mathcal{L} and (φ_t) are C^1 and if $\varphi_t|_{\mathcal{L}}$ is C^1 conjugate to a rotation on a torus for some $t \neq 0$, then \mathcal{L} is isotropic; indeed, in this case, the entropy vanishes and the spectral radius of Df is 1. A simpler proof of this statement is given in [2].
- if \mathcal{L} and (φ_t) are smooth and if $\varphi_t|_{\mathcal{L}}$ is C^0 conjugate to a rotation on a torus for some $t \neq 0$, then \mathcal{L} is isotropic; indeed, in this case, the entropy vanishes.

3. ENTROPY

The purpose of this section is to improve regularity in Proposition 4. We will start by giving an abstract result on a manifold endowed with a form with constant rank and then we will give an application to invariant submanifolds of conformal symplectic dynamics.

Let

- $\mathcal{N}^{(n)}$ be a compact Riemannian C^2 manifold and d its distance
- \mathcal{F} be a C^2 foliation induced by a subbundle F of $T\mathcal{N}$ of rank $p \leq n - 1$
- Ω be an $(n - p)$ -form on \mathcal{N} which induces a volume on submanifolds transverse to \mathcal{F}
- f be a C^1 -diffeomorphism of \mathcal{N} preserving \mathcal{F} and such that

$$f^*\Omega = b\Omega$$

for some $b > 1$.

Theorem 5. *The topological entropy of f satisfies*

$$\text{ent } f \geq \ln b.$$

Proof. Key to the proof is the refined distance $d_{\mathcal{F}}$ on \mathcal{N} defined by

$$d_{\mathcal{F}}(x, y) = \begin{cases} \infty & \text{if } x \text{ and } y \text{ are not on the same leaf} \\ \text{distance from } x \text{ to } y \text{ along their common leaf} & \text{otherwise.} \end{cases}$$

Lemma 1. *There exist $\varepsilon > 0$ and $K > 0$ such that for every $x, y \in \mathcal{N}$*

$$(3) \quad d_{\mathcal{F}}(x, y) < \varepsilon \quad \Rightarrow \quad d_{\mathcal{F}}(x, y) \leq Kd(x, y).$$

Replacing the Riemannian metric d by $\frac{1}{\varepsilon}d$, we will assume that $\varepsilon = 1$.

Proof of Lemma 1. We choose $\varepsilon > 0$ that is strictly less than the radius of injectivity of the metric d restricted to every leaf and introduce

$$\mathcal{D} = \{(x, y) \in \mathcal{N} \times \mathcal{N}; d_{\mathcal{F}}(x, y) \leq \varepsilon\}.$$

This set is closed and due to our choice of ε , $d_{\mathcal{F}}$ is continuous on \mathcal{D} . If we use the notation

$$\Delta = \{(x, x); x \in \mathcal{N}\},$$

then the continuous function $\frac{d_{\mathcal{F}}}{d}$ is bounded on the complement of every neighbourhood of Δ in \mathcal{D} .

The exponential maps for the Riemannian form g and for the Riemannian form $g_{\mathcal{F}}$ restricted to the leaves are tangent along the tangent bundle to the leaves, hence

$$\lim_{(x,y) \rightarrow \Delta} \frac{d_{\mathcal{F}}(x, y)}{d(x, y)} = 1.$$

□

For every $x \in \mathcal{N}$, let $\mathcal{U}_x^{(n-p)}$ be a submanifold through x of dimension $n - p$, transverse to \mathcal{F} and homeomorphic to a ball, such that its normal bundle is trivial. Let \mathcal{V}_x be a tubular neighborhood of \mathcal{U}_x , of the form

$$\mathcal{V}_x = \cup_{y \in \mathcal{U}_x} \{z \in \mathcal{N}, d_{\mathcal{F}}(y, z) < \epsilon_x\}.$$

We choose \mathcal{U}_x and $\epsilon_x < 1$ small enough so that \mathcal{V}_x has a product structure. Furthermore, let

$$\mathcal{W}_x = \cup_{y \in \mathcal{U}_x} \{z \in \mathcal{N}, d_{\mathcal{F}}(y, z) < \epsilon_x/2\}.$$

Let $\mathcal{F}_{\mathcal{W}_x}$ be the foliation induced on \mathcal{W}_x by \mathcal{F} . (Due to the product structure, leaves of $\mathcal{F}_{\mathcal{W}_x}$ are of the form $\mathcal{W}_x \cap L_x$, where L_x is the leaf through x of the foliation induced on \mathcal{V}_x .) The neighborhood \mathcal{W}_x has the property that for any two points y and z of \mathcal{W}_x , if $d_{\mathcal{F}}(y, z) < \epsilon_x/2$ then y and z must belong to the same leaf of $\mathcal{F}_{\mathcal{W}_x}$; indeed, if y and z do not lie on the same leaf of $\mathcal{F}_{\mathcal{W}_x}$, their distance must be $\geq \epsilon_x$ since any path from y to z along a leaf of \mathcal{F} runs twice across $\mathcal{V}_x \setminus \mathcal{W}_x$.




FIGURE 1. Construction of the finite covering of \mathcal{N}

Let $\mathcal{W}_{x_1}, \dots, \mathcal{W}_{x_I}$ be a finite subcovering of \mathcal{N} . Denote \mathcal{W}_{x_i} by \mathcal{W}_i , and let $\epsilon = \min_i \epsilon_{x_i}/2$. So, the following property holds:

- (*) For every $i = 1, \dots, I$ and $y, z \in \mathcal{W}_i$ such that $d_{\mathcal{F}}(y, z) < \epsilon$, y and z belong to the same leaf of the foliation $\mathcal{F}_{\mathcal{W}_i}$ induced by \mathcal{F} on \mathcal{W}_i .

Moreover, since f^{-1} and \mathcal{F} are continuous and f preserves \mathcal{F} , there exists $\eta < \epsilon$ such that

- (**) For every $x, y \in \mathcal{N}$ such that $d_{\mathcal{F}}(x, y) < \eta$, $d_{\mathcal{F}}(f^{-1}x, f^{-1}y) < \epsilon$.

According to Lebesgue covering lemma, there exists $\theta < \eta/K$ such that every ball of radius θ is inside at least one of the \mathcal{W}_i 's.

Let $(Q_j)_{1 \leq j \leq J}$ be a decomposition of \mathcal{N} into cubes (or compact submanifolds with boundaries) such that each cube is contained in a ball of radius $< \theta$.

Let \mathcal{S} be a submanifold of \mathcal{N} of dimension $n - p$, included into some cube Q_j and transverse to \mathcal{F} . \mathcal{S} must lie into some \mathcal{W}_i . For any \mathcal{W}_i

containing \mathcal{S} , \mathcal{S} meets each leaf of $\mathcal{F}_{\mathcal{W}_i}$ at isolated points. By narrowing \mathcal{S} , we may assume that \mathcal{S} meets each leaf of $\mathcal{F}_{\mathcal{W}_i}$ at one point at most.

We claim that

(***) For every k and $j_1, \dots, j_k \in \{1, \dots, J\}$,

$$f^k(\mathcal{S}) \cap f^{k-1}(Q_{j_1}) \cap \dots \cap Q_{j_k}$$

meets each leaf of any \mathcal{W}_i containing Q_{j_k} at one point at most.

Let $j \in \{1, \dots, J\}$. Then $\mathcal{S}' = f(\mathcal{S}) \cap Q_j$ is also transverse to the foliation. Let $x, y \in \mathcal{S}'$ be on a common leaf of $\mathcal{F}_{\mathcal{W}_i}$, with $Q_j \subset \mathcal{W}_{i_0}$. Since such leaves have a diameter < 1 (due to our choice $\epsilon_x < 1$), using (3)⁸, we see that

$$d_{\mathcal{F}}(x, y) \leq Kd(x, y) \leq K \text{diam } Q_j \leq K\theta \leq \eta.$$

Using (**), $d_{\mathcal{F}}(f^{-1}x, f^{-1}y) < \epsilon$. But using (*), $f^{-1}x$ and $f^{-1}y$ belong to the same leaf of $\mathcal{F}_{\mathcal{W}_{i_0}}$. So, by the constructing property of \mathcal{W}_{i_0} , $f^{-1}x = f^{-1}y$ and $x = y$. By induction, (***) holds.

If $\mathcal{S} \subset \mathcal{W}_i$, we have

$$|\Omega(f^k(\mathcal{S}) \cap f^{k-1}(Q_{j_1}) \cap \dots \cap Q_{j_k})| \leq \max\{|\Omega(\mathcal{U}_1)|, \dots, |\Omega(\mathcal{U}_I)|\} = M,$$

uniformly with respect to k . Let

$$N_k = \#\{(j_1, \dots, j_k), f^k(\mathcal{S}) \cap f^{k-1}(Q_{j_1}) \cap \dots \cap Q_{j_k} \neq \emptyset\}.$$

Then

$$b^k |\Omega(\mathcal{S})| \leq N_k M,$$

hence

$$\frac{1}{k} \ln N_k \geq \frac{1}{k} \ln \frac{|\Omega(\mathcal{S})|}{M} + \ln b,$$

hence the wanted inequality. \square

Now assume that ω is a presymplectic form⁹ of \mathcal{N} of (even) rank $2\ell \geq 2$ and

$$f^*\omega = a\omega, \quad a > 1.$$

The kernel of ω is a uniquely integrable subbundle F of corank 2ℓ . Setting $\Omega = \omega^\ell$ and $b = a^\ell$ brings us back to the prior setting.

Corollary 1. *The topological entropy of $f : \mathcal{N} \hookrightarrow$ satisfies*

$$\text{ent } f \geq \frac{\text{rank}(\omega)}{2} \ln a.$$

Let us now return to our usual setting, where (\mathcal{M}, ω) is a symplectic manifold.

⁸Recall the metric was changed in order to have $\epsilon = 1$ in (3).

⁹A presymplectic form is a closed 2-form with constant rank.

Corollary 2. *Let $f : \mathcal{M} \hookrightarrow$ be a C^3 conformal symplectic diffeomorphism such that $f^*\omega = a\omega$ with $a > 1$. Suppose that \mathcal{N} is an invariant C^3 submanifold such that the induced form $\omega|_{\mathcal{N}}$ on \mathcal{N} has constant rank. Then*

$$\text{ent } f|_{\mathcal{N}} \geq \frac{\text{rank}(\omega|_{\mathcal{N}})}{2} \ln a;$$

in particular, if the entropy of $f|_{\mathcal{N}}$ vanishes, \mathcal{N} is isotropic.

Note that if \mathcal{N} is a compact submanifold such that $f|_{\mathcal{N}}$ is minimal,¹⁰ $\omega|_{\mathcal{N}}$ has constant rank and so the corollary applies.

Proof. As \mathcal{N} is C^3 , its tangent bundle is C^2 . Then Frobenius Theorem applies to $F = \ker \omega|_{\mathcal{N}}$ ¹¹ and the characteristic foliation \mathcal{F} exists. \square

4. LIOUVILLE CLASS OF INVARIANT SUBMANIFOLDS

In this section we assume that $(\mathcal{M}, \omega = -d\lambda)$ is an exact symplectic manifold. The goal is to prove that, given a conformal dynamics, there is only one Liouville class that an isotropic invariant submanifold may have.

4.1. Action of conformal dynamics on Liouville classes.

Definition 6. Let $j : \mathcal{S} \hookrightarrow \mathcal{M}$ be an isotropic embedding.

- Its *Liouville class* $[j] \in H^1(\mathcal{S}, \mathbb{R})$ is the cohomology class of the induced form $j^*\lambda$.
- It is *exact* if its Liouville class vanishes.

So, the notion of exactness is independent of the embedding with a given image.

When $\mathcal{M} = T^*\mathcal{Q}$ is the cotangent bundle of a closed manifold endowed with its tautological 1-form λ and \mathcal{L} is a Lagrangian submanifold of $T^*\mathcal{Q}$ that is homotopic to the zero section \mathcal{Z} , the restriction to \mathcal{L} of the canonical projection $\pi : T^*\mathcal{Q} \rightarrow \mathcal{Q}$ is a homotopy equivalence between \mathcal{L} and \mathcal{Q} and induces an isomorphism between $H^1(\mathcal{L}, \mathbb{R})$ and $H^1(\mathcal{Q}, \mathbb{R})$. Denoting by $j_{\mathcal{L}} : \mathcal{L} \hookrightarrow T^*\mathcal{Q}$ the canonical injection defined by $j_{\mathcal{L}}(x) = x$, the Liouville class of the submanifold \mathcal{L} is the cohomological class

$$[\mathcal{L}] = \left[(\pi|_{\mathcal{L}})_* (j_{\mathcal{L}}^*\lambda) \right] \in H^1(\mathcal{Q}, \mathbb{R}).$$

In this case, we may thus update the definition of Liouville classes.

¹⁰By definition, it is minimal if every orbit is dense

¹¹The infinitesimal integrability condition is well known: if X, Y are sections of F and Z is a section of $T\mathcal{N}$, $0 = d\omega(X, Y, Z) = -\omega([X, Y], Z)$, which shows that $[X, Y]$ itself is a section of F .

Definition 7. Let \mathcal{L} be a Lagrangian submanifold of $T^*\mathcal{Q}$ that is homotopic to the zero section, the Liouville class $[\mathcal{L}]$ of \mathcal{L} is the cohomology class on \mathcal{Q} whose pull back by $\pi|_{\mathcal{L}}$ is the cohomology class of $\lambda|_{T\mathcal{L}}$.

The following straightforward proposition explains that the group of conformal dynamics acts on the set of Liouville classes of isotropic embeddings that are homotopic to a given isotropic embedding of a given manifold \mathcal{S} by homotheties (translations when the dynamics is symplectic).

Proposition 5. *Let $f : \mathcal{M} \hookrightarrow$ be a conformal diffeomorphism with conformality ratio a . Then $\eta = f^*\lambda - a\lambda$ is a closed 1-form.*

Let $j_0 : \mathcal{S} \hookrightarrow \mathcal{M}$ be an isotropic embedding. For every isotropic embedding $j : \mathcal{S} \hookrightarrow \mathcal{M}$ that is homotopic to j_0 , the Liouville class of the isotropic embedding $f \circ j : \mathcal{S} \hookrightarrow \mathcal{M}$ is

$$[f \circ j] = a[j] + [j_0^*\eta].$$

Proof. We have $d\eta = -f^*\omega + af^*\omega = 0$ and η is closed. For $\gamma : \mathbb{T} \hookrightarrow \mathcal{M}$, let us compute

$$\begin{aligned} [f \circ j][\gamma] &= \int_{\gamma} (f \circ j)^*\lambda = \int_{\gamma} j^*(f^*\lambda) = \int_{\gamma} j^*(\eta + a\lambda) \\ &= \int_{\gamma} j^*\eta + a \int_{j \circ \gamma} \lambda = a[j][\gamma] + [\eta][j \circ \gamma] \\ &= a[j][\gamma] + [\eta][j_0 \circ \gamma] = a[j][\gamma] + [j_0^*\eta][\gamma]. \end{aligned}$$

□

Definition 8. A diffeomorphism $f : \mathcal{M} \hookrightarrow$ is λ conformal Hamiltonian (CH) if there exists an isotopy $(f_t)_{t \in [0,1]}$ such that $f_0 = \text{Id}_{\mathcal{M}}$, $f_1 = f$ and two functions $H : [0, 1] \times \mathcal{M} \rightarrow \mathbb{R}$ and $\alpha : [0, 1] \rightarrow \mathbb{R}$ such that

$$\forall (t, x) \in [0, 1] \times \mathcal{M}, i_{j_t(x)}\omega = \alpha(t)\lambda + \partial_x H(t, x).$$

Remark 9. A diffeomorphism $f : \mathcal{M} \hookrightarrow$ is conformal Hamiltonian if and only if there exists an isotopy $(f_t)_{t \in [0,1]}$ of CES diffeomorphisms such that $f_0 = \text{Id}_{\mathcal{M}}$ and $f_1 = f$.

Definition 10. The flow (φ_t) associated to the vector field X on \mathcal{M} is λ conformal Hamiltonian if there exists $\alpha \in \mathbb{R}$ and $H : \mathcal{M} \rightarrow \mathbb{R}$ such that $i_X\omega = \alpha\lambda + dH$.

Remark 11. A flow is a flow of λ conformal exact symplectic diffeomorphisms if and only if it is λ conformal Hamiltonian.

To describe the behavior of Lagrangian submanifolds of $T^*\mathcal{Q}$ that are H-isotopic to a graph, we first need the following invariance result.

Proposition 6. *Let (\mathcal{L}_t) be an isotopy of Lagrangian submanifolds of $T^*\mathcal{Q}$ such that $\mathcal{L}_0 = \mathcal{Z}$. Then \mathcal{L}_1 is H -isotopic to a graph.*

Corollary 3. *Let $(g_t)_{t \in [0,1]}$ be an isotopy of conformal symplectic diffeomorphisms such that $g_0 = \text{Id}_{T^*\mathcal{Q}}$. Let \mathcal{L} be a Lagrangian submanifold of $T^*\mathcal{Q}$ that is H -isotopic to a graph. Then $g_1(\mathcal{L})$ is H -isotopic to a graph. If moreover \mathcal{L} is H -isotopic to the zero-section and the isotopy is conformal Hamiltonian, then $g_1(\mathcal{L})$ is H -isotopic to the zero-section.*

Proof of Proposition 6. We will prove

Lemma 2. *Assume that \mathcal{L} is H -isotopic to the zero section and that $(\mathcal{L}_t)_{t \in [-\varepsilon, \varepsilon]}$ is an isotopy of exact Lagrangian submanifolds such that $\mathcal{L}_0 = \mathcal{L}$. Then there exists a neighbourhood \mathcal{N} of 0 in $[-\varepsilon, \varepsilon]$ such that for every $t \in \mathcal{N}$, \mathcal{L}_t is H -isotopic to the zero section.*

Proof of Lemma 2. We use Weinstein tubular neighbourhood Theorem, [21]. Let \mathcal{T} be a symplectic tubular of \mathcal{L} , i.e. there exists a neighbourhood \mathcal{U} of the zero section in $T^*\mathcal{L}$ and a symplectic embedding $\phi : \mathcal{U} \hookrightarrow T^*\mathcal{Q}$ with image \mathcal{T} that is $\text{Id}_{\mathcal{L}}$ on \mathcal{L} . As Φ maps the exact Lagrangian submanifold \mathcal{L} of $T^*\mathcal{L}$ onto the exact Lagrangian submanifold \mathcal{L} of $T^*\mathcal{Q}$, then Φ is exact symplectic.

This implies that every submanifold $\phi^{-1}(\mathcal{L}_t)$ is exact Lagrangian. Moreover, there exists a neighbourhood \mathcal{N} of 0 in $[-\varepsilon, \varepsilon]$ such that for every $t \in \mathcal{N}$, $\phi^{-1}(\mathcal{L}_t)$ is a graph. Hence this is the graph of an exact 1-form du_t . Then $\phi^{-1}(\mathcal{L}_t)$ is the image by the time-1 Hamiltonian flow of $H = -\frac{du_t}{dt} \circ \pi$. Using a bump function, we can assume that H has support in \mathcal{U} , and then the time-1 map of the Hamiltonian $H \circ \phi$ maps \mathcal{L} onto \mathcal{L}_t . \square

We now prove Proposition 6. Let us firstly deal with the case when all the \mathcal{L}_t s are exact. We introduce

$$\{t \in [0, 1]; \forall s \in [0, t], g_s(\mathcal{L}) \text{ is } H\text{-isotopic to the zero section}\}.$$

Lemma 2 and the transitivity of the relation of H -isotopy imply that this set is closed and open in $[0, 1]$, hence equal to $[0, 1]$.

Now we just assume that (\mathcal{L}_t) is an isotopy of Lagrangian submanifolds of $T^*\mathcal{Q}$ such that $\mathcal{L}_0 = \mathcal{Z}$. We choose an arc $(\eta_t)_{t \in [0,1]}$ of closed 1-forms on \mathcal{Q} whose cohomology class $[\eta_t] = [\mathcal{L}_t]$ is the Liouville class of \mathcal{L}_t . We denote by $T_t : T^*\mathcal{Q} \hookrightarrow T^*\mathcal{Q}$ the symplectic diffeomorphisms such that $T_t(p) = p + \eta_t \circ \pi(p)$. Then $\mathcal{L}_t^* = T_{-t}(\mathcal{L}_t)$ defines a homotopy of exact Lagrangian submanifolds of $T^*\mathcal{Q}$. A result of the first part of the proof is that \mathcal{L}_t^* is H -isotopic to the zero section, i.e. there exists a H -isotopy $(\phi_t)_{t \in [0,1]}$ such that $\phi_0 = \text{Id}$ and $\phi_1(\mathcal{Z}) = \mathcal{L}_1^*$. Hence $\mathcal{L}_1 = T_1(\mathcal{L}_1^*)$ is H -isotopic to the graph of η_1 via the H -isotopy

$$(\psi_t)_{t \in [0,1]} = (T_1 \circ \phi_t \circ T_1^{-1})_{t \in [0,1]}.$$

□

Proof of Corollary 3. We assume that $(g_t)_{t \in [0,1]}$ is an isotopy of conformal symplectic diffeomorphisms such that $g_0 = \text{Id}_{T^*\mathcal{Q}}$ and that \mathcal{L} is a Lagrangian submanifold of $T^*\mathcal{Q}$ that is H-isotopic to a graph. Then there exist a closed 1-form η on \mathcal{Q} and a H -isotopy $(h_t)_{t \in [0,1]}$ such that $h_0 = \text{Id}_{T^*\mathcal{Q}}$ and $\mathcal{L} = h_1(\text{graph}(\eta))$. We introduce the symplectic diffeomorphisms $(T_t)_{t \in [0,1]}$ of $T^*\mathcal{Q}$ that are defined by $T_t(p) = p + t\eta \circ \pi(q)$. Then

$$(\mathcal{L}_t)_{t \in [0,1]} = (g_t \circ h_t \circ T_t(\mathcal{Z}))_{t \in [0,1]}$$

is a isotopy of Lagrangian submanifolds such that $\mathcal{L}_0 = \mathcal{Z}$ and $\mathcal{L}_1 = g_1(\mathcal{L})$. A result of Proposition 6 is that $g_1(\mathcal{L})$ is H-isotopic to a graph.

If moreover \mathcal{L} is H-isotopic to the zero-section and the isotopy is conformal Hamiltonian, then all the maps $g_t \circ h_t \circ T_t$ are conformal Hamiltonian and thus every manifold \mathcal{L}_t is exact Lagrangian. The conclusion is a result of the second part of Corollary 3. □

4.2. Liouville classes of invariant submanifolds. Let $j_0 : \mathcal{S} \hookrightarrow \mathcal{M}$ be an isotropic embedding. We denote by $\mathcal{J}(j_0)$ the set of isotropic embeddings $j : \mathcal{S} \hookrightarrow \mathcal{M}$ that are homotopic to j_0 .

A consequence of Proposition 5 is

Proposition 7. *Let $f : \mathcal{M} \hookrightarrow \mathcal{M}$ be a conformal diffeomorphism. Let $j \in \mathcal{J}(j_0)$ be an isotropic embedding which is strongly f -invariant in the sense that*

- $j(\mathcal{S}) = f \circ j(\mathcal{S})$
- f acts trivially on $H^1(j(\mathcal{S}), \mathbb{R})$.

Then j may have only one Liouville class, that we denote by $[\ell_f(\mathcal{J}(j_0))]$. In particular, when f is CES, then $[\ell_f(\mathcal{J}(j_0))] = 0$ and j has to be exact.

Proof. Let $j : \mathcal{S} \hookrightarrow \mathcal{M}$ be such an embedding. With the notations of Proposition 5, we have

$$[f \circ j] = a[j] + [j_0^* \eta].$$

Let us denote by $i : j(\mathcal{S}) \hookrightarrow \mathcal{M}$ the canonical injection. As f acts trivially on $H^1(j(\mathcal{S}), \mathbb{R})$, we have

$$[f \circ j] = \left[(i \circ f \circ j)^* \lambda \right] = j^* \left[f^*(i^* \lambda) \right] = [j^* \lambda] = [j]$$

and finally $[j]$ has to be the only fixed point of the homothety that maps $[j]$ on $a[j] + [j_0^* \eta]$. □

As a consequence:

Proposition 8. *Let $f : \mathcal{M} \looparrowright$ be a λ CES diffeomorphism. Then every invariant isotropic submanifold \mathcal{S} such that $f|_{\mathcal{S}}$ acts trivially on $H^1(\mathcal{S})$ is exact.*

Corollary 4. *Let X be a CS vector field on \mathcal{M} with flow (φ_t) . Let $j_0 : \mathcal{S} \hookrightarrow \mathcal{M}$ be an isotropic embedding. We denote by $\mathcal{J}(j_0)$ the set of isotropic embeddings $j : \mathcal{S} \hookrightarrow \mathcal{M}$ that are homotopic to j_0 . Then there is only one Liouville class that we denote by $[\ell_X(\mathcal{J})]$, that an isotropic embedding $j \in \mathcal{J}(j_0)$ such that*

$$\forall t \in \mathbb{R}, \varphi_t(j(\mathcal{S})) = j(\mathcal{S})$$

may have.

In particular, when X is CH, then $[\ell_X(\mathcal{J})] = 0$.

Corollary 5. *Let $f : T^*\mathcal{Q} \looparrowright$ be a CS-diffeomorphism that is homotopic to $\text{Id}_{T^*\mathcal{Q}}$. Then there is only one Liouville class that we denote by $[\ell_f]$, that a homotopic to the zero section and f -invariant submanifold may have.*

Proof of corollary 5. Let $j_0 : \mathcal{Q} \hookrightarrow T^*\mathcal{Q}$ be the canonical injection onto the zero-section.

We assume that \mathcal{L} is an f -invariant submanifold that is isotopic to the zero section. Because

- $\pi|_{\mathcal{L}}$ defines an homotopy equivalence between \mathcal{L} and \mathcal{Q} ;
- π defines an homotopy equivalence between $T^*\mathcal{Q}$ and \mathcal{Q} ;
- f is homotopic to $\text{Id}_{T^*\mathcal{Q}}$,

then f acts trivially on $H^1(\mathcal{L}, \mathbb{R})$.

Let (ψ_t) be an isotopy of diffeomorphisms of $T^*\mathcal{Q}$ such that $\psi_0 = \text{Id}_{T^*\mathcal{Q}}$ and $\psi_1(\mathcal{Z}) = \mathcal{L}$. Then $\psi_1 \circ j_0 \in \mathcal{J}(j_0)$ and a result of Proposition 7 is that $[\psi_1 \circ j_0] = [\ell_f(\mathcal{J}(j_0))]$.

Moreover, if $i_{\mathcal{L}} : \mathcal{L} \hookrightarrow T^*\mathcal{Q}$ is the canonical injection, we have

$$[\psi_1 \circ j_0] = [i_{\mathcal{L}} \circ \psi_1 \circ j_0] = [(\psi_1 \circ j_0)^*(i_{\mathcal{L}}^*\lambda)].$$

Observe that $\psi_1 \circ j_0 : \mathcal{Q} \rightarrow \mathcal{L}$ is an homotopy equivalence such that and $\pi \circ (\psi_1 \circ j_0)$ acts trivially on $H^1(\mathcal{Q}, \mathbb{R})$. We deduce that

$$[\psi_1 \circ j_0] = [(\psi_1 \circ j_0)^*(i_{\mathcal{L}}^*\lambda)] = [\pi_*(i_{\mathcal{L}}^*\lambda)] = [\mathcal{L}].$$

and then $[\mathcal{L}] = [\ell_f(\mathcal{J}(j_0))]$. □

5. LIOUVILLE CLASS OF LAGRANGIAN SUBMANIFOLDS OF $T^*\mathcal{Q}$ WITH COMPACT ORBITS

The goal of this section is to prove that, given a conformal dynamics on $T^*\mathcal{Q}$, there is only one Liouville class that a Lagrangian submanifold with compact orbit may have.

We assume that $\mathcal{M} = T^*\mathcal{Q}$ and that $f : \mathcal{M} \looparrowright$ is CS-isotopic to $\text{Id}_{\mathcal{M}}$.

We suppose that $j : \mathcal{Q} \hookrightarrow \mathcal{M}$ is a Lagrangian embedding such that $j(\mathcal{Q}) = \mathcal{L}$ is H-isotopic to a graph and has compact orbit (for example is contained in some compact attracting set).

Theorem 12. *Let $f : \mathcal{M} \hookrightarrow \mathcal{M}$ be a diffeomorphism that is CS-isotopic to $\text{Id}_{\mathcal{M}}$ and let \mathcal{L} be a Lagrangian submanifold that is isotopic to the zero section among Lagrangian submanifolds and such that $\bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L})$ is relatively compact, then $[\mathcal{L}] = \ell_f$.*

Corollary 6. *Let (φ_t) be the flow of the conformal symplectic vector field X and let \mathcal{L} be a Lagrangian submanifold that is isotopic to the zero section among the Lagrangian submanifolds of $T^*\mathcal{Q}$ such that $\bigcup_{t \in \mathbb{R}} \varphi_t(\mathcal{L})$ is relatively compact, then $[\mathcal{L}] = \ell_X$.*

Remark 13. We give a proof of Theorem 12 that uses the notion of graph selector. If \mathcal{Q} (as \mathbb{T}^n) satisfies that every element of $H^1(\mathcal{Q}, \mathbb{R}) \setminus \{0\}$ contains a non-vanishing 1-form, we can give a simpler proof. Indeed, in the proof, we are reduced to prove that if we have a sequence (\mathcal{L}_n) of Lagrangian submanifolds such that $[\mathcal{L}_n] = a^n([\mathcal{L}_0] - \ell_f) + \ell_f$ tends to infinity as $n \rightarrow \infty$, then $\bigcup_{n \in \mathbb{N}} \mathcal{L}_n$ is not relatively compact. If $\eta = f^*\lambda - \lambda$ and the 1-form ν_1 on \mathcal{Q} is non-vanishing and represents $[\mathcal{L}_0] - \ell_f$, then \mathcal{L}_n and the graph of $\frac{1}{1-a}\eta + a^n\nu_1$ intersect. As ν_1 doesn't vanish, we can conclude.

Proof of Theorem 12. We endow \mathcal{Q} with a Riemannian metric and denote by $\|\cdot\|$ the norm on $T\mathcal{Q}$. Changing f into f^{-1} , we can assume that $a > 1$. As f is CS, then $f^*\lambda - a\lambda = \eta$ is closed, We deduce from the proof of Proposition 7 that $\ell_f = \frac{1}{1-a}[j^*\eta]$ where j is the canonical injection from \mathcal{Q} in $T^*\mathcal{Q} = \mathcal{M}$ on the zero section.

Then f^k is also CS with

$$(f^k)^*\lambda - a^k\lambda = \sum_{j=0}^{k-1} a^{k-j-1}(f^j)^*(f^*\lambda - a\lambda) = \sum_{j=0}^{k-1} a^{k-j-1}(f^j)^*\eta.$$

Suppose ad absurdum that $[\mathcal{L}]$ is not ℓ_f . Let ν be a closed 1-form on \mathcal{Q} such that $\ell_f + [\nu] = [\mathcal{L}]$. There is a loop $\gamma : \mathbb{T} \rightarrow \mathcal{Q}$ such that $\int_{\gamma} \nu \neq 0$.

As f is CS-isotopic to $\text{Id}_{\mathcal{M}}$ and by transitivity of the relation of CS-isotopy, f^k is also CS-isotopic to $\text{Id}_{\mathcal{M}}$. Hence by Corollary 3, $f^k(\mathcal{L})$ is H-isotopic to a Lagrangian graph. The submanifold \mathcal{L} is H-isotopic to the graph of $\frac{1}{1-a}j^*\eta + \nu$. A result of Proposition 5 is that $f^k(\mathcal{L})$ is H-isotopic to the graph of

$$a^k \left(\frac{1}{1-a} j^* \eta + \nu \right) + \sum_{j=0}^{k-1} a^{k-j-1} j^* \eta = a^k \nu + \frac{1}{1-a} j^* \eta.$$

If we denote by $\tau_k : \mathcal{M} \looparrowright$ the symplectic diffeomorphisms $\tau_k(p) = p + a^k \nu(\pi(p)) + \frac{1}{1-a} \eta(j(\pi(p)))$, then $\tau_k^{-1} \circ f^k(\mathcal{L})$ is H -isotopic to the zero section and then admits a generating function and a graph selector that is (see e.g. [14] p 98 and references herein) a Lipschitz function $u_k : \mathcal{Q} \rightarrow \mathbb{R}$ that is C^1 on an open subset \mathcal{U}_k of \mathcal{Q} with full Lebesgue measure such that

$$\forall q \in \mathcal{U}_k, du_k(q) \in \tau_k^{-1} \circ f^k(\mathcal{L}).$$

Using Fubini theorem, we find a loop γ_k that is C^1 close to γ and such that

- γ_k is smooth and isotopic to γ ;
- for Lebesgue almost $s \in \mathbb{T}$, we have $\gamma_k(s) \in \mathcal{U}_k$.

As $u_k \circ \gamma_k$ is Lipschitz and then absolutely continuous, we have

$$0 = \int_{\mathbb{T}} \frac{d(u_k \circ \gamma_k)}{ds}(s) ds.$$

Because $\gamma_k(s) \in \mathcal{U}_0$ for almost every s , we deduce

$$0 = \int_{\mathbb{T}} du_k(\gamma_k(s)) \cdot \gamma'_k(s) ds$$

and because γ_k is homotopic to γ and $a^k \nu + \frac{1}{1-a} j^* \eta$ is closed,

$$\int_{\mathbb{T}} \left(a^k \nu(\gamma_k(s)) + \frac{1}{1-a} \eta(j(\gamma_k(s))) + du_k(\gamma_k(s)) \right) \cdot \gamma'_k(s) ds = a^k \int_{\gamma} \nu + \frac{1}{1-a} \int_{\gamma} j^* \eta$$

As the loops γ_k are C^1 -close to γ , there exists a constant K that is an upper bound for all the $\|\gamma'_k(s)\|$. Hence there is a subset E_k with non-zero Lebesgue measure of \mathbb{T} such that for every $s \in E_k$, we have

$$(4) \quad \left\| a^k \nu(\gamma_k(s)) + \frac{1}{1-a} \eta(j(\gamma_k(s))) + du_k(\gamma_k(s)) \right\| \geq \frac{a^k}{2K} \left| \int_{\gamma} \nu \right| - \frac{1}{(1-a)K} \left| \int_{\gamma} \eta \right|.$$

Moreover, for almost every $s \in \mathbb{T}$, we have

$$du_k(\gamma_k(s)) \in \tau_k^{-1} \circ f^k(\mathcal{L})$$

i.e.

$$(5) \quad a^k \nu(\gamma_k(s)) + \frac{1}{1-a} \eta(j(\gamma_k(s))) + du_k(\gamma_k(s)) \in f^k(\mathcal{L}).$$

We deduce from (4) and (5) that there is $p \in f^k(\mathcal{L})$ such that

$$\|p\| \geq \frac{a^k}{2K} \left| \int_{\gamma} \nu \right| - \frac{1}{(1-a)K} \left| \int_{\gamma} \eta \right|.$$

□

Question. Is the hypothesis on H -isotopy to the zero section necessary?

6. UNIQUENESS

We work on the cotangent bundle $(T^*\mathcal{Q}, -d\lambda)$ of a closed orientable manifold.

Viterbo introduced in the seminal paper [17], see also [19], the spectral distance γ that is defined on the set of H-isotopic to the zero-section Lagrangian submanifolds.

We will recall the main results of this theory and apply this to prove that if two submanifolds $\mathcal{L}, \mathcal{L}'$ are H-isotopic to the zero section and if (φ_t) is a CH flow of $T^*\mathcal{Q}$, then

$$\text{either } \gamma(\varphi_t(\mathcal{L}), \varphi_t(\mathcal{L}')) \xrightarrow{t \rightarrow +\infty} +\infty \quad \text{or} \quad \gamma(\varphi_t(\mathcal{L}), \varphi_t(\mathcal{L}')) \xrightarrow{t \rightarrow -\infty} +\infty.$$

Using a recent result due to Shelukhin, [15], we will deduce that for certain manifolds \mathcal{Q} , e.g. tori \mathbb{T}^n , there is at most one H-isotopic to the zero section submanifold whose orbit is compact and when it exists, this submanifold is in fact invariant.

6.1. On Viterbo spectral distance γ . If $\mathcal{L}, \mathcal{L}'$ are H-isotopic to the zero section submanifolds of $T^*\mathcal{Q}$, they have quadratic at infinity *generating functions* $S : \mathcal{Q} \times \mathbb{R}^k \rightarrow \mathbb{R}$ and $S' : \mathcal{Q} \times \mathbb{R}^{k'} \rightarrow \mathbb{R}$.

We recall that a generating function S for \mathcal{L} is such that

- if we use the notation $(q, \xi) \in \mathcal{Q} \times \mathbb{R}^k$, on $\Sigma_S = \left(\frac{\partial S}{\partial \xi}\right)^{-1}(0)$, $\frac{\partial S}{\partial \xi}$ has maximal rank;
- the map $j_S : \Sigma_S \hookrightarrow T^*\mathcal{Q}$ defined by $j_S(q, \xi) = \frac{\partial S}{\partial q}(q, \xi)$ is an embedding and its image is \mathcal{L} .

The generating function is *quadratic at infinity* if there exists a non-degenerate quadratic form $Q : \mathbb{R}^k \rightarrow \mathbb{R}$ such that outside a compact subset of $\mathcal{Q} \times \mathbb{R}^k$, we have $S(q, \xi) = Q(\xi)$.

The function $S \ominus S' : \mathcal{Q} \times \mathbb{R}^k \times \mathbb{R}^{k'} \rightarrow \mathbb{R}$ is defined by

$$(S \ominus S')(q, \xi, \chi) = S(q, \xi) - S'(q, \chi).$$

Observe that

$$\mathcal{L} \cap \mathcal{L}' = \left\{ \frac{\partial S}{\partial q}(q, \xi); d(S \ominus S')(q, \xi, \chi) = 0 \right\} = \left\{ \frac{\partial S'}{\partial q}(q, \chi); d(S \ominus S')(q, \xi, \chi) = 0 \right\}.$$

The function $S \ominus S'$ is not quadratic at infinity, but it satisfies conditions of Proposition 1.6. of [18] that ensure that it can be replaced by such a function, which we also denote by $S \ominus S'$. There exists a compact set $K \subset \mathcal{Q} \times \mathbb{R}^k \times \mathbb{R}^{k'}$ such that

$$\forall (q, \xi, \chi) \notin K, (S \ominus S')(q, \xi, \chi) = Q(\xi, \chi)$$

where Q is a non degenerate quadratic form on $\mathbb{R}^k \times \mathbb{R}^{k'}$. We denote by m its index. Moreover, there exist $a, b \in \mathbb{R}$ such that

$$K \cap \left(\{(S \ominus S') \geq b\} \cup \{(S \ominus S') \leq a\} \right) = \emptyset.$$

For $c \in \mathbb{R}$, we denote by \mathcal{E}^c and \mathcal{F}^c the sublevels

$$\mathcal{E}^c = \{(q, \xi, \chi); (S \ominus S')(q, \xi, \chi) \leq c\} \quad \text{and} \quad \mathcal{F}^c = \{(\xi, \chi); Q(\xi, \chi) \leq c\}.$$

As $(S \ominus S')(q, \xi, \chi)$ and $Q(\xi, \chi)$ are equal on \mathcal{E}^a and outside \mathcal{E}^b , we have

$$\forall c \notin]a, b[, \mathcal{E}^c = \mathcal{Q} \times \mathcal{F}^c.$$

Hence, by Kunneth theorem [3], there is an isomorphism

$$K : H(\mathcal{F}^b, \mathcal{F}^a) \otimes H(\mathcal{Q}) \rightarrow H(\mathcal{E}^b, \mathcal{E}^a).$$

As Q is a non-degenerate quadratic form with index m , we have $H^p(\mathcal{F}^b, \mathcal{F}^a) = \{0\}$ for $p \neq m$ and $H^m(\mathcal{F}^b, \mathcal{F}^a) = \mathbb{R}C$ is one dimensional. We deduce an isomorphism

$$T : \mathbb{R}C \otimes H^*(\mathcal{Q}) \rightarrow H^{*+m}(\mathcal{E}^b, \mathcal{E}^a).$$

Then, if $\alpha \in H^*(\mathcal{Q})$ is non-zero,

$$c(\alpha, S \ominus S') = \inf\{t \in [a, b], j_t^*(C \otimes \alpha) \neq 0\}$$

where $j_t : (\mathcal{E}^t, \mathcal{E}^a) \rightarrow (\mathcal{E}^b, \mathcal{E}^a)$ is the inclusion. The number $c(\alpha, S \ominus S')$ is then a critical value of $S \ominus S'$ that continuously depend on S and S' for the uniform C^0 distance.

Viterbo proved that $c(\alpha, S \ominus S')$ depends only on \mathcal{L} and \mathcal{L}' and not on the choice of generating functions. It is then denoted by $c(\alpha, \mathcal{L}, \mathcal{L}')$.

If μ is the orientation class of \mathcal{Q} , the distance $\gamma(\mathcal{L}, \mathcal{L}')$ is defined by

$$\gamma(\mathcal{L}, \mathcal{L}') = c(\mu, \mathcal{L}, \mathcal{L}') - c(1, \mathcal{L}, \mathcal{L}').$$

Theorem 14. *Let $f : \mathcal{M} \hookrightarrow \mathcal{M}$ be a CES diffeomorphism that is CH-isotopic to $\text{Id}_{T^*\mathcal{Q}}$. Let $\mathcal{L}, \mathcal{L}'$ be two distinct submanifolds of $T^*\mathcal{Q}$ which are H-isotopic to the zero section, then*

$$\text{either } \gamma(f^n(\mathcal{L}), f^n(\mathcal{L}')) \xrightarrow{n \rightarrow +\infty} +\infty$$

$$\text{or } \gamma(f^{-n}(\mathcal{L}), f^{-n}(\mathcal{L}')) \xrightarrow{n \rightarrow +\infty} +\infty.$$

Corollary 7. *Let $f : \mathcal{M} \hookrightarrow \mathcal{M}$ be a CES diffeomorphism that is CH-isotopic to $\text{Id}_{T^*\mathcal{Q}}$. Then there exists at most one H-isotopic to the zero section submanifold of $T^*\mathcal{Q}$ that is invariant by f .*

Proof of Theorem 14. This is direct application of the following result of which we provide a proof.

Lemma 3. *Let $\mathcal{L}, \mathcal{L}'$ be two H -isotopic to the zero section submanifolds of $T^*\mathcal{Q}$. Let (ϕ_t) be an isotopy of exact conformal symplectic diffeomorphisms of $T^*\mathcal{Q}$ such that $\phi_0 = \text{Id}_{T^*\mathcal{Q}}$ and $\phi_t^*\omega = a(t)\omega$. Then*

$$\gamma(\phi_t(\mathcal{L}), \phi_t(\mathcal{L}')) = a(t)\gamma(\mathcal{L}, \mathcal{L}').$$

Proof. As the distance γ continuously depends on the generating functions, we only need to prove the results for submanifolds \mathcal{L} and \mathcal{L}' whose intersections are all transverse. In this case, there is only a finite number of critical points and critical values for $S \ominus S'$. If $x, y \in \mathcal{L} \cap \mathcal{L}'$, we denote by $\Delta(x, y, \mathcal{L}, \mathcal{L}')$ the difference of the corresponding critical values of $S \ominus S'$, i.e.

$$\Delta(x, y, \mathcal{L}, \mathcal{L}') = \left(S \circ j_S^{-1}(y) - S' \circ j_{S'}^{-1}(y) \right) - \left(S \circ j_S^{-1}(x) - S' \circ j_{S'}^{-1}(x) \right).$$

Then if η_1 is a path in \mathcal{L} joining x to y and η_2 a path in \mathcal{L}' joining y to x , the difference of the two corresponding critical values of $S \ominus S'$ is

$$\Delta(x, y, \mathcal{L}, \mathcal{L}') = \int_{\eta_1 \vee \eta_2} \lambda.$$

We can always choose η_1 and η_2 that are homotopic with fixed ends. Then, if \mathcal{D} is a disc with boundary $\eta_1 \vee \eta_2$, we have

$$\Delta(x, y, \mathcal{L}, \mathcal{L}') = \int_{\mathcal{D}} \omega.$$

The intersection points of $\phi_t(\mathcal{L})$ and $\phi_t(\mathcal{L}')$ are the points $\phi_t(x)$ with $x \in \mathcal{L} \cap \mathcal{L}'$. For x, y in $\mathcal{L} \cap \mathcal{L}'$, we have

$$\Delta(\phi_t(x), \phi_t(y), \phi_t(\mathcal{L}), \phi_t(\mathcal{L}')) = \int_{\Phi_t(\mathcal{D})} \omega = a(t) \int_{\mathcal{D}} \omega = a(t)\Delta(x, y, \mathcal{L}, \mathcal{L}').$$

Hence $t \mapsto \frac{1}{a(t)} \left(c(\mu, \phi_t(\mathcal{L}), \phi_t(\mathcal{L}')) - c(1, \phi_t(\mathcal{L}), \phi_t(\mathcal{L}')) \right)$ is a continuous map that takes its values in a fixed finite set, it has to be constant. \square

\square

6.2. An application of a result of Shelukhin.

Theorem 15. *Let $f : T^*\mathbb{T}^n \hookrightarrow$ be a CES diffeomorphism that is CH -isotopic to $\text{Id}_{T^*\mathbb{T}^n}$. Then there exists at most one H -isotopic to the zero section submanifold \mathcal{L} such that*

$$\bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L}) \text{ is relatively compact.}$$

Hence when it exists, \mathcal{L} is invariant by the f .

Proof. In [15], Shelukhin defines a notion of string-point invertible manifold. The tori \mathbb{T}^n are examples of such manifolds. His result implies

Theorem (Shelukhin,[15]). *Let g be a Riemannian metric on \mathbb{T}^n . Then there exists a constant $C(g)$ such that for all exact Lagrangian submanifolds $\mathcal{L}_0, \mathcal{L}_1$ contained in the unit codisk bundle $D^*(g) \subset T^*\mathbb{T}^n$, we have $\gamma(\mathcal{L}_0, \mathcal{L}_1) \leq C(g)$.*

The Liouville vector field Z_λ that is defined by $i_{Z_\lambda}\omega = \lambda$ satisfies

$$L_{Z_\lambda}\omega = d\lambda = -\omega.$$

Hence its flow (φ_t^λ) is conformal symplectic with $(\varphi_t^\lambda)_*\omega = e^{-t}\omega$ and even exact conformal symplectic because it preserves the zero section (and then the zero Liouville class). We have seen in Lemma 3 that φ_t^λ alters the distance γ up to the scaling factor e^{-t} .

Observe also that this flow is a homothety the fiber direction: $\varphi_t^\lambda(p) = e^{-t}p$. Hence the image of the unit codisk bundle $D^*(g)$ by φ_t is the codisk bundle $D_{e^{-t}}^*(g)$ with radius e^{-t} .

Let us introduce the following notation for $K \subset T^*\mathbb{T}^n$.

$$\delta_g(K) = \min\{r \geq 0; K \subset D_r^*(g)\}.$$

Finally, we have that for every H-isotopic to the zero section submanifolds $\mathcal{L}, \mathcal{L}'$ of $T^*\mathbb{T}^n$,

$$(6) \quad \gamma(\mathcal{L}, \mathcal{L}') \leq 2C(g) \max\{\delta_g(\mathcal{L}), \delta_g(\mathcal{L}')\}.$$

If now \mathcal{L} and \mathcal{L}' are two distinct H-isotopic to the zero section submanifolds of $T^*\mathbb{T}^n$ and $f : T^*\mathbb{T}^n \hookrightarrow T^*\mathbb{T}^n$ is a CES diffeomorphism that is CH isotopic to $\text{Id}_{T^*\mathbb{T}^n}$, we deduce from Theorem 14 that

$$\begin{aligned} \text{either } & \gamma(f^n(\mathcal{L}), f^n(\mathcal{L}')) \xrightarrow{n \rightarrow +\infty} +\infty \\ \text{or } & \gamma(f^{-n}(\mathcal{L}), f^{-n}(\mathcal{L}')) \xrightarrow{n \rightarrow +\infty} +\infty. \end{aligned}$$

By (6), one of the two sets

$$\bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L}); \quad \bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L}').$$

is not relatively compact. We deduce that there is at most one \mathcal{L} H-isotopic to the zero section such that

$$\bigcup_{k \in \mathbb{Z}} f^k(\mathcal{L}) \text{ is relatively compact.}$$

When \mathcal{L} is H-isotopic to the zero-section, $f(\mathcal{L})$ is also H-isotopic to the zero-section because f is CH-isotopic to $\text{Id}_{T^*\mathbb{T}^n}$, see Corollary 3. Moreover, the orbits of \mathcal{L} and $f(\mathcal{L})$ coincide. This implies that $\mathcal{L} = f(\mathcal{L})$. \square

7. EXAMPLES

7.1. Mañé example. This example was introduced by Mañé in the Hamiltonian setting, [11]. It can be extended to the conformal symplectic setting. For every vector field X of a closed manifold \mathcal{Q} , it provides a conformal Hamiltonian Tonelli flow of $T^*\mathcal{Q}$ such that the zero section is invariant and the flow restricted to this zero section is conjugated to the flow of X .

Let \mathcal{Q} be a closed manifold endowed with a Riemannian metric, $T^*\mathcal{Q}$ is endowed with its tautological 1-form λ and the symplectic form $\omega = -d\lambda$. We denote by $\|\cdot\|$ the norm on the fibers of $T^*\mathcal{Q}$ that is dual to the Riemannian norm of \mathcal{Q} and by p_q a point of $T^*\mathcal{Q}$ above $q \in \mathcal{Q}$.

If X is a vector field on \mathcal{Q} , we denote by p_X the 1-form on \mathcal{Q} that is dual to X via the Riemannian scalar product. We define the Hamiltonian

$$H_X(p_q) = \frac{1}{2}\|p_q + p_X(q)\|^2 - \frac{1}{2}\|p_X(q)\|^2.$$

Since the zero-section $\mathcal{Z} = \{p = 0\}$ is contained in the zero-energy level and is Lagrangian, \mathcal{Z} is invariant by the Hamiltonian flow of H_X . The restriction to \mathcal{Z} of the vector field is dual via ω to the derivative of H in the fiber direction, so if we denote by $\sharp : T_q^*M \rightarrow T_qM$ the duality that is defined by the Riemannian metric, we have

$$\dot{q}|_{\mathcal{Z}} = \sharp(p + p_X(q))|_{\mathcal{Z}} = \sharp p_X(q) = X(q).$$

Hence on the zero-section, the vector field is X .


In the conformal Hamiltonian setting, we add α times the Liouville vector field to the Hamiltonian vector field X_H of H , for some $\alpha \in \mathbb{R}$. Since the Liouville vector field vanishes on \mathcal{Z} , the dynamics remains conjugate to X .

Remark 16. The global attractor may differ from the zero section. For example, X may have an attractive fixed point whose unstable manifold is not contained in the \mathcal{Z} , in which case the global attractor is not a submanifold either.

7.2. An example of a Tonelli Hamiltonian that has an invariant Lagrangian submanifold that is not a graph. The example we are about to describe is inspired by an example of Le Calvez [9].

Let $\beta > 0$ be a positive number and let $\alpha \in (\beta, 2\beta)$. On $T^*\mathbb{R} = \mathbb{R}^2$, let H be the quadratic Tonelli Hamiltonian

$$H(x, y) = y^2 - \beta xy.$$



Consider the sum of the Hamiltonian vector field of H and of α times the Liouville vector field $-y \partial_y$:

$$(7) \quad \begin{cases} \dot{x} = -\beta x + 2y \\ \dot{y} = (\beta - \alpha)y. \end{cases}$$

The matrix of this linear system is $\begin{pmatrix} -\beta & 2 \\ 0 & \beta - \alpha \end{pmatrix}$. Hence $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is an eigenvector for the eigenvalue $-\beta$ and $\begin{pmatrix} 1 \\ \beta - \frac{\alpha}{2} \end{pmatrix}$ is an eigenvector for the eigenvalue $\beta - \alpha$. As $\alpha \in (\beta, 2\beta)$, $(0, 0)$ is an attracting fixed point and the line $\mathbb{R} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is the strong stable eigenspace. Every solution that is not contained in an eigenspace is contained in a curve whose equation is

$$x = \frac{2}{2\beta - \alpha}y + K|y|^{\frac{\beta}{\alpha - \beta}}$$

where $K \neq 0$, and then is not a graph if $x(0) \cdot y(0) > 0$.

Let us choose two large real numbers $B > A > 0$ and let $V : \mathbb{R} \rightarrow [-1, 0]$ be a function with support in $[-B, B]$ such that $V|_{[-A, A]} = -1$, $V|_{[-B, -A]}$ is non-increasing and $V|_{[A, B]}$ is non-decreasing. Then we add $V(x)$ to $H(x, y)$ and the equations become

$$(8) \quad \begin{cases} \dot{x} = -\beta x + 2y \\ \dot{y} = -V'(x) + (\beta - \alpha)y. \end{cases}$$

As the support of V' is in $[-B, -A] \cup [A, B]$, the two vector fields are equal in the complement of $([-B, -A] \cup [A, B]) \times \mathbb{R}$. As $V'|_{[-B, -A]} \leq 0$, the orbit on the x -axis for $x \leq -B$ is pushed to the half plane $y > 0$ and then coincides with an orbit of (7) which tends to $(0, 0)$. In the same way, the orbit that coincides with the x -axis for $x \geq B$ tends to $(0, 0)$ at $+\infty$ with an incursion into the half-plane $y < 0$. Hence the union of these two orbits and $\{(0, 0)\}$ is an invariant curve Γ for (8) that is not a graph.

Now, let us choose $D > C > B$. Let $X : \mathbb{R} \rightarrow \mathbb{R}$ be a vector field such that

- $\forall x \in [-\frac{D+C}{2}, -B] \cup [B, \frac{C+D}{2}]$, $X(x) = -\beta x$;
- $X(-D) = X(D) = 0$ and all the derivatives of X are the same at $-D$ and D ;
- $(-D, -B]$ (resp. $[B, D)$) is a piece of unstable manifold of the equilibrium $-D$ (resp. D).

Then X defines also a vector field on the circle $\mathcal{C}_D = [-D, D]/D \sim -D$. Let H_X be the Hamiltonian that is associated to X on $T^*\mathbb{R} = \mathbb{R}^2$ via the Mañé construction

$$H_X(x, y) = \frac{1}{2}y(y + 2X(x)).$$

Let us eventually define

$$\begin{aligned} K(x, y) &= (1 - \eta(x))H_X(x, y) + \eta(x)(H(x, y) + V(x)) \\ &= \frac{1 - \eta(x)}{2}y(y + 2X(x)) + \eta(x)(y^2 - \beta xy + V(x)), \end{aligned}$$

where $\eta : \mathbb{R} \rightarrow [0, 1]$ is a bump function with support in $[-C, C]$ that is equal to 1 on $[-B, B]$. The function K is also Hamiltonian on the annulus $\mathcal{C}_D \times \mathbb{R}$ and, since

$$\frac{\partial^2 K}{\partial y^2}(x, y) = (1 - \eta(x)) + 2\eta(x) \geq 1,$$

K is Tonelli.

Note the following:

- $([-D, -B] \cup [B, D]) \times \{0\}$ is in the zero level of K and then is locally invariant by the Hamiltonian flow of K and also by the conformal Hamiltonian flow $(\frac{\partial K}{\partial y}, -\frac{\partial K}{\partial x} - \alpha y)$;
- $K|_{[-B, B] \times \mathbb{R}} = (H + V)|_{[-B, B] \times \mathbb{R}}$.

Finally, the vector field $(\frac{\partial K}{\partial y}, -\frac{\partial K}{\partial x} - \alpha y)$ has an invariant curve that is not a graph, which is the union of $([-D, -B] \cup [B, D]) \times \{0\}$ and the part of Γ that is between $x = -B$ and $x = B$.

APPENDIX A. YOMDIN'S INEQUALITY

Let \mathcal{L} be a compact Riemannian C^r manifold, $\mathcal{S} \subset \mathcal{L}$ be a compact C^r submanifold of dimension s and $f : \mathcal{L} \hookrightarrow \mathcal{L}$ be a C^r -diffeomorphism ($r \geq 1$). (The general statement does not require f to be invertible.)

Define the logarithmic volume growth of $f|_{\mathcal{S}}$ as

$$\log \text{vol}(f|_{\mathcal{S}}) = \limsup_{n \rightarrow +\infty} \frac{1}{n} \log |\text{vol}(f^n(\mathcal{S}))|,$$

where vol is the s -dimensional Riemannian volume, and

$$\text{rad}(Df) = \limsup_{n \rightarrow +\infty} \|Df^n\|_\infty^{1/n}, \quad \|Df\|_\infty = \sup_x \|Df_x\|.$$

Theorem 17 (Yomdin [22], Gromov [7]).

$$\log \text{vol}(f|_{\mathcal{S}}) \leq \text{ent}(f) + \log^+(\text{rad}(Df)^{s/r})$$

(where $\log^+ t = \max(0, \log t)$).

In particular, if \mathcal{L} and f are smooth,

$$\log \text{vol}(f|_{\mathcal{S}}) \leq \text{ent}(f|_{\mathcal{S}}) \leq \text{ent} f.$$

APPENDIX B. CONFORMAL DYNAMICS ARE EXACT

We assume that $(\mathcal{M}, \omega = -d\lambda)$ is an exact symplectic manifold. We prove that every conformal dynamics is symplectically conjugate to a CES dynamics.

Our first result explains that every conformal dynamics on an exact symplectic manifold is exact conformal with respect to some primitive of the symplectic form.

Proposition 9. *Let $f : \mathcal{M} \rightarrow \mathcal{M}$ be a (CS) diffeomorphism that is homotopic to $\text{Id}_{\mathcal{M}}$ and such that $f^*\omega = a\omega$. Then there exists a primitive λ_1 of $-\omega$, namely*

$$\lambda_1 = \frac{1}{1-a}(\lambda - f^*\lambda)$$

such that f is λ_1 CES. Hence is $j : \mathcal{S} \hookrightarrow \mathcal{M}$ is an isotropic embedding such that $j(\mathcal{S})$ is f invariant, $j(\mathcal{S})$ is λ_1 exact.

Proof of Proposition 9. We denote $\eta = f^*\lambda - a\lambda$. Then $d\eta = -f^*\omega + a\omega = 0$ and so η is closed. Observe that

$$\lambda - \frac{1}{1-a}\eta = \frac{1}{1-a}(\lambda - f^*\lambda) = \lambda_1,$$

so λ_1 is a primitive of $-\omega$.

We have

$$f^*\lambda_1 - a\lambda_1 = \eta - \frac{1}{1-a}(f^*\eta - a\eta).$$

Because f is homotopic to $\text{Id}_{\mathcal{M}}$, $f^*\eta - \eta$ is exact and

$$f^*\lambda_1 - a\lambda_1 = \frac{1}{1-a}(\eta - f^*\eta).$$

is exact. The conclusion comes from Proposition 8 for the 1-form λ_1 instead of λ . \square

Proposition 10. *Let $f : \mathcal{M} \looparrowright$ be a conformal symplectic diffeomorphism that is homotopic to $\text{Id}_{\mathcal{M}}$ and such that $f^*\omega = a\omega$ with $a > 0$ and $a \neq 1$. Then $\eta = f^*\lambda - a\lambda$ is a closed 1-form that is dual to a symplectic vector field Y such that $i_Y\omega = \eta$. When Y is complete, there exists a symplectically isotopic to $\text{Id}_{\mathcal{M}}$ diffeomorphism $g : \mathcal{M} \looparrowright$ such that $g^*\lambda - \lambda + \frac{1}{1-a}\eta$ is exact and then $g \circ f \circ g^{-1}$ is λ CES.*

Proof. We denote $\eta = f^*\lambda - a\lambda$. Then $d\eta = -f^*\omega + a\omega = 0$ and so η is closed. We denote by λ_1 the primitive of ω that was defined in Proposition 9.

Lemma 4. *There exists a symplectic vector field X with flow (g_t) such that $g_1^*\lambda - \lambda_1$ is exact.*

Proof. We consider the vector field X that is defined by $i_X\omega = \frac{1}{1-a}\eta$. As η is closed, X is symplectic. As Y is assumed to be complete and $X = \frac{1}{1-a}Y$, the vector field X is also complete and defines a flow (g_t) .

Then we have

$$L_X\lambda = -i_X\omega + d(i_X\lambda) = -\frac{1}{1-a}\eta + d(i_X\lambda).$$

If we denote by $[\cdot]$ the cohomology class, this gives

$$[L_X\lambda] = -\frac{1}{1-a}[\eta]$$

i.e.

$$\frac{d[g_t^*\lambda - \lambda]}{dt} = -\frac{1}{1-a}[g_t^*\eta].$$

We deduce that for all t we have $g_t^*\lambda - \lambda + \frac{t}{1-a}\eta$ is exact. In particular, $g_1^*\lambda - \lambda_1$ is exact. \square

We now consider $F = g_1 \circ f \circ g_1^{-1}$. We have

$$F^*\lambda = (g_1^{-1})^* \circ f^* \circ g_1^*(\lambda) = (g_1^{-1})^* \circ f^*(\lambda_1 + \nu_1)$$

where ν_1 is exact by lemma 4. By Proposition 9, $\nu_2 = f^*\lambda_1 - a\lambda_1$ is exact and we have

$$F^*\lambda = (g_1^{-1})^*(a\lambda_1 + \nu_2 + f^*\nu_1) = a\lambda + (g_1^{-1})^*(-a\nu_1 + \nu_2 + f^*\nu_1).$$

\square

Proposition 11. *Let X be a conformal symplectic vector field on \mathcal{M} such that $L_X\omega = \alpha\omega$ with $\alpha \in \mathbb{R}^*$. The 1-form $\xi = i_X\omega + \alpha\lambda$ is closed and the vector field X_1 defined by $i_{X_1}\omega = \xi$ is symplectic. When X_1 is complete, there exists a symplectically isotopic to $\text{Id}_{\mathcal{M}}$ diffeomorphism $g : \mathcal{M} \looparrowright$ such that $g^*\lambda - \lambda + \frac{1}{\alpha}\xi$ is exact. Then g^*X is λ conformal Hamiltonian.*

Proof. We have $d\xi = L_X\omega - \alpha\omega$ hence ξ is closed.

Lemma 5. *There exists a primitive λ_1 of $-\omega$, namely*

$$\lambda_1 = \lambda - \frac{1}{\alpha}\xi = -\frac{1}{\alpha}i_X\omega,$$

such that X is λ_1 Hamiltonian.

Proof. The 1-form

$$i_X\omega + \alpha\lambda_1 = i_X\omega + \alpha\lambda - \xi = 0$$

is exact. □

Lemma 6. *There exists a symplectic vector field Y with flow (ψ_t) such that $\psi_1^*\lambda - \lambda_1$ is exact.*

Proof. We consider the vector field Y that is defined by $i_Y\omega = \frac{1}{\alpha}\xi$. As ξ is closed, Y is symplectic. As X_1 is complete and $Y = \frac{1}{\alpha}X_1$, Y is also complete and defines a flow. Then we have

$$L_Y\lambda = -i_Y\omega + d(i_Y\lambda) = -\frac{1}{\alpha}\xi + d(i_Y\lambda).$$

We deduce that the flow (ψ_t) of Y satisfies

$$\frac{d}{dt}[\psi_t^*\lambda - \lambda] = -\frac{1}{\alpha}[\xi].$$

Hence $\psi_1^*\lambda - \lambda_1 = \psi_1^*\lambda - \lambda + \frac{1}{\alpha}\xi$ is exact. □

We denote $g = \psi_1$. Let us prove that g^*X is λ conformal Hamiltonian. Because g is symplectic, we have

$$i_{g^*X}\omega = g_*(i_X\omega) = g_*(\xi - \alpha\lambda).$$

Because $g^*\lambda - \lambda_1$ is exact, $g_*(\xi - \alpha\lambda) + \alpha\lambda$ is exact and $i_{g^*X}\omega + \alpha\lambda$ is exact and so g^*X is conformal Hamiltonian. □

REFERENCES

1. V.I. Arnold & A. Avez, Problèmes ergodiques de la mécanique classique. (French) Monographies Internationales de Mathématiques Modernes, No. 9 Gauthier-Villars, Editeur, Paris 1967 ii+243 pp.
2. R.C. Calleja, A. Celletti & R. de la Llave, Local behavior near quasi-periodic solutions of conformal symplectic systems. J. Dynam. Differential Equations 25 (2013), no. 3, 821–41.
3. C. Godbillon, Éléments de topologie algébrique (French) Hermann, Paris, 1971. 249 pp.
4. H. Geiges, Symplectic manifolds with disconnected boundary of contact type. Internat. Math. Res. Notices 1994, no. 1, 23–30.
5. H. Geiges, Examples of symplectic 4-manifolds with disconnected boundary of contact type. Bull. London Math. Soc. 27 (1995), no. 3, 278–280.
6. M. Gromov, Pseudoholomorphic curves in symplectic manifolds. Invent. Math. 81 (1985), 307–347.

7. M. Gromov, Entropy, homology and semialgebraic geometry. Séminaire Bourbaki, Vol. 1985/86. Astérisque No. 145-146 (1987), 5, 225–240.
 8. A. Katok & B. Hasselblatt, Introduction to the modern theory of dynamical systems. Encyclopedia of Mathematics and its Applications, 54. Cambridge University Press, Cambridge, 1995. xviii+802 pp.
 9. P. Le Calvez, Propriétés des attracteurs de Birkhoff. (French) Ergodic Theory Dynam. Systems 8 (1988), no. 2, 241–310.
 10. P. Libermann, Sur les automorphismes infinitésimaux des structures symplectiques et des structures de contact. (French) 1959 Colloque Géom. Diff. Globale (Bruxelles, 1958) pp. 37–59 Centre Belge Rech. Math., Louvain
 11. R. Mañé, On the minimizing measures of Lagrangian dynamical systems. Nonlinearity 5 (1992), no. 3, 623–638
 12. S. Marò & A. Sorrentino, Aubry-Mather theory for conformal symplectic systems. Comm. Math. Phys. 354 (2017), no. 2, 775–808.
 13. D. McDuff, Symplectic manifolds with contact type boundaries. Invent. Math. 103 (1991), no. 3, 651–671.
 14. K. F. Siburg, The principle of least action in geometry and dynamics. Lecture Notes in Mathematics, 1844. Springer-Verlag, Berlin, 2004. xii+128 pp
 15. E. Shelukhin, Symplectic cohomology and a conjecture of Viterbo arXiv:1904.06798
 16. J.-C. Sikorav, Sur les immersions lagrangiennes dans un fibré cotangent admettant une phase génératrice globale. C. R. Acad. Sci. Paris Sér. I Math. 302 (1986), no. 3, 119–122
 17. C. Viterbo, Symplectic topology as the geometry of generating functions. Math. Ann. 292 (1992), no. 4, 685–710.
 18. C. Viterbo, Symplectic topology and Hamilton-Jacobi equations. Morse theoretic methods in nonlinear analysis and in symplectic topology, 439–459, NATO Sci. Ser. II Math. Phys. Chem., 217, Springer, Dordrecht, 2006.
 19. C. Viterbo, Symplectic Homogenization, arXiv:0801.0206
 20. I. Vaisman, Locally conformal symplectic manifolds, *Internat. J. Math. Math. Sci.*, 8(3):521-536, 1985
 21. A. Weinstein, Lectures on symplectic manifolds. Expository lectures from the CBMS Regional Conference held at the University of North Carolina, March 8–12, 1976. Regional Conference Series in Mathematics, No. 29. American Mathematical Society, Providence, R.I., 1977
 22. Y. Yomdin, Volume growth and entropy. Israel J. Math. 57 (1987), no. 3, 285–300.
- Email address:* Marie-Claude.Arnaud@imj-prg.fr, jacques.fejoz@dauphine.fr