

Constructing and destructing tori on singular symplectic manifolds

Workshop on Classical Integrability and Perturbations

Eva Miranda (UPC-CEREMADE-IMCEE)

Journée Astronomie et Systèmes Dynamiques
Paris

- 1 Motivating examples
- 2 b^m -Symplectic manifolds
- 3 Integrable systems on b -symplectic manifolds
- 4 Debugging b^m -symplectic manifolds
- 5 Debugging integrable systems

The restricted 3-body problem

- Simplified version of the general 3-body problem. One of the bodies has **negligible mass**.
- The other two bodies move independently of it following **Kepler's laws** for the 2-body problem.




Figure: Circular 3-body problem


Planar restricted 3-body problem

- The time-dependent self-potential of the small body is
$$U(q, t) = \frac{1-\mu}{|q-q_1|} + \frac{\mu}{|q-q_2|},$$
with $q_1 = q_1(t)$ the position of the planet with mass $1 - \mu$ at time t and $q_2 = q_2(t)$ the position of the one with mass μ .
- The Hamiltonian of the system is
$$H(q, p, t) = p^2/2 - U(q, t), \quad (q, p) \in \mathbf{R}^2 \times \mathbf{R}^2,$$
where $p = \dot{q}$ is the momentum of the planet.
- Consider the canonical change $(X, Y, P_X, P_Y) \mapsto (r, \alpha, P_r =: y, P_\alpha =: G)$.
- Introduce **McGehee coordinates** (x, α, y, G) , where $r = \frac{2}{x^2}$, $x \in \mathbf{R}^+$, can be then extended to infinity ($x = 0$).
- The symplectic structure becomes a singular object
$$\omega = -\frac{4}{x^3} dx \wedge dy + d\alpha \wedge dG.$$
for $x > 0$
- The integrable 2-body problem for $\mu = 0$ is integrable with respect to the singular ω .


Model for these systems

$$\omega = \frac{1}{x_1^n} dx_1 \wedge dy_1 + \sum_{i \geq 2} dx_i \wedge dy_i$$

Close to $x_1 = 0$, the systems behave like,




and not like,



Other examples

- Kustaanheimo-Stiefel regularization for n -body problem \rightsquigarrow folded-type symplectic structures



- two fixed-center problem via Appell's transformation (Albouy) \rightsquigarrow combination of folded-type and b^m -symplectic structures \rightsquigarrow Dirac structures.

Can we get any singularity we want in physical examples?

Consider a system of two particles moving under the influence of the generalized potential $U(x) = -|x|^{-\alpha}$, $\alpha > 0$, with $|x|$ the distance.

Answer

The McGehee change of coordinates used to study collisions provides b^m -symplectic and m -folded symplectic forms for any m in the problem of a particle moving in a central force field with general potential depending on m .



Figure: The proof was somewhere on this greenboard

Definition

Let (M^{2n}, Π) be an (oriented) Poisson manifold such that the map

$$p \in M \mapsto (\Pi(p))^n \in \Lambda^{2n}(TM)$$

is transverse to the zero section, then $Z = \{p \in M \mid (\Pi(p))^n = 0\}$ is a hypersurface called *the critical hypersurface* and we say that Π is a **b -Poisson structure** on (M, Z) .

Symplectic foliation of a b -Poisson manifold

The symplectic foliation has dense symplectic leaves and codimension 2 symplectic leaves whose union is Z .

Darboux normal forms

Theorem (Guillemin-M.-Pires)

For all $p \in Z$, there exists a Darboux coordinate system $x_1, y_1, \dots, x_n, y_n$ centered at p such that Z is defined by $x_1 = 0$ and

$$\Pi = x_1 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y_1} + \sum_{i=2}^n \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_i}$$

Darboux for b^n -symplectic structures

$$\Pi = x_1^m \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial y_1} + \sum_{i=2}^n \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial y_i}$$


or dually

$$\omega = \frac{1}{x_1^n} dx_1 \wedge dy_1 + \sum_{i=2}^n dx_i \wedge dy_i$$

Dimension 2

Radko classified b-Poisson structures on compact oriented surfaces:

- **Geometrical invariants:** The topology of S and the curves γ_i where Π vanishes.
- **Dynamical invariants:** The periods of the “**modular vector field**” along γ_i .
- **Measure:** The regularized Liouville volume of S , $V_h^\varepsilon(\Pi) = \int_{|h|>\varepsilon} \omega_\Pi$ for h a function vanishing linearly on the curves $\gamma_1, \dots, \gamma_n$ and ω_Π the “dual” form to the Poisson structure.



Higher dimensions: Some compact examples.

- The product of (R, π_R) a Radko compact surface with a compact symplectic manifold (S, ω) is a b -Poisson manifold.
- corank 1 Poisson manifold (N, π) and X Poisson vector field $\Rightarrow (S^1 \times N, f(\theta) \frac{\partial}{\partial \theta} \wedge X + \pi)$ is a b -Poisson manifold if,
 - 1 f vanishes linearly.
 - 2 X is transverse to the symplectic leaves of N .

We then have as many copies of N as zeroes of f .

Poisson Geometry of the critical hypersurface


This last example is semilocally the *canonical* picture of a b -Poisson structure .

- 1 The critical hypersurface Z has an **induced regular Poisson** structure of corank 1.
- 2 There exists a **Poisson vector field** transverse to the symplectic foliation induced on Z .

The singular hypersurface

Theorem (Guillemin-M.-Pires)

If \mathcal{L} contains a compact leaf L , then Z is the mapping torus of the symplectomorphism $\phi : L \rightarrow L$ determined by the flow of a Poisson vector field v transverse to the symplectic foliation.




This description also works for b^m -symplectic structures.

A dual approach...

- b -Poisson structures can be seen as symplectic structures modeled over a Lie algebroid (the b -cotangent bundle).
- A vector field v is a **b -vector field** if $v_p \in T_p Z$ for all $p \in Z$. The **b -tangent bundle** bTM is defined by

$$\Gamma(U, {}^bTM) = \left\{ \begin{array}{l} \text{b-vector fields} \\ \text{on } (U, U \cap Z) \end{array} \right\}$$



- The **b -cotangent bundle** ${}^bT^*M$ is $({}^bTM)^*$. Sections of $\Lambda^p({}^bT^*M)$ are **b -forms**, ${}^b\Omega^p(M)$. The standard differential extends to

$$d : {}^b\Omega^p(M) \rightarrow {}^b\Omega^{p+1}(M)$$

- A **b -symplectic form** is a closed, nondegenerate, b -form of degree 2.
- This dual point of view, allows to prove a **b -Darboux theorem and semilocal forms** via an adaptation of Moser's path method because we can play the same tricks as in the symplectic case.

What else?

b -integrable systems

Definition

b -integrable system A set of b -functions^a f_1, \dots, f_n on (M^{2n}, ω) such that

- f_1, \dots, f_n Poisson commute.
- $df_1 \wedge \dots \wedge df_n \neq 0$ as a section of $\Lambda^n(bT^*(M))$ on a dense subset of M

$$^a c \log |x| + g$$

Example

The symplectic form $\frac{1}{h} dh \wedge d\theta$ defined on the interior of the upper hemisphere H_+ of S^2 extends to a b -symplectic form ω on the double of H_+ which is S^2 . The triple $(S^2, \omega, \log|h|)$ is a b -integrable system.

Example

If (f_1, \dots, f_n) is an integrable system on M , then $(\log|h|, f_1, \dots, f_n)$ on $H_+ \times M$ extends to a b -integrable on $S^2 \times M$.

Action-angle coordinates for b -integrable systems

The compact regular level sets of a b -integrable system are (Liouville) tori.

Theorem (Kiesenhofer-M.-Scott)

Around a Liouville torus there exist coordinates

$(p_1, \dots, p_n, \theta_1, \dots, \theta_n) : U \rightarrow B^n \times \mathbb{T}^n$ such that

$$\omega|_U = \frac{c}{p_1} dp_1 \wedge d\theta_1 + \sum_{i=2}^n dp_i \wedge d\theta_i, \quad (1)$$

and the level sets of the coordinates p_1, \dots, p_n correspond to the Liouville tori of the system.

Reformulation of the result

Integrable systems semilocally \longleftrightarrow twisted cotangent lift^a of a \mathbb{T}^n action by translations on itself to $(T^*\mathbb{T}^n)$.

^aWe replace the Liouville form by $\log |p_1| d\theta_1 + \sum_{i=2}^n p_i d\theta_i$.

Intermezzo on twisted b -cotangent lifts

Consider $G := S^1 \times \mathbf{R}^+ \times S^1$ acting on $M := S^1 \times \mathbf{R}^2$:

$(\varphi, a, \alpha) \cdot (\theta, x_1, x_2) := (\theta + \varphi, aR_\alpha(x_1, x_2))$, with R_α rotation.

Its twisted b -cotangent lift gives **focus-focus** singularities on b -symplectic manifolds.

The logarithmic Liouville one-form is $\lambda := \log |p| d\theta + y_1 dx_1 + y_2 dx_2$ and the moment map is $\mu := (f_1, f_2, f_3)$ with

$$f_1 = \langle \lambda, X_1^\# \rangle = \log |p|,$$

$$f_2 = \langle \lambda, X_2^\# \rangle = x_1 y_1 + x_2 y_2,$$

$$f_3 = \langle \lambda, X_3^\# \rangle = x_1 y_2 - y_1 x_2.$$



- 1 **Topology of the foliation.** In a neighbourhood of a compact connected fiber the b -integrable system F is diffeomorphic to the b -integrable system on $W := \mathbf{T}^n \times B^n$ given by the projections p_1, \dots, p_{n-1} and $\log |p_n|$.
- 2 **Uniformization of periods:** We want to define integrals whose $(b-)$ Hamiltonian vector fields induce a \mathbf{T}^n action. Start with \mathbf{R}^n -action:

$$\begin{aligned}\Phi &: \mathbf{R}^n \times (\mathbf{T}^n \times B^n) \rightarrow \mathbf{T}^n \times B^n \\ ((t_1, \dots, t_n), m) &\mapsto \Phi_{t_1}^{(1)} \circ \dots \circ \Phi_{t_n}^{(n)}(m).\end{aligned}$$

Uniformize to get a \mathbf{T}^n action with fundamental vector fields Y_i .

- 3 The vector fields Y_i are **Poisson vector fields** (check $\mathcal{L}_{Y_i} \mathcal{L}_{Y_i} \omega = 0$).
- 4 The vector fields Y_i are **Hamiltonian** with primitives $\sigma_1, \dots, \sigma_n \in {}^b C^\infty(W)$. In this step the properties of b -cohomology are essential. Use this action to drag a local normal form (**Darboux-Carathéodory**) in a whole neighbourhood.

- ① **Topology of the foliation.** In a neighbourhood of a compact connected fiber the b -integrable system F is diffeomorphic to the b -integrable system on $W := \mathbf{T}^n \times B^n$ given by the projections p_1, \dots, p_{n-1} and $\log |p_n|$.
- ② **Uniformization of periods:** We want to define integrals whose (b) -Hamiltonian vector fields induce a \mathbf{T}^n action. Start with \mathbf{R}^n -action:

$$\begin{aligned} \Phi &: \mathbf{R}^n \times (\mathbf{T}^n \times B^n) \rightarrow \mathbf{T}^n \times B^n \\ ((t_1, \dots, t_n), m) &\mapsto \Phi_{t_1}^{(1)} \circ \dots \circ \Phi_{t_n}^{(n)}(m). \end{aligned}$$

Uniformize to get a \mathbf{T}^n action with fundamental vector fields Y_i .

- ③ The vector fields Y_i are **Poisson vector fields** (check $\mathcal{L}_{Y_i} \mathcal{L}_{Y_i} \omega = 0$).
- ④ The vector fields Y_i are **Hamiltonian** with primitives $\sigma_1, \dots, \sigma_n \in {}^b C^\infty(W)$. In this step the properties of b -cohomology are essential. Use this action to drag a local normal form (**Darboux-Carathéodory**) in a whole neighbourhood.

- 1 **Topology of the foliation.** In a neighbourhood of a compact connected fiber the b -integrable system F is diffeomorphic to the b -integrable system on $W := \mathbf{T}^n \times B^n$ given by the projections p_1, \dots, p_{n-1} and $\log |p_n|$.
- 2 **Uniformization of periods:** We want to define integrals whose (b) -Hamiltonian vector fields induce a \mathbf{T}^n action. Start with \mathbf{R}^n -action:

$$\begin{aligned}\Phi &: \mathbf{R}^n \times (\mathbf{T}^n \times B^n) \rightarrow \mathbf{T}^n \times B^n \\ ((t_1, \dots, t_n), m) &\mapsto \Phi_{t_1}^{(1)} \circ \dots \circ \Phi_{t_n}^{(n)}(m).\end{aligned}$$

Uniformize to get a \mathbf{T}^n action with fundamental vector fields Y_i .

- 3 The vector fields Y_i are **Poisson vector fields** (check $\mathcal{L}_{Y_i} \mathcal{L}_{Y_i} \omega = 0$).
- 4 The vector fields Y_i are **Hamiltonian** with primitives $\sigma_1, \dots, \sigma_n \in {}^b C^\infty(W)$. In this step the properties of b -cohomology are essential. Use this action to drag a local normal form (**Darboux-Carathéodory**) in a whole neighbourhood.

- ① **Topology of the foliation.** In a neighbourhood of a compact connected fiber the b -integrable system F is diffeomorphic to the b -integrable system on $W := \mathbf{T}^n \times B^n$ given by the projections p_1, \dots, p_{n-1} and $\log |p_n|$.
- ② **Uniformization of periods:** We want to define integrals whose (b) -Hamiltonian vector fields induce a \mathbf{T}^n action. Start with \mathbf{R}^n -action:

$$\begin{aligned} \Phi &: \mathbf{R}^n \times (\mathbf{T}^n \times B^n) \rightarrow \mathbf{T}^n \times B^n \\ ((t_1, \dots, t_n), m) &\mapsto \Phi_{t_1}^{(1)} \circ \dots \circ \Phi_{t_n}^{(n)}(m). \end{aligned}$$

Uniformize to get a \mathbf{T}^n action with fundamental vector fields Y_i .

- ③ The vector fields Y_i are **Poisson vector fields** (check $\mathcal{L}_{Y_i} \mathcal{L}_{Y_i} \omega = 0$).
- ④ The vector fields Y_i are **Hamiltonian** with primitives $\sigma_1, \dots, \sigma_n \in {}^b C^\infty(W)$. In this step the properties of b -cohomology are essential. Use this action to drag a local normal form (**Darboux-Carathéodory**) in a whole neighbourhood.

A picture is worth more than a thousand words...




Figure: Fibration by Liouville tori

Applications to **KAM theory** (surviving torus under perturbations) on b -symplectic manifolds (Kiesenhofer-M.-Scott).

Theorem (Kiesenhofer-M.-Scott)

Consider $\mathbf{T}^n \times B_r^n$ with the standard b -symplectic structure and the b -function $H = k \log |y_1| + h(y)$ with h analytic. If the frequency map has a Diophantine value and is non-degenerate, then a Liouville torus on Z persists under sufficiently small perturbations of H . More precisely, if $|\epsilon|$ is sufficiently small, then the perturbed system


$$H_\epsilon = H + \epsilon P$$

(with $P(\varphi, y) = \log |y_1| + f_1(\tilde{\varphi}, y) + y_1 f_2(\varphi, y) + f_3(\varphi_1, y_1)$) admits an invariant torus \mathcal{T} .

Moreover, there exists a diffeomorphism $\mathbf{T}^n \rightarrow \mathcal{T}$ close to the identity taking the flow γ^t of the perturbed system on \mathcal{T} to the linear flow on \mathbf{T}^n with frequency vector $(\frac{k+\epsilon k'}{c}, \tilde{\omega})$.


Global classification


Circle actions on b -surfaces:



Further results


- Delzant theorem and a convexity theorem for \mathbb{T}^k -actions on b -symplectic manifolds (Guillemin- M.-Pires-Scott).
- What about b^m -symplectic manifolds? Guillemin-M.-Weitsman






Symplectic manifolds

- Darboux theorem
- Delzant and convexity theorems
- Action-Angle coordinates



b-Symplectic manifolds

- Darboux theorem
- Delzant and convexity theorems
- Action-Angle theorems



Folded symplectic manifolds

- Darboux theorem (Martinet)
- Delzant-type theorems (Cannas da Silva-Guillemin-Pires)
- ?

Examples and counterexamples

Orientable Surface

- Is symplectic
- Is folded symplectic
- (orientable or not) is b-symplectic

\mathbb{CP}^2

- Is symplectic
- Is folded symplectic
- Is **not** b-symplectic

S^4

- Is **not** symplectic
- Is **not** b-symplectic
- Is folded-symplectic

Desingularizing b^m -symplectic structures

Theorem (Guillemin-M.-Weitsman)


Given a b^m -symplectic structure ω on a compact manifold (M^{2n}, Z) :

- If $m = 2k$, there exists a family of **symplectic forms** ω_ϵ which coincide with the b^m -symplectic form ω outside an ϵ -neighbourhood of Z and for which the family of bivector fields $(\omega_\epsilon)^{-1}$ **converges** in the C^{2k-1} -topology to the Poisson structure ω^{-1} as $\epsilon \rightarrow 0$.
- If $m = 2k + 1$, there exists a family of **folded symplectic forms** ω_ϵ which coincide with the b^m -symplectic form ω outside an ϵ -neighbourhood of Z .

Deblogging b^{2k} -symplectic structures

$$\omega = \frac{dx}{x^{2k}} \wedge \left(\sum_{i=0}^{2k-1} \alpha_i x^i \right) + \beta \quad (2)$$

- Let $f \in \mathcal{C}^\infty(\mathbb{R})$ be an odd smooth function satisfying $f'(x) > 0$ for all $x \in [-1, 1]$,



and satisfying

$$f(x) = \begin{cases} \frac{-1}{(2k-1)x^{2k-1}} - 2 & \text{for } x < -1 \\ \frac{-1}{(2k-1)x^{2k-1}} + 2 & \text{for } x > 1 \end{cases}$$

Deblogging b^{2k} -symplectic structures

- Scaling:

$$f_{\epsilon}(x) := \frac{1}{\epsilon^{2k-1}} f\left(\frac{x}{\epsilon}\right). \quad (3)$$

Outside the interval $[-\epsilon, \epsilon]$,

$$f_{\epsilon}(x) = \begin{cases} \frac{-1}{(2k-1)x^{2k-1}} - \frac{2}{\epsilon^{2k-1}} & \text{for } x < -\epsilon \\ \frac{-1}{(2k-1)x^{2k-1}} + \frac{2}{\epsilon^{2k-1}} & \text{for } x > \epsilon \end{cases}$$

- Replace $\frac{dx}{x^{2k}}$ by df_{ϵ} to obtain

$$\omega_{\epsilon} = df_{\epsilon} \wedge \left(\sum_{i=0}^{2k-1} \alpha_i x^i \right) + \beta$$

which is symplectic.

Deblogging b^{2k} -symplectic structures

- Scaling:

$$f_\epsilon(x) := \frac{1}{\epsilon^{2k-1}} f\left(\frac{x}{\epsilon}\right). \quad (4)$$

Outside the interval $[-\epsilon, \epsilon]$,

$$f_\epsilon(x) = \begin{cases} \frac{-1}{(2k-1)x^{2k-1}} - \frac{2}{\epsilon^{2k-1}} & \text{for } x < -\epsilon \\ \frac{-1}{(2k-1)x^{2k-1}} + \frac{2}{\epsilon^{2k-1}} & \text{for } x > \epsilon \end{cases}$$

- Replace $\frac{dx}{x^{2k}}$ by df_ϵ to obtain

$$\omega_\epsilon = df_\epsilon \wedge \left(\sum_{i=0}^{2k-1} \alpha_i x^i \right) + \beta$$

which is symplectic.

New toy: Deblogging integrable systems

Denote $F_\epsilon^{m-i}(x) = (\frac{d}{dx} f_\epsilon(x))x^i$, and hence $F_\epsilon^i(x) = (\frac{d}{dx} f_\epsilon(x))x^{m-i}$.
The desingularized ω_ϵ reads

$$\omega_\epsilon = \sum_{i=0}^{m-1} F_\epsilon^{m-i}(x) dx \wedge \alpha_{m-i} + \beta.$$

Definition

The desingularization of a b^m -integrable system $\mu = (f_1, \dots, f_n)$ is given by:

$$\mu = (f_1 = c_0 \log(x) + \sum_{i=1}^{m-1} c_i \frac{1}{x^i}, \dots, f_n) \mapsto \mu_\epsilon = (f_{1\epsilon} = \sum_{i=1}^m \hat{c}_i G_\epsilon^i(x), f_2, \dots, f_n)$$

with $G_\epsilon^i(x) = \int_0^x F_\epsilon^i(\tau) d\tau$, and $\hat{c}_1 = c_0$ and $\hat{c}_{i-1} = -ic_i$ if $i \neq 0$.

limits

When ϵ tends to 0, μ_ϵ tends to μ .

Debugging everything...

