

Well-posedness in critical spaces for barotropic viscous fluids

Boris Haspot *

Abstract

This paper is dedicated to the study of viscous compressible barotropic fluids in dimension $N \geq 2$. We address the question of well-posedness for *large* data having critical Besov regularity. Our result improve the analysis of R. Danchin in [13], by the fact that we choose initial density more general in $B_{p,1}^{\frac{N}{p}}$ with $1 \leq p < +\infty$. Our result relies on a new a priori estimate for the velocity, where we introduce a new structure to *kill* the coupling between the density and the velocity. In particular our result is the first where we obtain uniqueness without imposing hypothesis on the gradient of the density.

1 Introduction

The motion of a general barotropic compressible fluid is described by the following system:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \operatorname{div}(\mu(\rho)D(u)) - \nabla(\lambda(\rho)\operatorname{div}u) + \nabla P(\rho) = \rho f, \\ (\rho, u)_{/t=0} = (\rho_0, u_0). \end{cases} \quad (1.1)$$

Here $u = u(t, x) \in \mathbb{R}^N$ stands for the velocity field and $\rho = \rho(t, x) \in \mathbb{R}^+$ is the density. The pressure P is a suitable smooth function of ρ . We denote by λ and μ the two viscosity coefficients of the fluid, which are assumed to satisfy $\mu > 0$ and $\lambda + 2\mu > 0$ (in the sequel to simplify the calculus we will assume the viscosity coefficients as constants). Such a conditions ensures ellipticity for the momentum equation and is satisfied in the physical cases where $\lambda + \frac{2\mu}{N} > 0$. We supplement the problem with initial condition (ρ_0, u_0) and an outer force f . Throughout the paper, we assume that the space variable $x \in \mathbb{R}^N$ or to the periodic box \mathcal{T}_a^N with period a_i , in the i -th direction. We restrict ourselves the case $N \geq 2$.

The problem of existence of global solution in time for Navier-Stokes equations was addressed in one dimension for smooth enough data by Kazhikov and Shelukin in [24], and for discontinuous ones, but still with densities away from zero, by Serre in [30] and Hoff in [17]. Those results have been generalized to higher dimension by Matsumura and Nishida in [26] for smooth data close to equilibrium and by Hoff in the case of discontinuous data in [20, 21]. All those results do not require to be far from the vacuum. The existence and uniqueness of local classical solutions for (1.1) with smooth initial data such that

*Karl Ruprecht Universität Heidelberg, Institut for Applied Mathematics, Im Neuenheimer Feld 294, D-69120 Heidelberg, Germany. Tel. 49(0)6221-54-6112

the density ρ_0 is bounded and bounded away from zero (i.e., $0 < \underline{\rho} \leq \rho_0 \leq M$) has been stated by Nash in [28]. Let us emphasize that no stability condition was required there. On the other hand, for small smooth perturbations of a stable equilibrium with constant positive density, global well-posedness has been proved in [26]. Many works on the case of the one dimension have been devoted to the qualitative behavior of solutions for large time (see for example [17, 24]). Refined functional analysis has been used for the last decades, ranging from Sobolev, Besov, Lorentz and Triebel spaces to describe the regularity and long time behavior of solutions to the compressible model [31], [32], [19], [23]. Let us recall that (local) existence and uniqueness for (1.1) in the case of smooth data with no vacuum has been stated for long in the pioneering works by J. Nash [28], and A. Matsumura, T. Nishida [26]. For results of weak-strong uniqueness, we refer to the work of P. Germain [14].

Guided in our approach by numerous works dedicated to the incompressible Navier-Stokes equation (see e.g [27]):

$$(NS) \quad \begin{cases} \partial_t v + v \cdot \nabla v - \mu \Delta v + \nabla \Pi = 0, \\ \operatorname{div} v = 0, \end{cases}$$

we aim at solving (1.1) in the case where the data (ρ_0, u_0, f) have *critical* regularity. By critical, we mean that we want to solve the system functional spaces with norm invariant by the changes of scales which leaves (1.1) invariant. In the case of barotropic fluids, it is easy to see that the transformations:

$$(\rho(t, x), u(t, x)) \longrightarrow (\rho(l^2 t, lx), lu(l^2 t, lx)), \quad l \in \mathbb{R}, \quad (1.2)$$

have that property, provided that the pressure term has been changed accordingly. The use of critical functional frameworks led to several new well-posedness results for compressible fluids (see [10, 11, 13]). In addition to have a norm invariant by (1.2), appropriate functional space for solving (1.1) must provide a control on the L^∞ norm of the density (in order to avoid vacuum and loss of ellipticity). For that reason, we restricted our study to the case where the initial data (ρ_0, u_0) and external force f are such that, for some positive constant $\bar{\rho}$:

$$(\rho_0 - \bar{\rho}) \in B_{p,1}^{\frac{N}{p}}, \quad u_0 \in B_{p_1,1}^{\frac{N}{p_1}-1} \quad \text{and} \quad f \in L_{loc}^1(\mathbb{R}^+, \in B_{p_1,1}^{\frac{N}{p_1}-1})$$

with $(p, p_1) \in [1, +\infty[$ good chosen. In [13], however, we had to have $p = p_1$, indeed in this article there exists a very strong coupling between the pressure and the velocity. To be more precise, the pressure term is considered as a term of rest for the elliptic operator in the momentum equation of (1.1). This paper improve the results of R. Danchin in [10, 13], in the sense that the initial density belongs to larger spaces $B_{p,1}^{\frac{N}{p}}$ with $p \in [1, +\infty[$. The main idea of this paper is to introduce a new variable than the velocity in the goal to *kill* the relation of coupling between the velocity and the density. In the present paper, we address the question of local well-posedness in the critical functional framework under the assumption that the initial density belongs to critical Besov space with a index of integrability different of this of the velocity. We adapt the spirit of the results of [1] and [16] which treat the case of Navier-Stokes incompressible with dependent density (at the

difference than in these works the velocity and the density are naturally decoupled). To simplify the notation, we assume from now on that $\bar{\rho} = 1$. Hence as long as ρ does not vanish, the equations for $(a = \rho^{-1} - 1, u)$ read:

$$\begin{cases} \partial_t a + u \cdot \nabla a = (1 + a) \operatorname{div} u, \\ \partial_t u + u \cdot \nabla u - (1 + a) \mathcal{A}u + \nabla(g(a)) = f, \end{cases} \quad (1.3)$$

In the sequel we will note $\mathcal{A} = \mu \Delta + (\lambda + \mu) \nabla \operatorname{div}$ and where g is a smooth function which may be computed from the pressure function P .

One can now state our main result.

Theorem 1.1 *Let P a suitably smooth function of the density and $1 \leq p_1 \leq p \leq 2N$ such that $\frac{1}{p_1} \leq \frac{1}{N} + \frac{1}{p}$ and $2\frac{N}{p} - 1 > 0$. Assume that $u_0 \in B_{p_1,1}^{\frac{N}{p_1}-1}$, $f \in L_{loc}^1(\mathbb{R}^+, B_{p_1,1}^{\frac{N}{p_1}-1})$ and $a_0 \in B_{p,1}^{\frac{N}{p}}$ with $1 + a_0$ bounded away from zero.*

If $\frac{1}{p} + \frac{1}{p_1} > \frac{1}{N}$ there exists a positive time T such that system (1.1) has a solution (a, u) with $1 + a$ bounded away from zero,

$$a \in \tilde{C}([0, T], B_{p,1}^{\frac{N}{p}}), \quad u \in \tilde{C}([0, T]; B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{L}^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2}).$$

Moreover this solution is unique if $\frac{2}{N} \leq \frac{1}{p} + \frac{1}{p_1}$.

Remark 1 *It seems possible to improve the theorem 1.1 by choosing initial data a_0 in $B_{p,\infty}^{\frac{N}{p}} \cap B_{\infty,1}^0$, however some supplementary conditions appear on p_1 in this case.*

The key to theorem 1.1 is to introduce a new variable v_1 to control the velocity where to avoid the coupling between the density and the velocity, we analyze by a new way the pressure term. More precisely we write the gradient of the pressure as a Laplacian of the variable v_1 , and we introduce this term in the linear part of the momentum equation. We have then a control on v_1 which can write roughly as $u - \mathcal{G}P(\rho)$ where \mathcal{G} is a pseudodifferential operator of order -1 . By this way, we have canceled the coupling between v_1 and the density, we next verify easily that we have a control Lipschitz of the gradient of u (it is crucial to estimate the density by the transport equation).

Remark 2 *In the present paper we did not strive for unnecessary generality which may hide the new ideas of our analysis. Hence we focused on the somewhat academic model of barotropic fluids. In physical contexts however, a coupling with the energy equation has to be introduced. Besides, the viscosity coefficients may depend on the density. We believe that our analysis may be carried out to these more general models. (See [15]).*

In [18], D. Hoff show a very strong theorem of uniqueness for the weak solution when the pressure is of the specific form $P(\rho) = K\rho$ with $K > 0$. Similarly in [20], [21], [19], D. Hoff get global weak solution with regularizing effects on the velocity. In particular when the pressure is on this form, he does not need to have estimate on the gradient of the density. In the following corollary, we will observe that this type of pressure assure a specific structure and avoid to impose that $p < 2N$.

Corollary 1 *Assume that $P(\rho) = K\rho$ with $K > 0$. Let $1 \leq p_1 \leq p \leq +\infty$ such that $\frac{1}{p_1} \leq \frac{1}{N} + \frac{1}{p}$. Assume that $u_0 \in B_{p_1,1}^{\frac{N}{p_1}-1}$, $f \in L_{loc}^1(\mathbb{R}^+, B_{p_1,1}^{\frac{N}{p_1}-1})$ and $a_0 \in B_{p,1}^{\frac{N}{p}}$ with $1 + a_0$ bounded away from zero. If $\frac{1}{p} + \frac{1}{p_1} > \frac{1}{N}$ there exists a positive time T such that system (1.1) has a solution (a, u) with $1 + a$ bounded away from zero,*

$$a \in \tilde{C}([0, T], B_{p,1}^{\frac{N}{p}}), \quad u \in \tilde{C}([0, T]; B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{L}^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1}).$$

If moreover we assume that $\sqrt{\rho_0}u_0 \in L^2$, $\rho_0 - \bar{\rho} \in L_2^1$, $u_0 \in L^\infty$ and $\lambda = 0$ then the solution (a, u) is unique.

Remark 3 *Here L_2^1 defines the corresponding Orlicz space.*

Remark 4 *Up to my knowledge, it seems that it is the first time that we get strong solution without condition of controll in space with positive regularity for the gradient of the density.*

Remark 5 *Moreover we can observe that with this type of pressure we are very close to have existence of strong in finite time solution for initial data (a_0, u_0) in $B_{\infty,1}^0 \times B_{N,1}^1$. It means that this theorem rely the result of D. Hoff where the initial density is assumed L^∞ but where the initial velocity is more regular and the results of R. Danchin in [13].*

Remark 6 *In particular we can show that the solution of D. Hoff in [19] are unique on a finite time interval $[0, T]$.*

The study of the linearization of (1.1) leads also the following continuation criterion:

Theorem 1.2 *Let $1 \leq p_1 \leq p \leq +\infty$ such that $\frac{N}{p_1} - 1 \leq \frac{N}{p}$ and $\frac{N}{p_1} - 1 + \frac{N}{p} > 0$. Assume that (1.1) has a solution $(a, u) \in C([0, T], B_{p,1}^{\frac{N}{p}} \times (B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})^N)$ with $p_1 > N$, $\rho_0^{\frac{1}{p_1}}u_0 \in L^{p_1}$ and:*

$$\lambda \leq \frac{4\mu}{N^2(p_1 - 1)}, \tag{1.4}$$

on the time interval $[0, T]$ which satisfies the following three conditions:

- *the function a belongs to $L^\infty(0, T; B_{p,1}^{\frac{N}{p}})$,*
- *the function $1 + a$ is bounded away from zero.*

Then (a, u) may be continued beyond T .

Remark 7 *Up my knowledge, it is the first time that we get a criterion of blow-up for strong solution for compressible Navier-Stokes system without imposing a controll Lipschitz of the norm ∇u .*

Our paper is structured as follows. In section , we give a few notation and briefly introduce the basic Fourier analysis techniques needed to prove our result. In section 3 and 4 are devoted to the proof of key estimates for the linearized system (1.1). In section 5, we prove the theorem 1.1 and corollary ?? whereas section 6 is devoted to the proof of continuation criterions of theorem ?? and 1.2. Two inescapable technical commutator estimates and some theorems of ellipticity are postponed in an appendix.

2 Littlewood-Paley theory and Besov spaces

Throughout the paper, C stands for a constant whose exact meaning depends on the context. The notation $A \lesssim B$ means that $A \leq CB$. For all Banach space X , we denote by $C([0, T], X)$ the set of continuous functions on $[0, T]$ with values in X . For $p \in [1, +\infty]$, the notation $L^p(0, T, X)$ or $L_T^p(X)$ stands for the set of measurable functions on $(0, T)$ with values in X such that $t \rightarrow \|f(t)\|_X$ belongs to $L^p(0, T)$. Littlewood-Paley decomposition corresponds to a dyadic decomposition of the space in Fourier variables. Let $\alpha > 1$ and (φ, χ) be a couple of smooth functions valued in $[0, 1]$, such that φ is supported in the shell supported in $\{\xi \in \mathbb{R}^N / \alpha^{-1} \leq |\xi| \leq 2\alpha\}$, χ is supported in the ball $\{\xi \in \mathbb{R}^N / |\xi| \leq \alpha\}$ such that:

$$\forall \xi \in \mathbb{R}^N, \quad \chi(\xi) + \sum_{l \in \mathbb{N}} \varphi(2^{-l}\xi) = 1.$$

Denoting $h = \mathcal{F}^{-1}\varphi$, we then define the dyadic blocks by:

$$\begin{aligned} \Delta_l u &= 0 \quad \text{if } l \leq -2, \\ \Delta_{-1} u &= \chi(D)u = \tilde{h} * u \quad \text{with } \tilde{h} = \mathcal{F}^{-1}\chi, \\ \Delta_l u &= \varphi(2^{-l}D)u = 2^{lN} \int_{\mathbb{R}^N} h(2^l y) u(x - y) dy \quad \text{with } h = \mathcal{F}^{-1}\chi, \quad \text{if } l \geq 0, \\ S_l u &= \sum_{k \leq l-1} \Delta_k u. \end{aligned}$$

Formally, one can write that: $u = \sum_{k \in \mathbb{Z}} \Delta_k u$. This decomposition is called nonhomogeneous Littlewood-Paley decomposition.

2.1 Nonhomogeneous Besov spaces and first properties

Definition 2.1 For $s \in \mathbb{R}$, $p \in [1, +\infty]$, $q \in [1, +\infty]$, and $u \in \mathcal{S}'(\mathbb{R}^N)$ we set:

$$\|u\|_{B_{p,q}^s} = \left(\sum_{l \in \mathbb{Z}} (2^{ls} \|\Delta_l u\|_{L^p})^q \right)^{\frac{1}{q}}.$$

The Besov space $B_{p,q}^s$ is the set of temperate distribution u such that $\|u\|_{B_{p,q}^s} < +\infty$.

Remark 8 The above definition is a natural generalization of the nonhomogeneous Sobolev and Hölder spaces: one can show that $B_{\infty,\infty}^s$ is the nonhomogeneous Hölder space C^s and that $B_{2,2}^s$ is the nonhomogeneous space H^s .

Proposition 2.1 The following properties holds:

1. there exists a constant universal C such that:
$$C^{-1} \|u\|_{B_{p,r}^s} \leq \|\nabla u\|_{B_{p,r}^{s-1}} \leq C \|u\|_{B_{p,r}^s}.$$
2. If $p_1 < p_2$ and $r_1 \leq r_2$ then $B_{p_1,r_1}^s \hookrightarrow B_{p_2,r_2}^{s-N(1/p_1-1/p_2)}$.
3. $B_{p,r_1}^{s'} \hookrightarrow B_{p,r}^s$ if $s' > s$ or if $s = s'$ and $r_1 \leq r$.

Before going further into the paraproduct for Besov spaces, let us state an important proposition.

Proposition 2.2 *Let $s \in \mathbb{R}$ and $1 \leq p, r \leq +\infty$. Let $(u_q)_{q \geq -1}$ be a sequence of functions such that*

$$\left(\sum_{q \geq -1} 2^{qs} \|u_q\|_{L^p}^r \right)^{\frac{1}{r}} < +\infty.$$

If $\text{supp } \hat{u}_1 \subset \mathcal{C}(0, 2^q R_1, 2^q R_2)$ for some $0 < R_1 < R_2$ then $u = \sum_{q \geq -1} u_q$ belongs to $B_{p,r}^s$ and there exists a universal constant C such that:

$$\|u\|_{B_{p,r}^s} \leq C^{1+|s|} \left(\sum_{q \geq -1} (2^{qs} \|u_q\|_{L^p})^r \right)^{\frac{1}{r}}.$$

Let now recall a few product laws in Besov spaces coming directly from the paradifferential calculus of J-M. Bony (see [4]) and rewrite on a generalized form in [1] by H. Abidi and M. Paicu (in this article the results are written in the case of homogeneous spaces but it can easily generalize for the nonhomogeneous Besov spaces).

Proposition 2.3 *We have the following laws of product:*

- For all $s \in \mathbb{R}$, $(p, r) \in [1, +\infty]^2$ we have:

$$\|uv\|_{B_{p,r}^s} \leq C(\|u\|_{L^\infty} \|v\|_{B_{p,r}^s} + \|v\|_{L^\infty} \|u\|_{B_{p,r}^s}). \quad (2.5)$$

- Let $(p, p_1, p_2, r, \lambda_1, \lambda_2) \in [1, +\infty]^2$ such that: $\frac{1}{p} \leq \frac{1}{p_1} + \frac{1}{p_2}$, $p_1 \leq \lambda_2$, $p_2 \leq \lambda_1$, $\frac{1}{p} \leq \frac{1}{p_1} + \frac{1}{\lambda_1}$ and $\frac{1}{p} \leq \frac{1}{p_2} + \frac{1}{\lambda_2}$. We have then the following inequalities: if $s_1 + s_2 + N \inf(0, 1 - \frac{1}{p_1} - \frac{1}{p_2}) > 0$, $s_1 + \frac{N}{\lambda_2} < \frac{N}{p_1}$ and $s_2 + \frac{N}{\lambda_1} < \frac{N}{p_2}$ then:

$$\|uv\|_{B_{p,r}^{s_1+s_2-N(\frac{1}{p_1}+\frac{1}{p_2}-\frac{1}{p})}} \lesssim \|u\|_{B_{p_1,r}^{s_1}} \|v\|_{B_{p_2,\infty}^{s_2}}, \quad (2.6)$$

when $s_1 + \frac{N}{\lambda_2} = \frac{N}{p_1}$ (resp $s_2 + \frac{N}{\lambda_1} = \frac{N}{p_2}$) we replace $\|u\|_{B_{p_1,r}^{s_1}} \|v\|_{B_{p_2,\infty}^{s_2}}$ (resp $\|v\|_{B_{p_2,\infty}^{s_2}}$) by $\|u\|_{B_{p_1,1}^{s_1}} \|v\|_{B_{p_2,r}^{s_2}}$ (resp $\|v\|_{B_{p_2,\infty}^{s_2} \cap L^\infty}$), if $s_1 + \frac{N}{\lambda_2} = \frac{N}{p_1}$ and $s_2 + \frac{N}{\lambda_1} = \frac{N}{p_2}$ we take $r = 1$.

If $s_1 + s_2 = 0$, $s_1 \in (\frac{N}{\lambda_1} - \frac{N}{p_2}, \frac{N}{p_1} - \frac{N}{\lambda_2}]$ and $\frac{1}{p_1} + \frac{1}{p_2} \leq 1$ then:

$$\|uv\|_{B_{p,\infty}^{-N(\frac{1}{p_1}+\frac{1}{p_2}-\frac{1}{p})}} \lesssim \|u\|_{B_{p_1,1}^{s_1}} \|v\|_{B_{p_2,\infty}^{s_2}}. \quad (2.7)$$

If $|s| < \frac{N}{p}$ for $p \geq 2$ and $-\frac{N}{p} < s < \frac{N}{p}$ else, we have:

$$\|uv\|_{B_{p,r}^s} \leq C \|u\|_{B_{p,r}^s} \|v\|_{B_{p,\infty}^{\frac{N}{p}} \cap L^\infty}. \quad (2.8)$$

Remark 9 *In the sequel p will be either p_1 or p_2 and in this case $\frac{1}{\lambda} = \frac{1}{p_1} - \frac{1}{p_2}$ if $p_1 \leq p_2$, resp $\frac{1}{\lambda} = \frac{1}{p_2} - \frac{1}{p_1}$ if $p_2 \leq p_1$.*

Corollary 2 *Let $r \in [1, +\infty]$, $1 \leq p \leq p_1 \leq +\infty$ and s such that:*

- $s \in (-\frac{N}{p_1}, \frac{N}{p_1})$ if $\frac{1}{p} + \frac{1}{p_1} \leq 1$,
- $s \in (-\frac{N}{p_1} + N(\frac{1}{p} + \frac{1}{p_1} - 1), \frac{N}{p_1})$ if $\frac{1}{p} + \frac{1}{p_1} > 1$,

then we have if $u \in B_{p,r}^s$ and $v \in B_{p_1,\infty}^{\frac{N}{p_1}} \cap L^\infty$:

$$\|uv\|_{B_{p,r}^s} \leq C \|u\|_{B_{p,r}^s} \|v\|_{B_{p_1,\infty}^{\frac{N}{p_1}} \cap L^\infty}.$$

The study of non stationary PDE's requires space of type $L^\rho(0, T, X)$ for appropriate Banach spaces X . In our case, we expect X to be a Besov space, so that it is natural to localize the equation through Littlewood-Paley decomposition. But, in doing so, we obtain bounds in spaces which are not type $L^\rho(0, T, X)$ (except if $r = p$). We are now going to define the spaces of Chemin-Lerner in which we will work, which are a refinement of the spaces $L_T^\rho(B_{p,r}^s)$.

Definition 2.2 Let $\rho \in [1, +\infty]$, $T \in [1, +\infty]$ and $s_1 \in \mathbb{R}$. We set:

$$\|u\|_{\tilde{L}_T^\rho(B_{p,r}^{s_1})} = \left(\sum_{l \in \mathbb{Z}} 2^{lrs_1} \|\Delta_l u(t)\|_{L^\rho(L^p)}^r \right)^{\frac{1}{r}}.$$

We then define the space $\tilde{L}_T^\rho(B_{p,r}^{s_1})$ as the set of temperate distribution u over $(0, T) \times \mathbb{R}^N$ such that $\|u\|_{\tilde{L}_T^\rho(B_{p,r}^{s_1})} < +\infty$.

We set $\tilde{C}_T(\tilde{B}_{p,r}^{s_1}) = \tilde{L}_T^\infty(\tilde{B}_{p,r}^{s_1}) \cap \mathcal{C}([0, T], B_{p,r}^{s_1})$. Let us emphasize that, according to Minkowski inequality, we have:

$$\|u\|_{\tilde{L}_T^\rho(B_{p,r}^{s_1})} \leq \|u\|_{L_T^\rho(B_{p,r}^{s_1})} \text{ if } r \geq \rho, \quad \|u\|_{\tilde{L}_T^\rho(B_{p,r}^{s_1})} \geq \|u\|_{L_T^\rho(B_{p,r}^{s_1})} \text{ if } r \leq \rho.$$

Remark 10 It is easy to generalize proposition 2.3, to $\tilde{L}_T^\rho(B_{p,r}^{s_1})$ spaces. The indices s_1 , p , r behave just as in the stationary case whereas the time exponent ρ behaves according to Hölder inequality.

Here we recall a result of interpolation which explains the link of the space $B_{p,1}^s$ with the space $B_{p,\infty}^s$, see [9].

Proposition 2.4 There exists a constant C such that for all $s \in \mathbb{R}$, $\varepsilon > 0$ and $1 \leq p < +\infty$,

$$\|u\|_{\tilde{L}_T^\rho(B_{p,1}^s)} \leq C \frac{1+\varepsilon}{\varepsilon} \|u\|_{\tilde{L}_T^\rho(B_{p,\infty}^s)} \left(1 + \log \frac{\|u\|_{\tilde{L}_T^\rho(B_{p,\infty}^{s+\varepsilon})}}{\|u\|_{\tilde{L}_T^\rho(B_{p,\infty}^s)}} \right).$$

Now we give some result on the behavior of the Besov spaces via some pseudodifferential operator (see [9]).

Definition 2.3 Let $m \in \mathbb{R}$. A smooth function $f : \mathbb{R}^N \rightarrow \mathbb{R}$ is said to be a \mathcal{S}^m multiplier if for all multi-index α , there exists a constant C_α such that:

$$\forall \xi \in \mathbb{R}^N, \quad |\partial^\alpha f(\xi)| \leq C_\alpha (1 + |\xi|)^{m-|\alpha|}.$$

Proposition 2.5 Let $m \in \mathbb{R}$ and f be a \mathcal{S}^m multiplier. Then for all $s \in \mathbb{R}$ and $1 \leq p, r \leq +\infty$ the operator $f(D)$ is continuous from $B_{p,r}^s$ to $B_{p,r}^{s-m}$.

3 Estimates for parabolic system with variable coefficients

Let us first state estimates for the following constant coefficient parabolic system:

$$\begin{cases} \partial_t u - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u = f, \\ u|_{t=0} = u_0. \end{cases} \quad (3.9)$$

Proposition 3.6 *Assume that $\mu \geq 0$ and that $\lambda + 2\mu \geq 0$. Then there exists a universal constant κ such that for all $s \in \mathbb{Z}$ and $T \in \mathbb{R}^+$,*

$$\begin{aligned} \|u\|_{\tilde{L}_T^\infty(B_{p_1,1}^s)} &\leq \|u_0\|_{B_{p_1,1}^s} + \|f\|_{L_T^1(B_{p_1,1}^s)}, \\ \kappa \nu \|u\|_{L_T^1(B_{p_1,1}^{s+2})} &\leq \sum_{l \in \mathbb{Z}} 2^{ls} (1 - e^{-\kappa \nu 2^{2l} T}) (\|\Delta_l u_0\|_{L^{p_1}} + \|\Delta_l f\|_{L_T^1(L^{p_1})}), \end{aligned}$$

with $\nu = \min(\mu, \lambda + 2\mu)$.

We now consider the following parabolic system which is obtained by linearizing the momentum equation:

$$\begin{cases} \partial_t u + v \cdot \nabla u + u \cdot \nabla w - b(\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u) = f + g, \\ u|_{t=0} = u_0. \end{cases} \quad (3.10)$$

Above u is the unknown function. We assume that $u_0 \in B_{p_1,1}^s$, $f \in L^1(0, T; B_{p_1,1}^s)$ and $g \in L^r(0, T; B_{q_1,1}^s)$, that v and w are time dependent vector-fields with coefficients in $L^1(0, T; B_{p,1}^{\frac{N}{p}+1})$, that b is bounded by below by a positive constant \underline{b} and b belongs to $L^\infty(0, T; B_{p,1}^{\frac{N}{p}})$ with $p \in [1, +\infty]$.

Proposition 3.7 *Let $g = 0$ and $\underline{\nu} = \underline{b} \min(\mu, \lambda + 2\mu)$ and $\bar{\nu} = \mu + |\lambda + \mu|$. Assume that $s \in (-\frac{N}{p}, \frac{N}{p}]$. Let $m \in \mathbb{Z}$ be such that $b_m = 1 + S_m a$ satisfies:*

$$\inf_{(t,x) \in [0,T] \times \mathbb{R}^N} b_m(t, x) \geq \frac{\underline{b}}{2}. \quad (3.11)$$

There exist three constants c, C and κ (with c, C , depending only on N and on s , and κ universal) such that if in addition we have:

$$\|1 - S_m a\|_{L^\infty(0,T; B_{p_1,1}^{\frac{N}{p_1}})} \leq c \frac{\underline{\nu}}{\bar{\nu}} \quad (3.12)$$

then setting:

$$V(t) = \int_0^t \|v\|_{B_{p,1}^{\frac{N}{p}+1}} d\tau, \quad W(t) = \int_0^t \|w\|_{B_{p,1}^{\frac{N}{p}+1}} d\tau, \quad \text{and} \quad Z_m(t) = 2^{2m} \bar{\nu}^2 \underline{\nu}^{-1} \int_0^t \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}}^2 d\tau,$$

We have for all $t \in [0, T]$,

$$\begin{aligned} \|u\|_{\tilde{L}^\infty((0,T) \times B_{p_1,1}^s)} + \kappa \underline{\nu} \|u\|_{\tilde{L}^1((0,T) \times B_{p_1,1}^{s+2})} &\leq e^{C(V+W+Z_m)(t)} (\|u_0\|_{B_{p_1,1}^s} \\ &\quad + \int_0^t e^{-C(V+W+Z_m)(\tau)} \|f(\tau)\|_{B_{p_1,1}^s} d\tau). \end{aligned}$$

Remark 11 Let us stress the fact that if $a \in \widetilde{L}^\infty((0, T) \times B_{p,1}^{\frac{N}{p}})$ then assumption (3.20) and (3.21) are satisfied for m large enough. This will be used in the proof of theorem 1.1. Indeed, according to Bernstein inequality, we have:

$$\|a - S_m a\|_{L^\infty((0,T) \times \mathbb{R}^N)} \leq \sum_{q \geq m} \|\Delta_q a\|_{L^\infty((0,T) \times \mathbb{R}^N)} \lesssim \sum_{q \geq m} 2^{q \frac{N}{p}} \|\Delta_q a\|_{L^\infty(L^p)}.$$

Because $a \in \widetilde{L}^\infty((0, T) \times B_{p,1}^{\frac{N}{p}})$, the right-hand side is the remainder of a convergent series hence tends to zero when m goes to infinity. For a similar reason, (3.21) is satisfied for m large enough.

Proof: Let us first rewrite (3) as follows:

$$\partial_t u + v \cdot \nabla u + u \cdot \nabla w - b_m(\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u) = f + E_m - u \cdot \nabla w, \quad (3.13)$$

with $E_m = (\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u)(\operatorname{Id} - S_m)a$. Note that, because $-\frac{N}{p} < s \leq \frac{N}{p}$, the error term E_m may be estimated by:

$$\|E_m\|_{B_{p_1,1}^s} \lesssim \|a - S_m a\|_{B_{p,1}^{\frac{N}{p}}} \|D^2 u\|_{B_{p_1,1}^s}. \quad (3.14)$$

and we have:

$$\|u \cdot \nabla w\|_{B_{p_1,1}^s} \lesssim \|\nabla w\|_{B_{p,1}^{\frac{N}{p}}} \|u\|_{B_{p_1,1}^s}. \quad (3.15)$$

Now applying Δ_q to equation (3.13) yields:

$$\begin{aligned} \frac{d}{dt} u_q + v \cdot \nabla u_q - \mu \operatorname{div}(b_m \nabla u_q) - (\lambda + \mu) \nabla(b_m \operatorname{div} u_q) &= f_q + E_{m,q} - \Delta_q(u \cdot \nabla w) \\ &+ R_q + \widetilde{R}_q. \end{aligned} \quad (3.16)$$

where we denote by $u_q = \Delta_q u$ and with:

$$\begin{aligned} R_q &= [v^j, \Delta_q] \partial_j u, \\ \widetilde{R}_q &= \mu(\Delta_q(b_m \Delta u) - \operatorname{div}(b_m \nabla u_q)) + (\lambda + \mu)(\Delta_q(b_m \nabla \operatorname{div} u) - \nabla(b_m \operatorname{div} u_q)). \end{aligned}$$

Next multiplying both sides by $|u_q|^{p_1-2} u_q$, and integrating by parts in the second, third and last term in the left-hand side, we get:

$$\begin{aligned} \frac{1}{p_1} \frac{d}{dt} \|u_q\|_{L^{p_1}}^{p_1} - \frac{1}{p_1} \int (|u_q|^{p_1} \operatorname{div} v + \mu \operatorname{div}(b_m \nabla u_q) |u_q|^{p_1-2} u_q + \xi \nabla(b_m \operatorname{div} u_q) |u_q|^{p_1-2} u_q) dx \\ \leq \|u_q\|_{L^{p_1}}^{p_1-1} (\|f_q\|_{L^{p_1}} + \|\Delta_q E_m\|_{L^{p_1}} + \|\Delta_q(u \cdot \nabla w)\|_{L^{p_1}} + \|R_q\|_{L^{p_1}} + \|\widetilde{R}_q\|_{L^{p_1}}). \end{aligned}$$

Hence denoting $\xi = \mu + \lambda$, $\nu = \min(\mu, \lambda + 2\mu)$ and using (3.20), lemma [A5] of [10] and Young's inequalities we get:

$$\begin{aligned} \frac{1}{p_1} \frac{d}{dt} \|u_q\|_{L^{p_1}}^{p_1} + \frac{\nu b(p_1 - 1)}{p_1^2} 2^{2q} \|u_q\|_{L^{p_1}}^{p_1} \leq \|u_q\|_{L^{p_1}}^{p_1-1} (\|f_q\|_{L^{p_1}} + \|E_{m,q}\|_{L^{p_1}} + \|\Delta_q(u \cdot \nabla w)\|_{L^{p_1}} \\ + \frac{1}{p_1} \|u_q\|_{L^{p_1}} \|\operatorname{div} u\|_{L^\infty} + \|R_q\|_{L^{p_1}} + \|\widetilde{R}_q\|_{L^{p_1}}), \end{aligned}$$

which leads, after time integration to:

$$\begin{aligned} \|u_q\|_{L^{p_1}} + \frac{\nu \underline{b}(p_1 - 1)}{p_1} 2^{2q} \int_0^t \|u_q\|_{L^{p_1}} d\tau &\leq \|\Delta_q u_0\|_{L^{p_1}} + \int_0^t (\|f_q\|_{L^{p_1}} + \|E_{m,q}\|_{L^{p_1}} \\ &+ \|\Delta_q(u \cdot \nabla w)\|_{L^{p_1}} + \frac{1}{p_1} \|u_q\|_{L^{p_1}} \|\operatorname{div} u\|_{L^\infty} + \|R_q\|_{L^{p_1}} + \|\tilde{R}_q\|_{L^{p_1}}) d\tau, \end{aligned} \quad (3.17)$$

where $\underline{\nu} = \underline{b}\nu$. For commutators R_q and \tilde{R}_q , we have the following estimates (see lemma 1 and 2 in the appendix)

$$\|R_q\|_{L^{p_1}} \lesssim c_q 2^{-qs} \|v\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} \|u\|_{B_{p_1,1}^s}, \quad (3.18)$$

$$\|\tilde{R}_q\|_{L^{p_1}} \lesssim c_q \bar{\nu} 2^{-qs} \|S_m a\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} \|Du\|_{B_{p_1,1}^s}, \quad (3.19)$$

where $(c_q)_{q \in \mathbb{Z}}$ is a positive sequence such that $\sum_{q \in \mathbb{Z}} c_q = 1$, and $\bar{\nu} = \mu + |\lambda + \mu|$. Note that, owing to Bernstein inequality, we have:

$$\|S_m a\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} \lesssim 2^m \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}}$$

Hence, plugging these latter estimates and (3.14), (3.15) in (3.17), then multiplying by 2^{qs} and summing up on $q \in \mathbb{Z}$, we discover that, for all $t \in [0, T]$:

$$\begin{aligned} \|u\|_{L_t^\infty(B_{p_1,1}^s)} + \frac{\nu \underline{b}(p_1 - 1)}{p} \|u\|_{L_t^1(B_{p_1,1}^{s+2})} &\leq \|u_0\|_{B_{p_1,1}^s} + \|f\|_{L_t^1(B_{p_1,1}^s)} + C \int_0^t (\|v\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} \\ &+ \|w\|_{B_{p_1,1}^{\frac{N}{p_1}+1}}) \|u\|_{B_{p_1,1}^s} d\tau + C \bar{\nu} \int_0^t (\|a - S_m a\|_{B_{p_1,1}^{\frac{N}{p_1}}} \|u\|_{B_{p_1,1}^{s+2}} + 2^m \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}} \|u\|_{B_{p_1,1}^{s+1}}) d\tau, \end{aligned}$$

for a constant C depending only on N and s . Let $X(t) = \|u\|_{L_t^\infty(B_{p_1,1}^s)} + \nu \underline{b} \|u\|_{L_t^1(B_{p_1,1}^{s+2})}$. Assuming that m has been chosen so large as to satisfy:

$$C \bar{\nu} \|a - S_m a\|_{L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}})} \leq \underline{\nu},$$

and using that by interpolation, we have:

$$C \bar{\nu} \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}} \|u\|_{B_{p_1,1}^{s+2}} \leq \kappa \underline{\nu} + \frac{C^2 \bar{\nu}^2 2^{2m}}{4\kappa \underline{\nu}} \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}}^2 \|u\|_{B_{p_1,1}^s},$$

we end up with:

$$X(t) \leq \|u_0\|_{B_{p_1,1}^s} + \|f\|_{L_t^1(B_{p_1,1}^s)} + C \int_0^t (\|v\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} + \|w\|_{B_{p_1,1}^{\frac{N}{p_1}+1}} + \frac{\bar{\nu}^2}{\underline{\nu}} 2^{2m} \|a\|_{B_{p_1,1}^{\frac{N}{p_1}}}^2) X d\tau$$

Grönwall lemma then leads to the desired inequality. \square

Remark 12 *The proof of the continuation criterion (theorem 1.1) relies on a better estimate which is available when $u = v = w$ and $s > 0$. In fact, by arguing as in the proof of the previous proposition and by making use of inequality (7.61) instead of (7.59), one can prove that under conditions (3.20) and (3.21), there exists constants C and κ such that:*

$$\forall t \in [0, T], \|u\|_{L_t^\infty(B_{p_1,1}^s)} + \kappa \underline{\nu} \|u\|_{L_t^1(B_{p_1,1}^{s+2})} \leq e^{C(U+Z_m)(t)} (\|u_0\|_{B_{p_1,1}^s} + \int_0^t e^{-C(U+Z_m)(\tau)} \|f(\tau)\|_{B_{p_1,1}^s} d\tau) \quad \text{with } U(t) = \int_0^t \|\nabla u\|_{L^\infty} d\tau.$$

In the following corollary, we generalize proposition 3.8 when $g \neq 0$ and $g \in \tilde{L}^r(B_{q_1,1}^{s'})$. Moreover here $u_0 = u_1 + u_2$ with $u_1 \in B_{p_1,1}^s$ and $u_2 \in B_{p_2,1}^{s'}$.

Corollary 3 *Let $\underline{\nu} = \underline{b} \min(\mu, \lambda + 2\mu)$ and $\bar{\nu} = \mu + |\lambda + \mu|$. Assume that $s, s' \in (-\frac{N}{p}, \frac{N}{p}]$. Let $m \in \mathbb{Z}$ be such that $b_m = 1 + S_m a$ satisfies:*

$$\inf_{(t,x) \in [0,T] \times \mathbb{R}^N} b_m(t,x) \geq \frac{\underline{b}}{2}. \quad (3.20)$$

There exist three constants c, C and κ (with c, C , depending only on N and on s , and κ universal) such that if in addition we have:

$$\|1 - S_m a\|_{L^\infty(0,T;B_{p,1}^{\frac{N}{p}})} \leq c \frac{\underline{\nu}}{\bar{\nu}} \quad (3.21)$$

then setting:

$$V(t) = \int_0^t \|v\|_{B_{p,1}^{\frac{N}{p}+1}} d\tau, \quad W(t) = \int_0^t \|w\|_{B_{p,1}^{\frac{N}{p}+1}} d\tau, \quad \text{and } Z_m(t) = 2^{2m} \bar{\nu}^2 \underline{\nu}^{-1} \int_0^t \|a\|_{B_{p,1}^{\frac{N}{p}}}^2 d\tau,$$

We have for all $t \in [0, T]$,

$$\|u\|_{\tilde{L}_T^\infty(B_{p_1,1}^s + B_{p_2,1}^{s'})} + \kappa \underline{\nu} \|u\|_{\tilde{L}_T^1(B_{p_1,1}^{s+2} + B_{p_2,1}^{s'+2})} \leq e^{C(V+W+Z_m)(t)} (\|u_1\|_{B_{p_1,1}^s} + \|u_2\|_{B_{p_2,1}^{s'}} + \int_0^t e^{-C(V+W+Z_m)(\tau)} (\|f(\tau)\|_{B_{p_1,1}^s} + \|g(\tau)\|_{B_{q_1,1}^{s'}}) d\tau).$$

Proof: We split the solution u in two parts u_1 and u_2 which verify the following equations:

$$\begin{cases} \partial_t u_1 + v \cdot \nabla u_1 + u_1 \cdot \nabla w - b(\mu \Delta u_1 + (\lambda + \mu) \nabla \operatorname{div} u_1) = f, \\ u_{/t=0} = u_1^0, \end{cases}$$

and:

$$\begin{cases} \partial_t u_2 + v \cdot \nabla u_2 + u_2 \cdot \nabla w - b(\mu \Delta u_2 + (\lambda + \mu) \nabla \operatorname{div} u_2) = g, \\ u_{/t=0} = u_2^0. \end{cases}$$

We have then $u = u_1 + u_2$ and we conclude by applying proposition 3.7. \square
Proposition 3.7 fails in the limit case $s = -\frac{N}{p}$. The reason why is that proposition 2.3 cannot be applied any longer. One can however state the following result which will be the key to the proof of uniqueness in dimension two.

Proposition 3.8 *Under condition (3.20), there exists three constants c , C and κ (with c , C , depending only on N , and κ universal) such that if:*

$$\|a - S_m a\|_{\tilde{L}_t^\infty(B_{p,1}^{\frac{N}{p}})} \leq c \frac{\underline{\nu}}{\bar{\nu}}, \quad (3.22)$$

then we have:

$$\|u\|_{L_t^\infty(B_{p_1,\infty}^{-\frac{N}{p_1}})} + \kappa \underline{\nu} \|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{2-\frac{N}{p_1}})} \leq 2e^{C(V+W)(t)} (\|u_0\|_{B_{p_1,\infty}^{-\frac{N}{p_1}}} + \|f\|_{\tilde{L}_t^1(B_{p_1,\infty}^{\frac{N}{p_1}})}),$$

whenever $t \in [0, T]$ satisfies:

$$\bar{\nu}^2 t \|a\|_{\tilde{L}_t^\infty(B_{p,1}^{\frac{N}{p}})}^2 \leq c 2^{-2m} \underline{\nu}. \quad (3.23)$$

Proof: We just point out the changes that have to be done compare to the proof of proposition 3.7. The first one is that instead of (3.14) and (3.15), we have in accordance with proposition 2.3:

$$\|E_m\|_{\tilde{L}_t^1(B_{p_1,\infty}^{-\frac{N}{p_1}})} \lesssim \|a - S_m a\|_{\tilde{L}_t^\infty(B_{p,1}^{\frac{N}{p}})} \|D^2 u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{-\frac{N}{p_1}})}, \quad (3.24)$$

$$\|u \cdot w\|_{B_{p,\infty}^{-\frac{N}{p}}} \lesssim \|u\|_{B_{p_1,\infty}^{-\frac{N}{p_1}}} \|\nabla w\|_{B_{p,1}^{\frac{N}{p}}}. \quad (3.25)$$

The second change concerns the estimates of commutator R_q and \tilde{R}_q . According to inequality (7.60) and remark 13, we now have for all $q \in \mathbb{Z}$:

$$\|R_q\|_{L^p} \lesssim 2^{q\frac{N}{p_1}} \|v\|_{B_{p,1}^{\frac{N}{p}+1}} \|u\|_{B_{p_1,\infty}^{-\frac{N}{p_1}}}, \quad (3.26)$$

$$\|\tilde{R}_q\| \lesssim \bar{\nu} 2^{q\frac{N}{p_1}} \|S_m a\|_{\tilde{L}_t^\infty(B_{p,1}^{\frac{N}{p}+1})} \|Du\|_{\tilde{L}_t^1(B_{p_1,\infty}^{-\frac{N}{p_1}})}. \quad (3.27)$$

Plugging all these estimates in (3.17) then taking the supremum over $q \in \mathbb{Z}$, we get:

$$\begin{aligned} & \|u\|_{L_t^\infty(B_{p_1,\infty}^{-\frac{N}{p_1}})} + 2\underline{\nu} \|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{2-\frac{N}{p_1}})} \leq \|u_0\|_{B_{p_1,\infty}^{-\frac{N}{p_1}}} + C_i n t_0^t (\|v\|_{B_{p,1}^{\frac{N}{p}+1}} + \|w\|_{B_{p,1}^{\frac{N}{p}+1}}) \|u\|_{B_{p_1,\infty}^{-\frac{N}{p_1}}} d\tau \\ & + C\bar{\nu} (\|a - S_m a\|_{\tilde{L}_t^\infty(B_{p,1}^{\frac{N}{p}})} \|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{2-\frac{N}{p_1}})} + 2^m \|a\|_{L_t^\infty(B_{p,1}^{\frac{N}{p}})} \|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{1-\frac{N}{p_1}})} + \|f\|_{\tilde{L}_t^1(B_{p_1,\infty}^{-\frac{N}{p_1}})}). \end{aligned}$$

Using that:

$$\|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{1-\frac{N}{p_1}})} \leq \sqrt{t} \|u\|_{\tilde{L}_t^1(B_{p_1,\infty}^{2-\frac{N}{p_1}})}^{\frac{1}{2}} \|u\|_{L_t^\infty(B_{p_1,\infty}^{\frac{N}{p_1}})}^{\frac{1}{2}},$$

and taking advantage of assumption (3.22) and (3.23), it is now easy to complete the proof. \square

4 The mass conservation equation

Let us first recall standard estimates in Besov spaces for the following linear transport equation:

$$(\mathcal{H}) \quad \begin{cases} \partial_t a + u \cdot \nabla a = g, \\ a|_{t=0} = a_0. \end{cases}$$

Proposition 4.9 *Let $1 \leq p_1 \leq p \leq +\infty$, $r \in [1, +\infty]$ and $s \in \mathbb{R}$ be such that:*

$$-N \min\left(\frac{1}{p_1}, \frac{1}{p'}\right) < s < 1 + \frac{N}{p_1}.$$

There exists a constant C depending only on N , p , p_1 , r and s such that for all $a \in L^\infty([0, T], B_{p,r}^s)$ of (\mathcal{H}) with initial data a_0 in $B_{p,r}^s$ and $g \in L^1([0, T], B_{p,r}^s)$, we have for a.e $t \in [0, T]$:

$$\|f\|_{\tilde{L}_t^\infty(B_{p,r}^s)} \leq e^{CU(t)} \left(\|f_0\|_{B_{p,r}^s} + \int_0^t e^{-CV(\tau)} \|F(\tau)\|_{B_{p_1,r}^s} d\tau \right), \quad (4.28)$$

$$\text{with: } U(t) = \int_0^t \|\nabla u(\tau)\|_{B_{p_1,\infty}^{\frac{N}{p_1}} \cap L^\infty} d\tau.$$

For the proof of proposition 4.9, see [3]. We now focus on the mass equation associated to (1.3):

$$\begin{cases} \partial_t a + v \cdot \nabla a = (1+a)\operatorname{div} v, \\ a|_{t=0} = a_0. \end{cases} \quad (4.29)$$

Here we generalize a proof of R. Danchin in [13].

Proposition 4.10 *Let $r \in 1, +\infty$, $1 \leq p_1 \leq p \leq +\infty$ and $s \in (-\min(\frac{N}{p_1}, \frac{N}{p}, \frac{N}{p}])$. Assume that $a_0 \in B_{p,r}^s \cap L^\infty$, $v \in L^1(0, T; B_{p_1,1}^{\frac{N}{p_1}+1})$ and that $a \in \tilde{L}_T^\infty(B_{p,r}^s) \cap L_T^\infty$ satisfies (4.29). Let $V(t) = \int_0^t \|\nabla v(\tau)\|_{B_{p_1,1}^{\frac{N}{p_1}}} d\tau$. There exists a constant C depending only on N such that for all $t \in [0, T]$ and $m \in \mathbb{Z}$, we have:*

$$\|a\|_{\tilde{L}_t^\infty(B_{p,r}^s \cap L^\infty)} \leq e^{2CV(t)} \|a_0\|_{B_{p,r}^s \cap L^\infty} + e^{2CV(t)} - 1, \quad (4.30)$$

$$\|a - S_m a\|_{B_{p,r}^s} \leq \|a_0 - S_m a_0\|_{B_{p,r}^s} + \frac{1}{2} (1 + \|a_0\|_{B_{p,r}^s \cap L^\infty}) (e^{2CV(t)} - 1) + C \|a\|_{L^\infty} V(t), \quad (4.31)$$

$$\begin{aligned} \left(\sum_{l \leq m} 2^{lrs} \|\Delta_l(a - a_0)\|_{L_t^\infty(L^p)}^r \right)^{\frac{1}{r}} &\leq (1 + \|a_0\|_{B_{p,r}^s}) (e^{CV(t)} - 1) \\ &+ C 2^m \|a_0\|_{B_{p,r}^s} \int_0^t \|v\|_{B_{p_1,1}^{\frac{N}{p_1}}} d\tau. \end{aligned} \quad (4.32)$$

Proof: Applying Δ_l to (4.29) yields:

$$\partial_t \Delta_l a + v \cdot \nabla \Delta_l a = R_l + \Delta_l((1+a)\operatorname{div} v) \quad \text{with } R_l = [v \cdot \nabla, \Delta_l]a.$$

Multiplying by $\Delta_l a |\Delta_l a|^{p-2}$ then performing a time integration, we easily get:

$$\|\Delta_l a(t)\|_{L^p} \lesssim \|\Delta_l a_0\|_{L^p} + \int_0^t (\|R_l\|_{L^p} + \|\operatorname{div} v\|_{L^\infty} \|\Delta_l a\|_{L^p} + \|\Delta_l((1+a)\operatorname{div} v)\|_{L^p}) d\tau.$$

According to proposition 2.3 and interpolation, there exists a constant C and a positive sequence $(c_l)_{l \in \mathbb{N}}$ in l^r with norm 1 such that:

$$\|\Delta_l((1+a)\operatorname{div} v)\|_{L^p} \leq C c_l 2^{-ls} (1 + \|a\|_{B_{p,r}^s \cap L^\infty}) \|\operatorname{div} v\|_{B_{p_1,1}^{\frac{N}{p_1}}}.$$

Next the term $\|R_l\|_{L^p}$ may be bounded according to lemma 1 in appendix. We end up with:

$$\forall t \in [0, T], \forall l \in \mathbb{Z}, \quad 2^{ls} \|\Delta_l a(t)\|_{L^p} \leq 2^{ls} \|\Delta_l a_0\|_{L^p} + C \int_0^t c_l (1 + \|a\|_{B_{p,r}^s \cap L^\infty}) V' d\tau, \quad (4.33)$$

hence, summing up on \mathbb{Z} in l^r ,

$$\forall t \in [0, T], \forall l \in \mathbb{Z}, \quad \|a(t)\|_{B_{p,r}^s} \leq \|a_0\|_{B_{p,r}^s} + \int_0^t C V' \|a(\tau)\|_{B_{p,r}^s} d\tau + \int_0^t C (1 + \|a\|_{L_T^\infty}) V' d\tau.$$

Next we have:

$$\|a\|_{L_t^\infty} \leq \int_0^t (1 + \|a(\tau)\|_{L^\infty}) V'(\tau) d\tau.$$

By summing the two previous inequalities, applying Gronwall lemma and proposition 2.2 yields inequality (4.30). Let us now prove inequality (4.31). Starting from (4.33) and summing up over $l \geq m$ in l^r , we get:

$$\begin{aligned} \left(\sum_{l \geq m} 2^{lsr} \|\Delta_l a\|_{L_t^\infty(L^p)}^r \right)^{\frac{1}{r}} &\leq \left(\sum_{l \geq m} 2^{lsr} \|\Delta_l a_0\|_{L^p}^r \right)^{\frac{1}{r}} + C \int_0^t V' (e^{2CV} \|a_0\|_{B_{p,r}^s \cap L^\infty} + e^{2CV} - 1) d\tau \\ &\quad + \int_0^t C (1 + \|a\|_{L^\infty}) V' d\tau. \end{aligned}$$

Straightforward calculations then leads to (4.31). In order to prove (4.32), we use the fact that $\tilde{a} = a - a_0$ satisfies:

$$\partial_t \tilde{a} + v \cdot \nabla \tilde{a} = (1 + \tilde{a})\operatorname{div} v + a_0 \operatorname{div} v - v \cdot \nabla a_0, \quad \tilde{a}|_{t=0} = 0.$$

Therefore, arguing as for proving (4.33), we get for all $t \in [0, T]$ and $l \in \mathbb{Z}$,

$$\begin{aligned} 2^{l \frac{N}{p}} \|\Delta_l \tilde{a}\|_{L^p} &\leq \int_0^t 2^{l \frac{N}{p}} (\|\Delta_l(a_0 \operatorname{div} v)\|_{L^p} + \|\Delta_l(v \cdot \nabla a_0)\|_{L^p}) d\tau \\ &\quad + C \int_0^t c_l (1 + \|a\|_{B_{p,1}^{\frac{N}{p}}}) V' d\tau. \end{aligned}$$

Since $B_{p,1}^{\frac{N}{p}}$ is an algebra and the product maps $B_{p,1}^{\frac{N}{p}} \times B_{p,1}^{\frac{N}{p}-1}$ in $B_{p,1}^{\frac{N}{p}-1}$, we discover that:

$$2^{l\frac{N}{p}} \|\Delta_l \tilde{a}\|_{L^\infty(L^p)} \leq C \left(\int_0^t 2^l c_l \|a_0\|_{B_{p,1}^{\frac{N}{p}}} \|v\|_{B_{p,1}^{\frac{N}{p}}} d\tau + \int_0^t c_l (1 + \|a_0\|_{B_{p,1}^{\frac{N}{p}}} + \|a\|_{B_{p,1}^{\frac{N}{p}}}) V' d\tau \right),$$

hence, summing up on $l \leq m$,

$$\sum_{l \leq m} 2^{l\frac{N}{p}} \|\Delta_l \tilde{a}\|_{L^\infty(L^p)} \leq C \left(\int_0^t 2^m \|a_0\|_{B_{p,1}^{\frac{N}{p}}} \|v\|_{B_{p,1}^{\frac{N}{p}}} d\tau + \int_0^t (1 + \|a_0\|_{B_{p,1}^{\frac{N}{p}}} + \|a\|_{B_{p,1}^{\frac{N}{p}}}) V' d\tau \right),$$

Plugging (4.30) in the right-hand side yields (4.32).

5 The proof of theorem 1.1

5.1 Strategy of the proof

To improve the results of R. Danchin in [10], [13], it is crucial to kill the coupling between the velocity and the pressure which intervene in the works of R. Danchin. In this goal, we need to integrate the pressure term in the study of the linearized equation of the momentum equation. For making, we will try to express the gradient of the pressure as a Laplacian term, so we set for $\bar{\rho} > 0$ a constant state:

$$\operatorname{div} v = P(\rho) - P(\bar{\rho}).$$

Let \mathcal{E} the fundamental solution of the Laplace operator.

We will set in the sequel: $v = \nabla \mathcal{E} * (P(\rho) - P(\bar{\rho})) = \nabla (\mathcal{E} * [P(\rho) - P(\bar{\rho})])$ (* here means the operator of convolution). We verify next that:

$$\nabla \operatorname{div} v = \nabla \Delta (\mathcal{E} * [P(\rho) - P(\bar{\rho})]) = \Delta \nabla (\mathcal{E} * [P(\rho) - P(\bar{\rho})]) = \Delta v = \nabla P(\rho).$$

By this way we can now rewrite the momentum equation of (1.3). We obtain the following equation where we have set $\nu = 2\mu + \lambda$:

$$\partial_t u + u \cdot \nabla u - \frac{\mu}{\rho} \Delta \left(u - \frac{1}{\nu} v \right) - \frac{\lambda + \mu}{\rho} \nabla \operatorname{div} \left(u - \frac{1}{\nu} v \right) = f.$$

We want now calculate $\partial_t v$, by the transport equation we get:

$$\partial_t v = \nabla \mathcal{E} * \partial_t P(\rho) = -\nabla \mathcal{E} * (P'(\rho) \operatorname{div}(\rho u)).$$

We have finally:

$$\Delta (\partial_t v) = -P'(\rho) \operatorname{div}(\rho u).$$

Notation 1 *To simplify the notation, we will note in the sequel*

$$\nabla \mathcal{E} * (P'(\rho) \operatorname{div}(\rho u)) = \nabla (\Delta)^{-1} (P'(\rho) \operatorname{div}(\rho u)).$$

Finally we can now rewrite the system (1.3) as follows:

$$\begin{cases} \partial_t a + (v_1 + \frac{1}{\nu}v) \cdot \nabla a = (1+a)\operatorname{div}(v_1 + \frac{1}{\nu}v), \\ \partial_t v_1 - (1+a)\mathcal{A}v_1 = f - u \cdot \nabla u + \frac{1}{\nu}\nabla(\Delta)^{-1}(P'(\rho)\operatorname{div}(\rho u)), \\ a|_{t=0} = a_0, (v_1)|_{t=0} = (v_1)_0. \end{cases} \quad (5.34)$$

where $v_1 = u - \frac{1}{\nu}v$. In the sequel we will study this system by extracting some uniform bounds in Besov spaces on (a, v_1) as the in the following works [1], [16]. The advantage of the system (5.34) is that we have *kill* the coupling between v_1 and a term of pressure. Indeed in the works of R. Danchin [10], [13], the pressure was considered as a term of rest in the momentum equation, so it implied a strong relationship between the density and the velocity. In particular it was impossible to distinguish the index of integration for the Besov spaces.

5.2 Proof of the existence

Construction of approximate solutions

We use a standard scheme:

1. We smooth out the data and get a sequence of smooth solutions $(a^n, u^n)_{n \in \mathbb{N}}$ to (1.3) on a bounded interval $[0, T^n]$ which may depend on n . We set $v_1^n = u^n - v^n$ where $\operatorname{div} v^n = P(\rho^n) - P(\bar{\rho})$.
2. We exhibit a positive lower bound T for T^n , and prove uniform estimates on (a^n, u^n) in the space

$$E_T = \tilde{\mathcal{C}}_T(B_{p,1}^{\frac{N}{p}}) \times (\tilde{\mathcal{C}}_T(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{\mathcal{L}}_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})).$$

More precisely to get this bounds we will need to study the behavior of (a^n, v_1^n) .

3. We use compactness to prove that the sequence (a^n, u^n) converges, up to extraction, to a solution of (5.34).

Throughtout the proof, we denote $\underline{\nu} = \underline{b} \min(\mu, \lambda + 2\mu)$ and $\bar{\nu} = \mu + |\mu + \lambda|$, and we assume (with no loss of generality) that f belongs to $\tilde{\mathcal{L}}_T^1(B_{p_1,1}^{\frac{N}{p_1}})$.

First step

We smooth out the data as follows:

$$a_0^n = S_n a_0, \quad u_0^n = S_n u_0 \quad \text{and} \quad f^n = S_n f.$$

Note that we have:

$$\forall l \in \mathbb{Z}, \quad \|\Delta_l a_0^n\|_{L^p} \leq \|\Delta_l a_0\|_{L^p} \quad \text{and} \quad \|a_0^n\|_{B_{p,\infty}^{\frac{N}{p}}} \leq \|a_0\|_{B_{p,\infty}^{\frac{N}{p}}},$$

and similar properties for u_0^n and f^n , a fact which will be used repeatedly during the next steps. Now, according [13], one can solve (1.3) with the smooth data (a_0^n, u_0^n, f^n) . We get a solution (a^n, u^n) on a non trivial time interval $[0, T_n]$ such that:

$$a^n \in \tilde{C}([0, T_n], B_{2,1}^N) \quad \text{and} \quad u^n \in \tilde{C}([0, T_n], B_{2,1}^{\frac{N}{2}-1}) \cap \tilde{L}_{T_n}^1(B_{2,1}^{\frac{N}{2}+1}). \quad (5.35)$$

Uniform bounds

Let T_n be the lifespan of (a_n, u_n) , that is the supremum of all $T > 0$ such that (1.1) with initial data (a_0^n, u_0^n) has a solution which satisfies (5.35). Let T be in $(0, T_n)$. We aim at getting uniform estimates in E_T for T small enough. For that, we need to introduce the solution u_L^n to the linear system:

$$\partial_t u_L^n - \mathcal{A}u_L^n = f^n, \quad u_L^n(0) = u_0^n - \frac{1}{\nu} \tilde{v}_0 \tilde{v}_0.$$

Now, we set $\tilde{u}^n = u^n - u_L^n$ and the vectorfield $\tilde{v}_1^n = \tilde{u}^n - \frac{1}{\nu} \tilde{v}^n$ with $\text{div} \tilde{v}^n = P(\rho^n)$. We can check that \tilde{v}_1^n satisfies the parabolic system:

$$\begin{cases} \partial_t \tilde{v}_1^n + (u_L^n + \frac{1}{\nu} \tilde{v}^n) \cdot \nabla \tilde{v}_1^n + \tilde{v}_1^n \cdot \nabla u^n - (1 + a^n) \mathcal{A} \tilde{v}_1^n = a^n \mathcal{A} u_L^n - \frac{1}{\nu} (u_L^n \cdot \nabla \tilde{v}^n \\ \quad + \frac{1}{\nu} \tilde{v}^n \cdot \nabla \tilde{v}^n) - u_L^n \cdot \nabla u_L^n + \frac{1}{\nu} \nabla (\Delta)^{-1} (P'(\rho^n) \text{div}(\rho^n u^n)), \\ (\tilde{v}_1^n)_{t=0} = 0. \end{cases} \quad (5.36)$$

which has been studied in proposition 3.7. Define $m \in \mathbb{Z}$ by:

$$m = \inf \{ p \in \mathbb{Z} / 2\bar{\nu} \sum_{l \geq p} 2^{l \frac{N}{p}} \|\Delta_l a_0\|_{L^p} \leq c\bar{\nu} \} \quad (5.37)$$

where c is small enough positive constant (depending only N) to be fixed hereafter. In the sequel we will need of a control on $a - S_m a$ small to apply proposition 3.7, so here m is enough big (we explain how in the sequel). Let:

$$\bar{b} = 1 + \sup_{x \in \mathbb{R}^N} a_0(x), \quad A_0 = 1 + 2 \|a_0\|_{B_{p,1}^{\frac{N}{p}}}, \quad U_0 = \|u_0\|_{B_{p,1}^{\frac{N}{p}-1}} + \|a_0\|_{B_{p,1}^{\frac{N}{p}+1}} + \|f\|_{L_T^1(B_{p,1}^{\frac{N}{p}-1})}$$

and $\tilde{U}_0 = 2CU_0 + 4C\bar{\nu}A_0$ (where C' is a constant embedding and C stands for a large enough constant depending only N which will be determined when applying proposition 2.3, 3.7 and 4.9 in the following computations.) We assume that the following inequalities are fulfilled for some $\eta > 0$:

$$(\mathcal{H}_1) \quad \|a^n - S_m a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq c\nu\bar{\nu}^{-1},$$

$$(\mathcal{H}_2) \quad C\bar{\nu}^2 T \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}^2 \leq 2^{-2m} \underline{\nu},$$

$$(\mathcal{H}_3) \quad \frac{1}{2} \bar{b} \leq 1 + a^n(t, x) \leq 2\bar{b} \quad \text{for all } (t, x) \in [0, T] \times \mathbb{R}^N,$$

$$(\mathcal{H}_4) \quad \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq A_0,$$

$$\begin{aligned}
(\mathcal{H}_5) \quad & \|u_L^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+3})} \leq \eta, \\
(\mathcal{H}_6) \quad & \|\tilde{v}_1^n\|_{\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})} + \underline{\nu} \|\tilde{v}_1^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})} \leq \tilde{U}_0 \eta, \\
(\mathcal{H}_7) \quad & \|\tilde{v}^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}+1})} \leq C' A_0, \\
(\mathcal{H}_8) \quad & \|\nabla u^n\|_{\tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}}) + \tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq (\underline{\nu}^{-1} \tilde{U}_0 + 1) \eta
\end{aligned}$$

Remark that since:

$$1 + S_m a^n = 1 + a^n + (S_m a^n - a^n),$$

assumptions (\mathcal{H}_1) and (\mathcal{H}_3) combined with the embedding $B_{p,1}^{\frac{N}{p}} \hookrightarrow L^\infty$ insure that:

$$\inf_{(t,x) \in [0,T] \times \mathbb{R}^N} (1 + S_m a^n)(t, x) \geq \frac{1}{4} \underline{b}, \quad (5.38)$$

provided c has been chosen small enough (note that $\frac{\underline{\nu}}{\nu} \leq \bar{b}$).

We are going to prove that under suitable assumptions on T and η (to be specified below) if condition (\mathcal{H}_1) to (\mathcal{H}_7) are satisfied, then they are actually satisfied with strict inequalities. Since all those conditions depend continuously on the time variable and are strictly satisfied initially, a basic boobstrap argument insures that (\mathcal{H}_1) to (\mathcal{H}_8) are indeed satisfied for T . First we shall assume that η and T satisfies:

$$C(1 + \underline{\nu}^{-1} \tilde{U}_0) \eta + \frac{C'}{\underline{\nu}} A_0 T < \log 2 \quad (5.39)$$

so that denoting $\tilde{V}_1^n(t) = \int_0^t \|\nabla \tilde{v}_1^n\|_{B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1}} d\tau$, $\tilde{V}^n(t) = \frac{1}{\underline{\nu}} \int_0^t \|\nabla \tilde{v}^n\|_{B_{p,1}^{\frac{N}{p}}} d\tau$ and $U_L^n(t) = \int_0^t \|\nabla u_L^n\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+3}} d\tau$, we have, according to (\mathcal{H}_5) and (\mathcal{H}_6) :

$$e^{C(U_L^n + \tilde{V}_1^n + \tilde{V}^n)(T)} < 2 \quad \text{and} \quad e^{C(U_L^n + \tilde{V}_1^n + \tilde{V}^n)(T)} - 1 \leq 1. \quad (5.40)$$

In order to bound a^n in $\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})$, we apply inequality (4.30) and get:

$$\|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} < 1 + 2\|a_0\|_{B_{p,1}^{\frac{N}{p}}} = A_0. \quad (5.41)$$

Hence (\mathcal{H}_4) is satisfied with a strict inequality. (\mathcal{H}_7) verifies a strict inequality, it follows from proposition 2.5 and (\mathcal{H}_4) . Next, applying proposition 3.6 and proposition 2.5 yields:

$$\|u_L^n\|_{\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})} \leq U_0, \quad (5.42)$$

$$\begin{aligned}
\kappa \nu \|u_L^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+3})} & \leq \sum_{l \in \mathbb{Z}} 2^{l(\frac{N}{p_1}-1)} (1 - e^{-\kappa \nu 2^{2l} T}) (\|\Delta_l u_0\|_{L^{p_1}} + \\
\|\Delta_l f\|_{L^1(\mathbb{R}^+, L^{p_1})}) & \leq \sum_{l \in \mathbb{Z}} 2^{l(\frac{N}{p}+1)} (1 - e^{-\kappa \nu 2^{2l} T}) \|\Delta_l a_0\|_{L^p}.
\end{aligned} \quad (5.43)$$

Hence taking T such that:

$$\begin{aligned} & \sum_{l \in \mathbb{Z}} 2^{l(\frac{N}{p_1}-1)} (1 - e^{-\kappa\nu 2^{2l}T}) (\|\Delta_l u_0\|_{L^{p_1}} + \|\Delta_l f\|_{L^1(\mathbb{R}^+, L^{p_1})}) \\ & + \leq \sum_{l \in \mathbb{Z}} 2^{l(\frac{N}{p}+1)} (1 - e^{-\kappa\nu 2^{2l}T}) \|\Delta_l a_0\|_{L^p} < \kappa\eta\nu, \end{aligned} \quad (5.44)$$

insures that (\mathcal{H}_5) is strictly verified. Since (\mathcal{H}_1) , (\mathcal{H}_2) , (\mathcal{H}_5) , (\mathcal{H}_6) , (\mathcal{H}_7) and (5.38) are satisfied, proposition 3.7 may be applied, we obtain:

$$\begin{aligned} & \|\tilde{v}_1^n\|_{\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})} + \nu \|\tilde{v}_1^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})} \\ & \leq C e^{C(2U_L^n + 2\tilde{V}^n + \tilde{V}_1^n)(T)} \int_0^T (\|a^n \mathcal{A}u_L^n\|_{B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}}} + \|u_L^n \cdot \nabla u_L^n\|_{B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}}}) \\ & + \|u_L^n \cdot \nabla \tilde{v}^n\|_{B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}}} + \|\tilde{v}^n \cdot \nabla \tilde{v}^n\|_{B_{p,1}^{\frac{N}{p}}} + \|\nabla(\Delta)^{-1}(P'(\rho^n) \operatorname{div}(\rho^n u^n))\|_{B_{p,1}^{\frac{N}{p}}} dt. \end{aligned}$$

As $\frac{N}{p} + \frac{N}{p_1} - 1 \geq 0$, $2\frac{N}{p} - 1 > 0$ and by taking advantage of proposition 2.3, 2.1 and 2.5, we get:

$$\begin{aligned} & \|\nabla(\Delta)^{-1}(P'(\rho^n) \operatorname{div}(\rho^n u^n))\|_{\tilde{L}_T^1(B_{p,1}^{\frac{N}{p}})} \leq C_P (1 + \|a^n\|_{\tilde{L}^\infty(B_{p,1}^{\frac{N}{p}})}) (\sqrt{T} \|\tilde{v}_1^n\|_{\tilde{L}_T^2(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})}) \\ & + \sqrt{T} \|u_L^n\|_{\tilde{L}_T^2(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})} + T \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}, \\ & \|\tilde{v}^n \cdot \nabla \tilde{v}^n\|_{\tilde{L}_T^1(B_{p,1}^{\frac{N}{p}})} \leq C_1 T \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}^2. \end{aligned}$$

We proceed similarly for the other terms and we end up with:

$$\begin{aligned} & \|\tilde{v}_1^n\|_{\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})} + \nu \|\tilde{v}_1^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})} \leq C e^{C(2U_L^n + 2\tilde{V}^n + \tilde{V}_1^n)(T)} \\ & \times \left(C \|u_L^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+3})} (\bar{\nu} \|a^n\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} + \|u_L^n\|_{L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})}) + \right. \\ & C_1 T \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}^2 + C_P (1 + \|a^n\|_{\tilde{L}^\infty(B_{p,1}^{\frac{N}{p}})}) (\sqrt{T} \|\tilde{v}_1^n\|_{\tilde{L}_T^2(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})}) \\ & \left. + \sqrt{T} \|u_L^n\|_{\tilde{L}_T^2(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})} + T \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \right) + T \|u_L^n\|_{L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})} \times \\ & \qquad \qquad \qquad \|a^n\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}. \end{aligned} \quad (5.45)$$

with $C = C(N)$, $C_1 = C_1(N)$ and $C_P = (N, P, \underline{b}, \bar{b})$. Now, using assumptions (\mathcal{H}_4) , (\mathcal{H}_5) and (\mathcal{H}_6) , and inserting (5.40) in (5.45) gives:

$$\|\tilde{v}_1^n\|_{\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1})} + \|\tilde{v}_1^n\|_{L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1})} \leq 2C(\bar{\nu}A_0 + U_0)\eta + C_1 T A_0(1 + A_0) + \sqrt{T} A_0 U_0,$$

hence (\mathcal{H}_6) is satisfied with a strict inequality provided when T verifies:

$$2C(\bar{\nu}A_0 + U_0)\eta + C_1 T A_0(1 + A_0) + \sqrt{T} A_0 U_0 < C\bar{\nu}\eta. \quad (5.46)$$

(\mathcal{H}_8) verifies a strict inequality, it follows from proposition (\mathcal{H}_5) , (\mathcal{H}_6) and (\mathcal{H}_7) . We now have to check whether (\mathcal{H}_1) is satisfied with strict inequality. For that we apply proposition (4.10) which yields for all $m \in \mathbb{Z}$,

$$\sum_{l \geq m} 2^{l \frac{N}{2}} \|\Delta_l a^n\|_{L_T^\infty(L^p)} \leq \sum_{l \geq m} 2^{l \frac{N}{p}} \|\Delta_l a_0\|_{L^p} + (1 + \|a_0\|_{B_{p,1}^{\frac{N}{2}}}) (e^{C(U_L^n + \tilde{U}^n)(T)} - 1). \quad (5.47)$$

Using (5.39) and (\mathcal{H}_5) , (\mathcal{H}_6) , we thus get:

$$\|a^n - S_m a^n\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq \sum_{l \geq m} 2^{l \frac{N}{p}} \|\Delta_l a_0\|_{L^p} + \frac{C}{\log 2} (1 + \|a_0\|_{B_{p,1}^{\frac{N}{p}}}) (1 + \nu^{-1} \tilde{L}_0) \eta.$$

Hence (\mathcal{H}_1) is strictly satisfied provided that η further satisfies:

$$\frac{C}{\log 2} (1 + \|a_0\|_{B_{p,1}^{\frac{N}{p}}}) (1 + \nu^{-1} \tilde{U}_0) \eta < \frac{c\nu}{2\nu}. \quad (5.48)$$

In order to check whether (\mathcal{H}_3) is satisfied, we use the fact that:

$$a^n - a_0 = S_m(a^n - a_0) + (Id - S_m)(a^n - a_0) + \sum_{l > n} \Delta_l a_0,$$

whence, using $B_{p,1}^{\frac{N}{p}} \hookrightarrow L^\infty$ and assuming (with no loss of generality) that $n \geq m$,

$$\begin{aligned} \|a^n - a_0\|_{L^\infty((0,T) \times \mathbb{R}^N)} &\leq C (\|S_m(a^n - a_0)\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} + \|a^n - S_m a^n\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} \\ &\quad + 2 \sum_{l \geq m} 2^{l \frac{N}{p}} \|\Delta_l a_0\|_{L^p}). \end{aligned}$$

Changing the constant c in the definition of m and in (5.48) if necessary, one can, in view of the previous computations, assume that:

$$C (\|a^n - S_m a^n\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} + 2 \sum_{l \geq m} 2^{l \frac{N}{p}} \|\Delta_l a_0\|_{L^p}) \leq \frac{b}{4}.$$

As for the term $\|S_m(a^n - a_0)\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})}$, it may be bounded according proposition 4.10:

$$\begin{aligned} \|S_m(a^n - a_0)\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} &\leq (1 + \|a_0\|_{B_{p,1}^{\frac{N}{p}}}) (e^{C(\tilde{V}_1^n + \tilde{V}^n + U_L^n)(T)} - 1) + C 2^{2m} \sqrt{T} \|a_0\|_{B_{p,1}^{\frac{N}{p}}} \\ &\quad \times \|u^n\|_{L_T^2(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}})}. \end{aligned}$$

Note that under assumptions (\mathcal{H}_5) , (\mathcal{H}_6) , (5.39) and (5.48) (and changing c if necessary), the first term in the right-hand side may be bounded by $\frac{b}{8}$. Hence using interpolation, (5.42) and the assumptions (5.39) and (5.48), we end up with:

$$\|S_m(a^n - a_0)\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq \frac{b}{8} + C 2^{2m} \sqrt{T} \|a_0\|_{B_{2,1}^{\frac{N}{2}}} \sqrt{\eta(U_0 + \tilde{U}_0) \eta} (1 + \nu^{-1} \tilde{U}_0).$$

Assuming in addition that T satisfies:

$$C2^m \sqrt{\bar{T}} \|a_0\|_{B_{p,1}^{\frac{N}{p}}} \sqrt{\eta(U_0 + \tilde{U}_0 \eta)(1 + \nu^{-1} \tilde{U}_0)} < \frac{b}{8}, \quad (5.49)$$

and using the assumption $\underline{b} \leq 1 + a_0 \leq \bar{b}$ yields (\mathcal{H}_3) with a strict inequality. One can now conclude that if $T < T^n$ has been chosen so that conditions (5.44), (5.46) and (5.49) are satisfied (with η verifying (5.39) and (5.48), and m defined in (5.37) and $n \geq m$ then (a^n, u^n) satisfies (\mathcal{H}_1) to (\mathcal{H}_8) , thus is bounded independently of n on $[0, T]$. We still have to state that T^n may be bounded by below by the supremum \bar{T} of all times T such that (5.44), (5.46) and (5.49) are satisfied. This is actually a consequence of the uniform bounds we have just obtained, and of remark 12 and proposition 4.9. Indeed, by combining all these informations, one can prove that if $T^n < \bar{T}$ then (a^n, u^n) is actually in:

$$\tilde{L}_{T^n}^\infty(B_{2,1}^{\frac{N}{2}} \cap B_{p,1}^{\frac{N}{p}}) \times \left(\tilde{L}_{T^n}^\infty(B_{2,1}^{\frac{N}{2}} \cap (B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})) \cap L_{T^n}^1(B_{2,1}^{\frac{N}{2}+1} \cap (B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+2})) \right)^N$$

hence may be continued beyond \bar{T} (see the remark on the lifespan following the statement in [10]). We thus have $T^n \geq \bar{T}$.

Compactness arguments

We now have to prove that $(a^n, u^n)_{n \in \mathbb{N}}$ tends (up to a subsequence) to some function (a, u) which belongs to E_T . Here we recall that:

$$E_T = \tilde{C}([0, T], B_{p,1}^{\frac{N}{p}}) \times (\tilde{L}^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{L}^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})).$$

The proof is based on Ascoli's theorem and compact embedding for Besov spaces. As similar arguments have been employed in [10] or [13], we only give the outlines of the proof.

- Convergence of $(a^n)_{n \in \mathbb{N}}$:

We use the fact that $\tilde{a}^n = a^n - a_0^n$ satisfies:

$$\partial_t \tilde{a}^n = -u^n \cdot \nabla a^n + (1 + a^n) \operatorname{div} u^n.$$

Since $(u^n)_{n \in \mathbb{N}}$ is uniformly bounded in $\tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1}) \cap L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})$, it is by interpolation and the fact that $p_1 \leq p$, also bounded in $L_T^r(B_{p,1}^{\frac{N}{p}-1+\frac{2}{r}})$ for any $r \in [1, +\infty]$. By using the standard product laws in Besov spaces, we thus easily gather that $(\partial_t \tilde{a}^n)$ is uniformly bounded in $\tilde{L}_T^2(B_{p,1}^{\frac{N}{p}-1})$. Hence $(\tilde{a}^n)_{n \in \mathbb{N}}$ is bounded in $\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}-1} \cap B_{p,1}^{\frac{N}{p}})$ and equicontinuous on $[0, T]$ with values in $B_{p,1}^{\frac{N}{p}-1}$. Since the embedding $B_{p,1}^{\frac{N}{p}-1} \cap B_{p,1}^{\frac{N}{p}}$ is (locally) compact, and $(a_0^n)_{n \in \mathbb{N}}$ tends to a_0 in $B_{p,1}^{\frac{N}{p}}$, we conclude that $(a^n)_{n \in \mathbb{N}}$ tends (up to extraction) to some distribution a . Given that $(a^n)_{n \in \mathbb{N}}$ is bounded in $\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})$, we actually have $a \in \tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})$.

- Convergence of $(u_L^n)_{n \in \mathbb{N}}$:

From the definition of u_L^n and proposition 3.6, it is clear that $(u_L^n)_{n \in \mathbb{N}}$ tends to solution u_L to:

$$\partial_t u_L - \mathcal{A}u_L = f, \quad u_L(0) = u_0 - \frac{1}{\nu}.$$

$$\text{in } \tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+3}).$$

- Convergence of $(\tilde{v}_1^n)_{n \in \mathbb{N}}$:

We use the fact that:

$$\begin{aligned} \partial_t \tilde{v}_1^n &= -(u_L^n + \frac{1}{\nu} \tilde{v}^n) \cdot \nabla \tilde{v}_1^n - \tilde{v}_1^n \cdot \nabla u^n - \frac{1}{\nu} (u_L^n \cdot \nabla \tilde{v}^n - \frac{1}{\nu} \tilde{v}^n \cdot \nabla \tilde{v}^n) + (1 + a^n) \mathcal{A} \tilde{v}_1^n \\ &\quad + a^n \mathcal{A} u_L^n - u_L^n \cdot \nabla u_L^n + \frac{1}{\nu} \nabla (\Delta)^{-1} (P'(\rho^n) \operatorname{div}(\rho^n u^n)), \end{aligned}$$

As $(a^n)_{n \in \mathbb{N}}$ is uniformly bounded in $L_T^\infty(B_{p,1}^{\frac{N}{p}})$ and $(u^n)_{n \in \mathbb{N}}$ is uniformly bounded in $L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap L^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1})$, it is easy to see that the the right-hand side is uniformly bounded in $\tilde{L}_T^{\frac{4}{3}}(B_{p_1,1}^{\frac{N}{p_1}-\frac{3}{2}}) + \tilde{L}^\infty(B_{p,1}^{\frac{N}{p}-1})$. Hence $(\tilde{v}_1^n)_{n \in \mathbb{N}}$ is bounded in $\tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1})$ and equicontinuous on $[0, T]$ with values in $B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}-\frac{3}{2}}$. This enables to conclude that $(\tilde{v}_1^n)_{n \in \mathbb{N}}$ converges (up to extraction) to some function $\tilde{v}_1 \in \tilde{L}_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})$.

By interpolating with the bounds provided by the previous step, one obtains better results of convergence so that one can pass to the limit in the mass equation and in the momentum equation. Finally by setting $u = \tilde{v}_1 + \tilde{v} + u_L$, we conclude that (a, u) satisfies (1.3).

In order to prove continuity in time for a it suffices to make use of proposition 4.9. Indeed, a_0 is in $B_{p,1}^{\frac{N}{p}}$, and having $a \in \tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})$ and $u \in \tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1})$ insure that $\partial_t a + u \cdot \nabla a$ belongs to $\tilde{L}_T^1(B_{p,1}^{\frac{N}{p}})$. Similarly, continuity for u may be proved by using that $(\tilde{v}_1)_0 \in B_{p_1,1}^{\frac{N}{p_1}-1}$ and that $(\partial_t v_1 - \mu \Delta v_1) \in \tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}})$. We conclude by using the fact that $u = v_1 + \frac{1}{\nu} v$.

5.3 The proof of the uniqueness

Uniqueness when $1 \leq p_1 < 2N$, $\frac{2}{N} < \frac{1}{p} + \frac{1}{p_1}$ and $N \geq 3$

In this section, we focus on the cases $1 \leq p_1 < 2N$, $\frac{2}{N} < \frac{1}{p} + \frac{1}{p_1}$, $N \geq 3$ and postpone the analysis of the other cases (which turns out to be critical) to the next section. Throughout the proof, we assume that we are given two solutions (a^1, u^1) and (a^2, u^2) of (1.3). In the sequel we will show that $a^1 = a^2$ and $v_1^1 = v_1^2$ where $u^i = v_1^i + \tilde{v}^i$. It will imply that $u^1 = u^2$. We know that (a^1, v_1^1) and (a^2, v_1^2) belongs to:

$$\tilde{C}([0, T]; B_{p,1}^{\frac{N}{p}}) \times (\tilde{C}([0, T]; B_{p_1,1}^{\frac{N}{p_1}-1} + B_{p,1}^{\frac{N}{p}+1}) \cap \tilde{L}^1(0, T; B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2}))^N.$$

Let $\delta a = a^2 - a^1$, $\delta v = \tilde{v}^2 - \tilde{v}^1$ and $\delta v_1 = v_1^2 - v_1^1$. The system for $(\delta a, \delta v_1)$ reads:

$$\left\{ \begin{array}{l} \partial_t \delta a + u^2 \cdot \nabla \delta a = \delta a \operatorname{div} u^2 + (\delta v_1 + \frac{1}{\nu} \delta v) \cdot \nabla a^1 + (1 + a^1) \operatorname{div} (\delta v_1 + \frac{1}{\nu} \delta v), \\ \partial_t \delta v_1 + u^2 \cdot \delta \nabla v_1 + \delta v_1 \cdot \nabla u^1 - (1 + a^1) \mathcal{A} \delta v_1 = \delta a \mathcal{A} v_1^2 - \frac{1}{\nu} (u^2 \cdot \nabla \delta \tilde{v} \\ - \delta \tilde{v} \cdot \nabla u^1) + \nabla (\Delta)^{-1} \left((P'(\rho^2) - P'(\rho^1)) \operatorname{div} (\rho^2 u^2) + P'(\rho^1) \operatorname{div} (\rho^1 \delta u) \right. \\ \left. + P'(\rho^1) \operatorname{div} ((\rho^2 - \rho^1) u^2) \right). \end{array} \right. \quad (5.50)$$

The function δa may be estimated by taking advantage of proposition 4.9 with $s = \frac{N}{p} - 1$.

Denoting $U^i(t) = \|\nabla u^i\|_{\tilde{L}^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1})}$ for $i = 1, 2$, we get for all $t \in [0, T]$,

$$\begin{aligned} \|\delta a(t)\|_{B_{p,1}^{\frac{N}{p}-1}} &\leq C e^{CU^2(t)} \int_0^t e^{-CU^2(\tau)} \|\delta a \operatorname{div} u^2 + (\delta v_1 + \frac{1}{\nu} \delta v) \cdot \nabla a^1 \\ &\quad + (1 + a^1) \operatorname{div} (\delta v_1 + \frac{1}{\nu} \delta v)\|_{B_{p,1}^{\frac{N}{p}-1}} d\tau, \end{aligned}$$

Next using proposition 2.3 and 2.5 we obtain:

$$\begin{aligned} \|\delta a(t)\|_{B_{p,1}^{\frac{N}{p}-1}} &\leq C e^{CU^2(t)} \int_0^t e^{-CU^2(\tau)} \|\delta a\|_{B_{p,1}^{\frac{N}{p}-1}} \left(\|u^2\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1}} + (1 + 2\|a_1\|_{B_{p,1}^{\frac{N}{p}}}) \right) \\ &\quad + (1 + 2\|a_1\|_{B_{p,1}^{\frac{N}{p}}}) \|\delta v_1\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1}} d\tau, \end{aligned}$$

Hence applying Grönwall lemma, we get:

$$\|\delta a(t)\|_{B_{p,1}^{\frac{N}{p}-1}} \leq C e^{CU^2(t)} \int_0^t e^{-CU^2(\tau)} (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}}) \|\delta v_1\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1}} d\tau. \quad (5.51)$$

For bounding δv_1 , we aim at applying proposition 3.7 to the second equation of (5.50). So let us fix an integer m such that:

$$1 + \inf_{(t,x) \in [0,T] \times \mathbb{R}^N} S_m a^1 \geq \frac{b}{2} \quad \text{and} \quad \|a^1 - S_m a^1\|_{L_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq c \frac{\nu}{\nu}. \quad (5.52)$$

Note since a^1 satisfies a transport equation with right-hand side in $\tilde{L}_T^1(B_{p,1}^{\frac{N}{p}-1})$, proposition 4.9 guarantees that a^1 is in $\tilde{C}_T(B_{p,1}^{\frac{N}{p}})$. Hence such an integer does exist (see remark 11). Now applying corollary 3.8 with $s = \frac{N}{p_1} - 2$ and $s' = \frac{N}{p} - 1$ insures that for all time $t \in [0, T]$, we have:

$$\begin{aligned} \|\delta v_1\|_{L_t^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+1})} &\leq C e^{CU(t)} \int_0^t e^{-CU(\tau)} \left(\|\delta a \mathcal{A} v_1^2 - \frac{1}{\nu} (\delta v \cdot \nabla v_1^1 + v_1^1 \cdot \nabla \delta v) \right. \\ &\quad \left. - \frac{1}{\nu^2} (v^1 \cdot \nabla \delta v + \delta v \cdot \nabla v^2) \right)_{B_{p_1,1}^{\frac{N}{p_1}-2} + B_{p,1}^{\frac{N}{p}-1}} d\tau, \end{aligned}$$

with $U(t) = U^1(t) + U^2(t) + 2^{2m} \underline{\nu}^{-1} \bar{\nu}^2 \int_0^t \|a^1\|_{B_{p,1}^{\frac{N}{p}}}^2 d\tau$.

Hence, applying proposition 2.3 we get:

$$\begin{aligned} \|\delta v_1\|_{\tilde{L}_t^1(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})} &\leq C e^{CU(t)} \int_0^t e^{-CU(\tau)} (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}} + \|a^2\|_{B_{p,1}^{\frac{N}{p}}} \\ &\quad + \|v_1^2\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2}}) \|\delta a\|_{B_{p,1}^{\frac{N}{p}-1}} d\tau. \end{aligned} \quad (5.53)$$

Finally plugging (5.51) in (5.53), we get for all $t \in [0, T_1]$,

$$\begin{aligned} \|\delta v_1\|_{\tilde{L}_t^1(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})} &\leq C e^{CU(t)} \int_0^t (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}} + \|a^2\|_{B_{p,1}^{\frac{N}{p}}} + \|v_1^2\|_{B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2}}) \\ &\quad \times \|\delta v_1\|_{B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1}} d\tau. \end{aligned}$$

Since a^1 and a^2 are in $L^\infty(B_{p,1}^{\frac{N}{p}})$ and v_1^2 belongs to $L_T^1(B_{p_1,1}^{\frac{N}{p_1}+1} + B_{p,1}^{\frac{N}{p}+2})$, applying Grönwall lemma yields $\delta v_1 = 0$, on $[0, T]$.

Uniqueness when: $\frac{2}{N} = \frac{1}{p_1} + \frac{1}{p}$ **or** $p_1 = 2N$ **or** $N = 2$.

The above proof fails in dimension two. One of the reasons why is that the product of functions does not map $B_{p,1}^{\frac{N}{p}} \times B_{p_1,1}^{\frac{N}{p_1}-2}$ in $B_{p_1,1}^{\frac{N}{p_1}-2}$ but only in the larger space $B_{p_1,\infty}^{\frac{N}{p_1}-2}$. This induces us to bound δa in $L_T^\infty(B_{p,\infty}^{\frac{N}{p}-1})$ and δv_1 in $L_T^\infty(B_{p_1,\infty}^{\frac{N}{p_1}-2} + B_{p,\infty}^{\frac{N}{p}}) \cap L_T^1(B_{p_1,\infty}^{\frac{N}{p_1}} + B_{p,\infty}^{\frac{N}{p}+1})$ (or rather, in the widetilde version of those spaces, see below). Yet, we are in trouble because due to $B_{p_1,\infty}^{\frac{N}{p_1}}$ is not embedded in L^∞ , the term $\delta v_1 \cdot \nabla a^1$ in the right hand-side of the first equation of (5.50) cannot be estimated properly. As noticed in [12], this second difficulty may be overcome by making use of logarithmic interpolation and Osgood lemma (a substitute for Gronwall inequality). Let us now tackle the proof. Fix an integer m such that:

$$1 + \inf_{(t,x) \in [0,T] \times \mathbb{R}^N} S_m a^1 \geq \frac{b}{2} \quad \text{and} \quad \|a^1 - S_m a^1\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq c \frac{\underline{\nu}}{\bar{\nu}}, \quad (5.54)$$

and define T_1 as the supremum of all positive times t such that:

$$t \leq T \quad \text{and} \quad t \bar{\nu}^2 \|a^1\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})} \leq c 2^{-2m} \underline{\nu}. \quad (5.55)$$

Remark that the proposition 4.9 ensures that a^1 belongs to $\tilde{C}_T(B_{p,1}^{\frac{N}{p}})$ so that the above two assumptions are satisfied if m has been chosen large enough. For bounding δa in $L_T^\infty(B_{p,\infty}^{\frac{N}{p}-1})$, we apply proposition 4.9 with $r = +\infty$ and $s = 0$. We get (with the notation of the previous section):

$$\begin{aligned} \forall t \in [0, T], \quad \|\delta a(t)\|_{B_{p,\infty}^{\frac{N}{p}-1}} &\leq C e^{CU^2(t)} \int_0^t e^{-CU^2(\tau)} \|\delta a \operatorname{div} u^2 + (\delta v_1 + \frac{1}{\nu} \delta v) \cdot \nabla a^1 \\ &\quad + (1 + a^1) \operatorname{div} (\delta v_1 + \frac{1}{\nu} \delta v)\|_{B_{p,\infty}^{\frac{N}{p}-1}} d\tau, \end{aligned}$$

hence using that the product of two functions maps $B_{p,\infty}^{\frac{N-1}{p}} \times B_{p,1}^{\frac{N}{p}}$ in $B_{p,\infty}^{\frac{N-1}{p}}$, and applying Gronwall lemma,

$$\|\delta a(t)\|_{B_{p,\infty}^{\frac{N-1}{p}}} \leq C e^{CU^2(t)} \int_0^t e^{-CU^2(\tau)} (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}}) \|\delta v_1\|_{B_{p,1}^{\frac{N}{p}} + B_{p,1}^{\frac{N+1}{p}}} d\tau. \quad (5.56)$$

Next, using proposition 3.8 combined with proposition 2.3 and corollary 2 in order to bound the nonlinear terms, we get for all $t \in [0, T_1]$:

$$\begin{aligned} \|\delta v_1\|_{\tilde{L}_T^1(B_{p,1}^{\frac{N-2}{p}} + B_{p,\infty}^{\frac{N}{p}+1})} &\leq C e^{C(U^1+U^2)(t)} \int_0^t (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}} + \|a^2\|_{B_{p,1}^{\frac{N}{p}}}) \\ &\quad + \|v_1^2\|_{B_{p,1}^{\frac{N}{p}+1} + B_{p,1}^{\frac{N}{p}+2}} \|\delta a\|_{B_{p,\infty}^{\frac{N-1}{p}}} d\tau. \end{aligned} \quad (5.57)$$

In order to control the term $\|\delta v_1\|_{B_{p,1}^{\frac{N}{p}} + B_{p,1}^{\frac{N+1}{p}}}$ which appears in the right-hand side of (5.56), we make use of the following logarithmic interpolation inequality whose proof may be found in [12], page 120:

$$\begin{aligned} \|\delta v_1\|_{L_t^1(B_{p,1}^{\frac{N}{p}} + B_{p,1}^{\frac{N+1}{p}})} &\lesssim \\ \|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N}{p}})} \log \left(e + \frac{\|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N-1}{p}})} + \|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N+1}{p}})}}{\|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N}{p}})}} \right) \\ &\quad + \|\delta v_1\|_{\tilde{L}_t^1(B_{p,\infty}^{\frac{N}{p}+1})} \log \left(e + \frac{\|\delta v_1\|_{\tilde{L}_t^1(B_{p,\infty}^{\frac{N}{p}})} + \|\delta v_1\|_{\tilde{L}_t^1(B_{p,\infty}^{\frac{N+2}{p}})}}{\|\delta v_1\|_{\tilde{L}_t^1(B_{p,\infty}^{\frac{N}{p}})}} \right). \end{aligned} \quad (5.58)$$

Because v_1^1 and v_2^2 belong to $\tilde{L}_T^\infty(B_{p,1}^{\frac{N-1}{p}} + B_{p,1}^{\frac{N+1}{p}}) \cap L_T^1(B_{p,1}^{\frac{N+1}{p}} + B_{p,1}^{\frac{N+2}{p}})$, the numerator in the right-hand side may be bounded by some constant C_T depending only on T and on the norms of v_1^1 and v_2^2 . Therefore inserting (5.56) in (5.57) and taking advantage of (5.58), we end up for all $t \in [0, T_1]$ with:

$$\begin{aligned} \|\delta v_1\|_{\tilde{L}_T^1(B_{p,1}^{\frac{N}{p}} + B_{p,1}^{\frac{N+1}{p}})} &\leq C (1 + \|a^1\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p}})}) \\ &\quad \times \int_0^t (1 + \|a^1\|_{B_{p,1}^{\frac{N}{p}}} + \|a^2\|_{B_{p,1}^{\frac{N}{p}}} + \|v_1^2\|_{B_{p,1}^{\frac{N}{p}+1} + B_{p,1}^{\frac{N}{p}+2}}) \|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N}{p}})} \\ &\quad \times \log(e + C_T \|\delta v_1\|_{\tilde{L}_t^1(B_{p,1}^{\frac{N}{p}} + B_{p,1}^{\frac{N+1}{p}})})^{-1} d\tau. \end{aligned}$$

Since the function $t \rightarrow \|a^1(t)\|_{B_{p,1}^{\frac{N}{p}}} + \|a^2(t)\|_{B_{p,1}^{\frac{N}{p}}} + \|v_1^2(t)\|_{B_{p,1}^{\frac{N}{p}+1} + B_{p,1}^{\frac{N}{p}+2}}$ is integrable on $[0, T]$, and:

$$\int_0^1 \frac{dr}{r \log(e + C_T r^{-1})} = +\infty$$

Osgood lemma yields $\|\delta v_1\|_{\tilde{L}_T^1(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1})} = 0$. Note that the definition of m depends only on T and that (5.52) is satisfied on $[0, T]$. Hence, the above arguments may be repeated on $[T_1, 2T_1]$, $[2T_1, 3T_1]$, etc. until the whole interval $[0, T]$ is exhausted. This yields uniqueness on $[0, T]$ for a and v_1 which implies uniqueness for u .

5.4 Proof of corollary 1

The proof follows the same line as theorem 1.1 except concerning the term of rest $\nabla(\Delta)^{-1}(P'(\rho)\text{div}(\rho u))$ in the momentum equation of system (5.34). Indeed in our case this term can be written simply on the form ρu . In this case we control this term in $\tilde{L}^2(B^{\frac{N}{p}} p, 1)$ without imposing additional conditions on p of type $2\frac{N}{p} - 1 > 0$.

Now the difficulty is to prove the uniqueness. For that we use the main theorem of D. Hoff in [18] which is a result weak-strong uniqueness. In this article, D. Hoff has two solutions (ρ, u) and (ρ_1, u_1) with the same initial data (ρ_0, u_0) and he shows that under some hypothesis of regularity on (ρ_1, u_1) and (ρ_2, u_2) then $\rho_1 = \rho_2$, $u_1 = u_2$. We now discuss that our solution checks the conditions required in [18]. More precisely we have to show that our solution (ρ, u) verifies all the hypotheses asked on (ρ_1, u_1) and (ρ_2, u_2) . The check is easy and tedious, but only one hypothesis is required to be careful and is in fact the main condition why D. Hoff does not get a global strong solution in dimension $N = 3$ for the solutions built in [19]. We need to check that $u \in L_{loc}^\infty((0, T], L^\infty)$ and $\nabla u \in L^1((0, T], L^\infty)$. In our case we have $\nabla u = \nabla v_1 + \frac{1}{\nu} \nabla v$ where we recall that $\text{div} v = P(\rho) - P(\bar{\rho})$. We know that by interpolation $\nabla v_1 \in L_T^1(B_{p_1,1}^{\frac{N}{p_1}} + B_{p,1}^{\frac{N}{p}+1}) \hookrightarrow L_T^1(L^\infty)$ and by proposition 2.5 $\nabla v \in L_T^\infty(B_{p,1}^{\frac{N}{p}})$. We obtain then $\nabla u \in L_T^1(L^\infty)$. We have now to show that $u \in L_T^\infty(L^\infty)$. In fact we have just to apply classical energy inequalities, so we multiply the momentum equation by $u|u|^{p_1-2}$

$$\begin{aligned} & \frac{1}{p_1} \int_{\mathbb{R}^N} \rho |u|^{p_1}(t, x) dx + \mu \int_0^t \int_{\mathbb{R}^N} |u|^{p_1-2} |\nabla u|^2(t, x) dt dx + \frac{p_1 - 2}{4} \mu \int_0^t \int_{\mathbb{R}^N} |u|^{p_1-4} |\nabla |u|^2|^2(t, x) dx dt \\ & + \int_0^t \int_{\mathbb{R}^N} (P(\rho) - P(\bar{\rho})) (\text{div} u |u|^{p_1-2} + (p_1 - 2) \sum_{i,k} u_i u_k \partial_i u_k |u|^{p_1-4})(t, x) dt dx \\ & \leq \int_{\mathbb{R}^N} \rho_0 |u_0|^{p_1} dx. \end{aligned}$$

By Young's inequalities and the fact that $P(\rho) - P(\bar{\rho})$ belongs in $L^\infty(L^2 \cap L^\infty)$ we obtain that for all $p_1 \in [1, +\infty[$, $\rho^{\frac{1}{p_1}} u \in L^\infty(L^{p_1})$ and:

$$\|\rho^{\frac{1}{p_1}} u\|_{L^\infty(L^{p_1})} \leq C_0,$$

where C_0 depends only on the initial data. As $\frac{1}{\rho} \in L^\infty$, we conclude that u is uniformly bounded in all spaces $L^\infty(L^{p_1})$ with $p_1 \in [1, +\infty[$. We conclude then $u \in L_T^\infty(L^\infty)$.

6 Continuation criterions

Proof of theorem 1.2

We now prove theorem 1.2. We have assumed here that $\rho_0^{\frac{1}{p_1}} u_0 \in L^{p_1}$ with $p_1 > N$. We want show that with our hypothesis in particular that $a \in L_T^\infty$ and $1+a$ bounded away on $[0, T]$, then we are able to show that $\rho^{\frac{1}{p_1}} u \in L_T^\infty(L^{p_1})$. In this case as $1+a$ is bounded away we show that $u \in L_T^\infty(L^{p_1})$ and by embedding we get $u \in L_T^\infty(B_{p_1,1}^{\frac{N}{p_1}-1})$ as $\frac{N}{p_1} \leq 0$. We can next conclude by the fact that $(a(T, \cdot), u(T, \cdot)) \in (B_{p,1}^{\frac{N}{p}} \times B_{p_1,1}^{\frac{N}{p_1}-1})$ so that we can extend our solutions. Finally we have just to show that $\rho^{\frac{1}{p_1}} u \in L_T^\infty(L^{p_1})$, in this goal we have just to apply classical energy inequality. We multiply the momentum equation by $u|u|^{p_1-2}$ and we get after integration by part:

$$\begin{aligned} & \frac{1}{p_1} \int_{\mathbb{R}^N} \rho |u|^{p_1}(t, x) dx + \mu \int_0^t \int_{\mathbb{R}^N} |u|^{p_1-2} |\nabla u|^2(t, x) dt dx + \frac{p_1-2}{4} \mu \int_0^t \int_{\mathbb{R}^N} |u|^{p_1-4} |\nabla |u|^2|^2(t, x) dx dt \\ & + \lambda \int_0^t \int_{\mathbb{R}^N} (\operatorname{div} u)^2 |u|^{p_1-2}(t, x) dt dx + \lambda \frac{p_1-2}{2} \int_0^t \int_{\mathbb{R}^N} \operatorname{div} u \sum_i u_i \partial_i |u|^2 |u|^{p_1-4}(t, x) dt dx - \\ & \int_0^t \int_{\mathbb{R}^N} (P(\rho) - P(\bar{\rho})) (\operatorname{div} u |u|^{p_1-2} + (p_1-2) \sum_{i,k} u_i u_k \partial_i u_k |u|^{p_1-4})(t, x) dt dx \\ & \leq \int_{\mathbb{R}^N} \rho_0 |u_0|^{p_1} dx. \end{aligned}$$

By Young's inequalities, inequality (1.4) and the fact that $P(\rho) - P(\bar{\rho})$ belongs in $L^\infty(L^1 \cap L^\infty)$ we conclude the proof.

7 Appendix

This section is devoted to the proof of commutator estimates which have been used in section 2 and 3. They are based on paradifferential calculus, a tool introduced by J.-M. Bony in [4]. The basic idea of paradifferential calculus is that any product of two distributions u and v can be formally decomposed into:

$$uv = T_u v + T_v u + R(u, v) = T_u v + T'_v u$$

where the paraproduct operator is defined by $T_u v = \sum_q S_{q-1} u \Delta_q v$, the remainder operator R , by $R(u, v) = \sum_q \Delta_q u (\Delta_{q-1} v + \Delta_q v + \Delta_{q+1} v)$ and $T'_v u = T_v u + R(u, v)$. Inequalities (3.18) and (3.26) are consequence of the following lemma:

Lemma 1 *Let $1 \leq p_1 \leq p \leq +\infty$ and $\sigma \in (-\min(\frac{N}{p}, \frac{N}{p_1}), \frac{N}{p} + 1]$. There exists a sequence $c_q \in l^1(\mathbb{Z})$ such that $\|c_q\|_{l^1} = 1$ and a constant C depending only on N and σ such that:*

$$\forall q \in \mathbb{Z}, \quad \|[v \cdot \nabla, \Delta_q] a\|_{L^{p_1}} \leq C c_q 2^{-q\sigma} \|\nabla v\|_{B_{p,1}^{\frac{N}{p}}} \|a\|_{B_{p_1,1}^\sigma}. \quad (7.59)$$

In the limit case $\sigma = -\min(\frac{N}{p}, \frac{N}{p_1})$, we have:

$$\forall q \in \mathbb{Z}, \quad \|[v \cdot \nabla, \Delta_q]a\|_{L^{p_1}} \leq Cc_q 2^{q\frac{N}{p}} \|\nabla v\|_{B_{p_1,1}^{\frac{N}{p}}} \|a\|_{B_{p,\infty}^{-\frac{N}{p_1}}}. \quad (7.60)$$

Finally, for all $\sigma > 0$ and $\frac{1}{p_2} = \frac{1}{p_1} - \frac{1}{p}$, there exists a constant C depending only on N and on σ and a sequence $c_q \in l^1(\mathbb{Z})$ with norm 1 such that:

$$\forall q \in \mathbb{Z}, \quad \|[v \cdot \nabla, \Delta_q]v\|_{L^p} \leq Cc_q 2^{-q\sigma} (\|\nabla v\|_{L^\infty} \|v\|_{B_{p_1,1}^\sigma} + \|\nabla v\|_{L^{p_2}} \|\nabla v\|_{B_{p_1,1}^{\sigma-1}}). \quad (7.61)$$

Proof: These results are proved in [3] chapter 2. \square

Inequality (3.19) is a consequence of the following lemma:

Lemma 2 Let $1 \leq p_1 \leq p \leq +\infty$ and $\alpha \in (1 - \frac{N}{p}, 1]$, $k \in \{1, \dots, N\}$ and $R_q = \Delta_q(a\partial_k w) - \partial_k(a\Delta_q w)$. There exists $c = c(\alpha, N, \sigma)$ such that:

$$\sum_q 2^{q\sigma} \|R_q\|_{L^{p_1}} \leq C \|a\|_{B_{p_1,1}^{\frac{N}{p}+\alpha}} \|w\|_{B_{p_1,1}^{\sigma+1-\alpha}} \quad (7.62)$$

whenever $-\frac{N}{p} < \sigma \leq \alpha + \frac{N}{p}$.

In the limit case $\sigma = -\frac{N}{p}$, we have for some constant $C = C(\alpha, N)$:

$$\sup_q 2^{-q\frac{N}{p}} \|R_q\|_{L^{p_1}} \leq C \|a\|_{B_{p_1,1}^{\frac{N}{p}+\alpha}} \|w\|_{B_{p_1,\infty}^{-\frac{N}{p}+1-\alpha}}. \quad (7.63)$$

Proof The proof is almost the same as the one of lemma A3 in [10]. It is based on Bony's decomposition which enables us to split R_q into:

$$R_q = \underbrace{\partial_k[\Delta_q, T_a]w}_{R_q^1} - \underbrace{\Delta_q T_{\partial_k a} w}_{R_q^2} + \underbrace{\Delta_q T_{\partial_k w} w}_{R_q^3} + \underbrace{\Delta_q R(\partial_k w, a)}_{R_q^4} - \underbrace{\partial_k T'_{\Delta_q w} a}_{R_q^5}.$$

Using the fact that:

$$R_q^1 = \sum_{q'=q-4}^{q+4} \partial_k[\Delta_q, S_{q'-1} a] \Delta_{q'} w,$$

and the mean value theorem, we readily get under the hypothesis that $\alpha \leq 1$,

$$\sum_q 2^{q\sigma} \|R_q^1\|_{L^{p_1}} \lesssim \|\nabla a\|_{B_{\infty,1}^{\alpha-1}} \|w\|_{B_{p_1,1}^{\sigma+1-\alpha}}. \quad (7.64)$$

Standard continuity results for the paraproduct insure that R_q^2 satisfies (7.64) and that:

$$\sum_q 2^{q\sigma} \|R_q^1\|_{L^{p_1}} \lesssim \|\nabla w\|_{B_{\infty,1}^{\sigma-\alpha-\frac{N}{p_1}}} \|a\|_{B_{p_1,1}^{\frac{N}{p}+\alpha}}. \quad (7.65)$$

provided $\sigma - \alpha - \frac{N}{p} \leq 0$. Next, standard continuity result for the remainder insure that under the hypothesis $\sigma > -\frac{N}{p}$, we have:

$$\sum_q 2^{q\sigma} \|R_q^1\|_{L^{p_1}} \lesssim \|\nabla w\|_{B_{p_1,1}^{\sigma-\alpha}} \|a\|_{B_{p_1,1}^{\frac{N}{p}+\alpha}}. \quad (7.66)$$

For bounding R_q^5 we use the decomposition: $R_q^5 = \sum_{q' \geq q-3} \partial_k (S_{q'+2} \Delta_q w \Delta_{q'} a)$, which leads (after a suitable use of Bernstein and Hölder inequalities) to:

$$2^{q\sigma} \|R_q^5\|_{L^{p_1}} \lesssim \sum_{q' \geq q-2} 2^{(q-q')(\alpha + \frac{N}{p_1} - 1)} 2^{q(\sigma+1-\alpha)} \|\Delta_q w\|_{L^{p_1}} 2^{q'(\frac{N}{p_1} + \alpha)} \|\Delta_{q'} a\|_{L^p}.$$

Hence, since $\alpha + \frac{N}{p} - 1 > 0$, we have:

$$\sum_q 2^{q\sigma} \|R_q^5\|_{L^p} \lesssim \|\nabla w\|_{B_{p_1,1}^{\sigma+1-\alpha}} \|a\|_{B_{p,1}^{\frac{N}{p} + \alpha}}.$$

Combining this latter inequality with (7.64), (7.65) and (7.66), and using the embedding $B_{p,1}^{\frac{N}{p}} \hookrightarrow B_{\infty,1}^{r-\frac{N}{p}}$ for $r = \frac{N}{p} + \alpha - 1$, σ_α completes the proof of (7.62).

The proof of (7.63) is almost the same: for bounding R_q^1 , R_q^2 , R_q^3 and R_q^5 , it is just a matter of changing \sum_q into \sup_q . \square

Remark 13 For proving proposition 3.8, we shall actually use the following non-stationary version of inequality (7.63):

$$\sup_q 2^{-q\frac{N}{p}} \|R_q\|_{L_T^1(L^{p_1})} \leq C \|a\|_{\tilde{L}_T^\infty(B_{p,1}^{\frac{N}{p} + \alpha})} \|w\|_{\tilde{L}_T^1(B_{p_1,\infty}^{-\frac{N}{p_1} + 1 - \alpha})},$$

which may be easily proved by following the computations of the previous proof, dealing with the time dependence according to Hölder inequality.

References

- [1] H. Abidi and M. Paicu. Équation de Navier-Stokes avec densité et viscosité variables dans l'espace critique. *Annales de l'institut Fourier*, 57 no. 3 (2007), p. 883-917.
- [2] H. Bahouri and J.-Y. Chemin, Équations d'ondes quasilineaires et estimation de Strichartz, *Amer. J. Mathematics*. 121 (1999) 1337-1377.
- [3] H. Bahouri, J.-Y. Chemin and R. Danchin. Fourier analysis and nonlinear partial differential equations, *to appear in Springer*.
- [4] J.-M. Bony, Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires, *Annales Scientifiques de l'école Normale Supérieure*. 14 (1981) 209-246.
- [5] D. Bresch and B. Desjardins, Existence of global weak solutions for a 2D Viscous shallow water equations and convergence to the quasi-geostrophic model. *Comm. Math. Phys.*, 238(1-2): 211-223, 2003.
- [6] J.-Y. Chemin, Théorèmes d'unicité pour le système de Navier-Stokes tridimensionnel, *J.d'Analyse Math.* 77 (1999) 27-50.
- [7] J.-Y. Chemin, About Navier-Stokes system, *Prépublication du Laboratoire d'Analyse Numérique de Paris 6*, R96023 (1996).

- [8] J.-Y. Chemin and N. Lerner, Flot de champs de vecteurs non lipschitziens et équations de Navier-Stokes, *J.Differential Equations*, 121 (1992) 314-328.
- [9] R. Danchin, Fourier analysis method for PDE's, *Preprint*, Novembre 2005.
- [10] R. Danchin, Local Theory in critical Spaces for Compressible Viscous and Heat-Conductive Gases, *Communication in Partial Differential Equations*, 26 (78),1183-1233, (2001).
- [11] R. Danchin, Global Existence in Critical Spaces for Flows of Compressible Viscous and Heat-Conductive Gases, *Arch.Rational Mech.Anal.*160, (2001), 1-39.
- [12] R. Danchin, On the uniqueness in critical spaces for compressible Navier-Stokes equations. *NoDEA Nonlinear Differential Equations Appl*, 12(1):111-128, 2005.
- [13] R. Danchin, Well-Posedness in Critical Spaces for Barotropic Viscous Fluids with Truly Not Constant Density, *Communications in Partial Differential Equations*,32:9,1373-1397.
- [14] Weak strong uniqueness for the isentropic compressible Navier-Stokes system, *preprint*.
- [15] B. Haspot, Cauchy problem for viscous shallow water equations with a term of capillarity , *accepted in HYP 2008*.
- [16] B. Haspot, Local well-posedness results for density-dependent incompressible fluids, *Arxiv:0902.1982*, (February 2009).
- [17] D. Hoff. Global existence for 1D, compressible, isentropic Navier-Stokes equations with large initial data. *Trans. Amer. Math. Soc*, 303(1), 169-181, 1987.
- [18] D. Hoff, Uniqueness of weak solutions of the NavierStokes equations of multidimensional, compressible flow, *SIAM J. Math. Anal.* 37 (6) (2006).
- [19] D. Hoff. Discontinuous solutions of the Navier-Stokes equations for multidimensional flows of the heat conducting fluids. *Arch. Rational Mech. Anal.*, 139, (1997), p. 303-354.
- [20] D. Hoff. Global solutions of the Navier-Stokes equations for multidimensional compressible flow with discontinuous initial data. *J. Differential Equations*, 120(1), 215-254, 1995.
- [21] D. Hoff. Strong convergence to global solutions for multidimensional flows of compressible, viscous fluids with polytropic equations of state and discontinuous initial data. *Arch. Rational Mech. Anal.*, 132(1), 1-14, 1995.
- [22] D. Hoff and K. Zumbrum. Multi-dimensional diffusion waves for the Navier-Stokes equations of compressible flow, *Indiana University Mathematics Journal*, 1995, 44, 603-676.
- [23] A. V. Kazhikov. The equation of potential flows of a compressible viscous fluid for small Reynolds numbers: existence, uniqueness and stabilization of solutions. *Sibirsk. Mat. Zh.*, 34 (1993), no. 3, p. 70-80.

- [24] A. V. Kazhikov and V. V. Shelukhin. Unique global solution with respect to time of initial-boundary value problems for one- dimensional equations of a viscous gas. *Prikl. Mat. Meh.*, 41(2): 282-291, 1977.
- [25] Akitaka Matsumura and Takaaki Nishida. The initial value problem for the equations of motion of compressible viscous and heat-conductive gases. *J. Math. Kyoto Univ.*, 20(1): 67-104, 1980.
- [26] Akitaka Matsumura and Takaaki Nishida. The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids. *Proc. Japan Acad. Ser. A Math. Sci*, 55(9):337-342, 1979.
- [27] Y. Meyer. Wavelets, paraproducts, and Navier-Stokes equation. *In Current developments in mathematics, 1996 (Cambridge, MA)*, page 105-212. Int. Press, Boston, MA, 1997.
- [28] J. Nash. Le problème de Cauchy pour les équations différentielles d'un fluide général. *Bull. Soc. Math. France*, 90: 487-497, 1962.
- [29] T. Runst and W. Sickel: Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations. *de Gruyter Series in Nonlinear Analysis and Applications*, 3. Walter de Gruyter and Co., Berlin (1996) na 21, no. 1 (2005), 1-24.
- [30] D. Serre. Solutions faibles globales des équations de Navier-Stokes pour un fluide compressible. *Comptes rendus de l'Académie des sciences. Série 1*, 303(13): 639-642, 1986.
- [31] V. A. Solonnikov. Estimates for solutions of nonstationary Navier-Stokes systems. *Zap. Nauchn. Sem. LOMI*, 38, (1973), p.153-231; *J. Soviet Math.* 8, (1977), p. 467-529.
- [32] V. Valli and W. Zajaczkowski. Navier-Stokes equations for compressible fluids: global existence and qualitative properties of the solutions in the general case. *Commun. Math. Phys.*, 103, (1986) no 2, p. 259-296.