

Equilibria in reflexive Banach lattices with a continuum of agents[★]

Aloisio Araujo¹, V. Filipe Martins-da-Rocha², and Paulo K. Monteiro³

¹ Instituto Nacional de Matemática Pura e Aplicada, IMPA and EPGE/FGV,
Estrada Dona Castorina 110, CEP 22460, Rio de Janeiro, RJ, BRAZIL
(e-mail: aloisio@impa.br)

² Ceremade, Université Paris-IX Dauphine, Place du Marchal de Lattre de Tassigny,
75775 Paris Cedex 16, FRANCE (e-mail: martins@ceremade.dauphine.fr)

³ EPGE/FGV, Praia de Botafogo 190, 11 andar Botafogo, 22250-900 Rio de Janeiro, RJ, BRAZIL
(e-mail: PKLM@fgv.br)

Received: October 20, 2003; revised version: February 2, 2004

Summary. We consider exchange economies with a measure space of agents and for which the commodity space is a separable and reflexive Banach lattice. Under assumptions imposing uniform bounds on marginal rates of substitution, positive results on core-Walras equivalence were established in Rustichini–Yannelis [27] and Podczeck [25]. In this paper we prove that under similar assumptions on marginal rates of substitution, the set of competitive equilibria (and thus the core) is non-empty.

Keywords and Phrases: Competitive equilibria, Continuum of agents, Reflexive Banach lattice commodity spaces, Uniform properness.

JEL Classification Numbers: C62, D41, D51.

1 Introduction

We consider an exchange economy with infinitely many agents and infinitely many commodities. Infinite dimensional commodity spaces arise very naturally in economics, in particular in problems involving the allocation of resources over an infinite time horizon (e.g. an ℓ^p commodity space) or uncertainty about the possibly infinite number of states of nature (e.g. an $L^p([0, 1])$ commodity space). In our model, the commodity space will be a reflexive and separable Banach space. In the formulation of the Arrow–Debreu–McKenzie model of an exchange economy

* Thanks to Charalambos D. Aliprantis, Konrad Podczeck, Rabee Tourky, Nicholas C. Yannelis and an anonymous referee for helpful discussions and suggestions.

Correspondence to: V.F. Martins-da-Rocha

(Arrow–Debreu [5], McKenzie [22], Debreu [11]), a finite number of agents take prices as given. This formulation raises a conceptual difficulty: a finite number of agents should mean that individuals are able to exercise some influence, which contradicts the price-taking behavior assumption. To model perfectly competitive markets, we follow Aumann [6, 7] and Hildenbrand [14], who suggested to model the set of agents by a finite complete measure space. The insignificance of individual agents is thus captured by the idea of a set of zero measure.

In the literature dealing with large economies (infinitely many agents), two solution concepts are used: the competitive (Walrasian) equilibrium and the core. For the first concept, agents are assumed to take prices as given and they engage in the sale and purchase of commodities in order to maximize their utilities subject to their budgets. Agents trade freely in a decentralized market and this process results in allocations which equate supply with demand. The second concept allows for the possibility of cooperation among agents. They are allowed to bargain multilaterally which leads to an allocation of resources where it is not possible for any coalition of agents to redistribute their initial endowments among themselves in any way that makes each member of the coalition better off. Aumann [6] proved that in perfectly competitive economies (i.e. economies with an atomless finite measure space of agents) with finitely many commodities, the core coincides with the set of competitive equilibria. He also proved in [7], that the set of competitive equilibria (and thus the core) is non-empty. The core-Walras equivalence theorem was extended by Rustichini–Yannelis [27], to commodity spaces being separable Banach spaces.

In the framework of large square economies (i.e. with infinitely many agents and infinitely many commodities), there are several equilibrium existence results: Bewley [9], Khan–Yannelis [16], Podczeck [24] and Martins-da-Rocha [19] for separable Banach commodity spaces with an interior point in the positive cone; and Mas-Colell [21], Jones [15], Ostroy–Zame [23], Podczeck [24, 26] and Martins-da-Rocha [20] for economies with differentiated commodities.¹ To the best of our knowledge, this paper is the first to provide an equilibrium existence result for economies with a separable and reflexive Banach lattice (i.e. ℓ^p or $L^p([0, 1])$ for $1 < p < +\infty$). Under assumptions imposing uniform bounds on marginal rates of substitution, positive results on core-Walras equivalence were established in Rustichini–Yannelis [27] and Podczeck [25]. In this paper we prove that under similar assumptions on marginal rates of substitution, the set of competitive equilibria (and thus the core) is non-empty. More precisely, we provide two frameworks to prove the existence of competitive equilibria. In the first one, existence is proved under an assumption (borrowed from Zame [30] and Podczeck [25]) imposing the existence, at each state of nature, of uniform (over consumption) upper and lower bounds on marginal rates of substitution. In the second one, the commodity space is ℓ^p and existence is proved for preference relations represented by separable utility functions. But for this framework, we only require the existence of an upper bound on the marginal rates of substitution at the initial endowment (and not uniformly

¹ The commodity space is $M([0, 1])$ the space of Radon measures on $[0, 1]$ and the price space is $C([0, 1])$ the space of continuous functions on $[0, 1]$.

over consumption) and we require the existence of a uniform (over consumption) lower bound on the marginal rates of substitution at only one state of nature.

Recently, Tourky–Yannelis [28] showed that, when aggregation of individual commodity bundles is formalized in terms of the Bochner integral, given a non-separable Hilbert space E , and given any atomless measure space $(\Omega, \mathcal{A}, \mu)$, there is an economy with $(\Omega, \mathcal{A}, \mu)$ as space of agents and E as commodity space that has a non-empty core but does not have a competitive equilibrium. Contrasting with the positive results of Aumann [6, 7] and their extensions to separable Banach commodity spaces by Rustichini–Yannelis [27], the crucial condition to get these results is that there are "many more agents than commodities". More precisely, Podczeck [25] proved that the class $\{E\}$ of Banach spaces such that, under a list of "desirable assumptions", any atomless economy with commodity space E exhibits core-Walras equivalence² is exactly the class of Banach spaces that are separable. However, there is no characterization in the literature of a class of Banach spaces as those spaces in which the existence of competitive equilibria holds. In our framework, the commodity space is a separable Banach space, thus under "desirable assumptions" the core-Walras equivalence theorem is valid. But, the separability assumption of the commodity space is no more sufficient to get the existence of competitive equilibria. We introduce an additional assumption which requires a compatibility (Definition 2.2 and Assumption 3.4) between the geometry of the lattice ordering of the commodity space and initial endowments. When $L^2([0, 1])$ is endowed with the natural pointwise lattice ordering, we construct an economy satisfying a list of "desirable assumptions", but not satisfying the compatibility assumption and for which there are no competitive equilibria at all. It appears that for the issue of the existence of competitive equilibria, the topological way of measuring the cardinality of the number of markets introduced by Tourky–Yannelis [28] is not appropriate. It is the geometric structure of the lattice ordering that matters. In this paper, several examples of lattice ordering which satisfy the compatibility condition are given. For these examples, the positive cone has countably many extreme rays. However the number of extreme directions of the positive cone is not the appropriate way of measuring the cardinality of the number of markets since positive results for smooth positive cones³ are given in Martins-da-Rocha [20].

Following the approach used in Martins-da-Rocha [19, 20], our proof of the existence of an equilibrium is based on the discretization of the set of agents. We approximate the initial economy \mathcal{E} by a sequence of economies (\mathcal{E}^n) with finitely many agents. To each *finite* economy \mathcal{E}^n , we use the lattice structure of the commodity space and the properness assumptions on preferences to get the existence of a quasi-equilibrium (x^n, p^n) . The last step consists on proving that the sequence (x^n, p^n) converges to an equilibrium (x, p) . In order to apply a Fatou type lemma to the sequence of mappings (x^n) , we need a specific compatibility (Definition 2.2 and Assumption 3.4) between the geometry of the lattice ordering and initial endowments.

² When feasibility of allocation is defined in terms of the Bochner integral.

³ The commodity space is $M([0, 1])$ ordered by the natural pointwise positive cone.

The paper is organized as follows. In Section 2 we define the model of an economy with infinitely many agents and commodities and we set out the main definitions and notations. In Section 3 we give the list of assumptions that economies will be required to satisfy and we present the two existence results. The different assumptions on the marginal rates of substitution are discussed in Section 4. Finally, Section 5 is devoted to the proof of the two theorems.

2 The model

2.1 Preliminaries

Let E be a separable and reflexive Banach lattice.⁴ We denote by E^* the dual space of E , i.e. the space of all continuous linear functions from E into \mathbb{R} . If $x \in E$ and $p \in E^*$, the value $p(x)$ of p at x will often be denoted $\langle p, x \rangle$. We write $\|\cdot\|$ for both the norm of E and the dual norm of E^* . We write w for the weak topology $\sigma(E, E^*)$ on E , w^* for the weak-star topology $\sigma(E^*, E)$ on E^* , and s for the norm-topology. As usual, the ordering of E is denoted by \geq , and E_+ denotes the positive cone of E , i.e. $E_+ = \{x \in E : x \geq 0\}$. The dual space E^* will always be regarded as endowed with the dual ordering, i.e. $E_+^* = \{p \in E^* : p(x) \geq 0, \forall x \in E_+\}$. A vector $x \in E$ is said positive if $x \geq 0$, a linear functional $q \in E^*$ is said *strictly positive* if $q(x) > 0$ whenever x belongs to $E_+ \setminus \{0\}$. For $x, y \in E$ the expressions $x^+, x^-, |x|$ have the usual lattice theoretical meaning. Let τ be a topology on E . If $(C_n)_n$ is a sequence of subsets of E , the τ -*sequential upper limit* of $(C_n)_n$, is denoted $\tau\text{-ls}_n C_n$ and is defined by

$$\tau\text{-ls}_n C_n := \{x \in E : x = \tau\text{-}\lim_k x^k, \quad x^k \in C_{n(k)}\}$$

where $(C_{n(k)})_k$ is a subsequence of $(C_n)_n$.

The Borel σ -algebra of E for the norm-topology or for the weak-topology coincide and is denoted by \mathcal{B} . Let $(\Omega, \mathcal{A}, \mu)$ be a complete finite positive measure space. A correspondence F from Ω to E is said to be graph measurable if $\{(a, x) \in \Omega \times E : x \in F(a)\}$ belongs to $\mathcal{A} \otimes \mathcal{B}$. A correspondence P from Ω to $E \times E$ is said to be graph measurable if $\{(a, x, z) \in \Omega \times E \times E : (x, z) \in P(a)\}$ belongs to $\mathcal{A} \otimes \mathcal{B} \otimes \mathcal{B}$. A mapping s from Ω to E is simple if there exist x_1, x_2, \dots, x_n in E and A_1, A_2, \dots, A_n in \mathcal{A} such that $s = \sum_{i=1}^n x_i \chi_{A_i}$ where $\chi_{A_i}(a) = 1$ if $a \in A_i$ and $\chi_{A_i}(a) = 0$ if $a \notin A_i$. A mapping $x : \Omega \rightarrow E$ is Bochner measurable if there is a sequence of simple mappings $s_n : \Omega \rightarrow E$ such that $\lim_n \|s_n(a) - x(a)\| = 0$ almost every where. Since E is separable, we know from Pettis' measurability theorem (see [12, Theorem II.1.2, p.42]) that a mapping $x : \Omega \rightarrow E$ is Bochner measurable if and only if for each $B \in \mathcal{B}$, $x^{-1}(B) := \{a \in \Omega : x(a) \in B\}$ belongs to \mathcal{A} . A Bochner measurable mapping x from Ω to E is Bochner integrable if there is a sequence of simple mappings $s_n : \Omega \rightarrow E$ such that $\lim_n \int_\Omega \|s_n(a) - x(a)\| d\mu(a) = 0$. For each measurable set A in \mathcal{A} , we denote by $\int_A x d\mu$ the limit $\lim_n \int_A s_n d\mu$. It can easily be shown (see [12, p.45]) that a

⁴ We refer to Aliprantis–Border [2] for definitions.

Bochner measurable mapping x is Bochner integrable if and only if the mapping $a \mapsto \|x(a)\|$ is integrable. In particular $\|\int_A x d\mu\| \leq \int_A \|x(a)\| d\mu(a)$.

2.2 Fatou's cone

We define hereafter a class of lattice orderings which will enable us to apply a Fatou type lemma.

Definition 2.1. Let E be a Banach lattice, a functional ρ from E into $[-\infty, +\infty]$ is a *positive extended linear functional* if

- (i) the space $E^\rho = \{x \in E : \rho(x) \in \mathbb{R}\}$ is a vector subspace of E ,
- (ii) the restriction of ρ to E^ρ is linear and,
- (iii) the functional ρ is positive, i.e. for any $y \geq x \geq 0$, we have $\rho(y) \geq \rho(x) \geq 0$.

Definition 2.2. Let E be a Banach lattice ordered by a positive cone E_+ and let $e : \Omega \rightarrow E_+$ be a Bochner integrable mapping. The cone E_+ is a *Fatou's cone relatively to e* if there exists a positive extended linear functional ρ such that

- (a) for every x in E_+ , $\|x\| \leq \rho(x)$,
- (b) the function $a \mapsto \rho[e(a)]$ from Ω to \mathbb{R}_+ is integrable.

Example 2.3. Take $E = \ell^p$ for any $1 < p < +\infty$, and $E_+ = \ell^p_+$ the natural pointwise positive cone.⁵ Then, for any Bochner integrable mapping $e : \Omega \rightarrow \ell^p_+$, if the function⁶ $a \mapsto \|e(a)\|_1$ is integrable, then E_+ is a Fatou's cone relatively to e .

Example 2.4. Take $E = L^2([0, 1])$ and let $(b_n)_n$ be an Hilbert basis of E . Let

$$E_+ = \{x \in E : \forall n \in \mathbb{N}, \langle x, b_n \rangle \geq 0\},$$

then E is a Banach lattice. Moreover, if $e : \Omega \rightarrow E_+$ is a Bochner integrable mapping such that the function

$$a \mapsto \sum_n \langle e(a), b_n \rangle$$

is integrable, then E_+ is a Fatou's cone relatively to e .

Remark 2.5. In the above two examples, the positive cone has countably many extreme rays. Let (T, \mathcal{T}, σ) be an atomless measure space and $1 < p < +\infty$. We prove in Appendix B.2 that when $E = L^p(T, \mathcal{T}, \sigma)$ is ordered by the pointwise "smooth" positive cone $E_+ = L^p(T, \mathcal{T}, \sigma)_+$, then for every Bochner integrable mapping $e : \Omega \rightarrow E_+$ with $\int_\Omega e d\mu > 0$, the cone E_+ is not a Fatou's cone relatively to e .

⁵ The natural pointwise positive cone of ℓ^p is $\ell^p_+ = \{x = (x_n) \in \ell^p : \forall n \in \mathbb{N}, x_n \geq 0\}$.

⁶ If $x = (x_n)$ belongs to ℓ^p , we let $\|x\|_1 = \sum_n |x_n|$.

2.3 The model

An economy \mathcal{E} is a list

$$\mathcal{E} = ((\Omega, \mathcal{A}, \mu), E, X, \succeq, e),$$

where X is a correspondence from Ω to E , \succeq is a correspondence from Ω to $E \times E$ and e is a mapping from Ω to E . The space of agents is $(\Omega, \mathcal{A}, \mu)$, a complete finite positive measure space. The commodity space is E . For each agent $a \in \Omega$, the consumption set is $X(a)$, the initial endowment is $e(a) \in E$ and the preference/indifference relation is $\succeq_a \subset X(a) \times X(a)$, a reflexive binary relation on $X(a)$.

We define the correspondence⁷ $P_a : X(a) \rightarrow X(a)$ by $P_a(x) = \{x' \in X(a) : x' \succ_a x\}$. In particular, if $x \in X(a)$ is a consumption bundle, the set $P_a(x)$ is the set of consumption bundles strictly preferred to x by agent a . We let P be the correspondence from Ω to $E \times E$ defined for each $a \in \Omega$ by $P(a) = \{(x, x') \in X(a) \times X(a) : x' \succ_a x\}$.

The set of allocations (or plans) of the economy is the set $S^1(X)$ of Bochner integrable selections of X , i.e. $S^1(X)$ is the set of mappings x from Ω to E which are Bochner integrable and which satisfies $x(a) \in X(a)$ for almost every $a \in \Omega$. An allocation $x \in S^1(X)$ is feasible if

$$\int_{\Omega} x d\mu = \int_{\Omega} e d\mu.$$

We assume that the mapping $e : \Omega \rightarrow E$ is a Bochner integrable mapping and we denote by $\omega := \int_{\Omega} e d\mu$ the aggregate initial endowment.

Definition 2.6. A pair (x, p) consisting of a feasible allocation x and a non-zero price p is said to be a *competitive equilibrium* if for almost every $a \in \Omega$, $\langle p, x(a) \rangle = \langle p, e(a) \rangle$, and $z \in P_a(x(a))$ implies $\langle p, z \rangle > \langle p, x(a) \rangle$.

3 Existence of a competitive equilibrium

We will maintain in this paper the following assumptions on the economy \mathcal{E} .

Assumption 3.1. For each $a \in \Omega$,

- (i) the consumption set is $X(a) = E_+$;
- (ii) the initial endowment is not zero, i.e. $e(a) > 0$;
- (iii) \succeq_a is reflexive, transitive and complete;
- (iv) \succeq_a is strictly monotone, i.e. for each $x \in X(a)$, if $z > x$ then $z \succ_a x$.

⁷ As usual, $y \succ_a x$ means $[y \succeq_a x$ and $x \not\succeq_a y]$. Note that the binary relation \succ_a coincide with the graph of the correspondence P_a .

Assumption 3.2. For each $a \in \Omega$, for every $x \in X(a)$,

- (i) the sets $P_a(x)$ and $P_a^{-1}(x) = \{z \in X(a) : x \succ_a z\}$ are norm-open in $X(a)$;
- (ii) the set $\{z \in X(a) : z \succeq_a x\}$ is convex.

Assumption 3.3. The correspondence P is graph measurable.

Assumption 3.4. The positive cone E_+ is a Fatou's cone relatively to e .

Remark 3.1. Assumptions 3.1–3.3 are standard in the literature dealing with exchange economies with finitely or infinitely many agents. We will see that in our framework, we can not dispense with Assumption 3.4.

We provide hereafter two frameworks to prove the existence of competitive equilibria. In the first one, existence is proved under an assumption imposing upper and lower uniform (over agents and consumption) bounds on marginal rates of substitution. In the second one, existence is proved for preference relations represented by separable utility functions defined on ℓ^p , but the assumption on the marginal rates of substitution required for the existence is weaker. We only require a uniform (over agents only) upper bound on the marginal rates of substitution at the initial endowment and a uniform (over agents and consumption) lower bound on the marginal rates of substitution at only one state of nature.

3.1 The general case

In this section, we consider economies with general preference relations. The following requirement is borrowed from Zame [30]. It is discussed in Section 4.

Definition 3.2. The preference relations (\succeq_a) are said to be *strong-uniformly proper*, if there exist strictly positive prices α and β in E^* with $\alpha \leq \beta$ and such that for every $a \in \Omega$, whenever $x, u, v \in E_+$ satisfy $v \leq x$ and $\langle \alpha, u \rangle > \langle \beta, v \rangle$ then $x - v + u \succ_a x$.

An economy \mathcal{E} is said strong-uniformly proper if it has strong-uniformly proper preference relations.

We can now state our first result for economies with general preference relations.

Theorem 3.3. *If the economy \mathcal{E} is strong-uniformly proper then there exists a competitive equilibrium.*

Remark 3.4. The strong-uniform properness assumption was already used in Zame [30]. Podczeck in [26] proved the equivalence between the core and the set of competitive equilibria under this assumption. Note that Rustichini–Yannelis [27] also proved the equivalence between the core and the set of competitive equilibria under another properness assumption.

Assumption 3.4 is unusual. Following Zame [30] we provide hereafter two examples of a strong-uniformly proper economies satisfying Assumptions 3.1–3.3 and not satisfying Assumption 3.4. For these economies the set of competitive equilibria is empty.

Counterexample 3.5. Consider the economy \mathcal{E} where $\Omega = [0, 1]$, \mathcal{A} is the Lebesgue σ -algebra and μ is the Lebesgue measure. The commodity space E is ℓ^p for $1 < p < +\infty$, ordered by the pointwise positive cone ℓ^p_+ . For each trader $a \in [0, 1]$, the utility function u_a is defined by

$$u_a(x) = \sum_{n \in \mathbb{N}} (2 + a)^{-n} x_n$$

and the initial endowment is defined by $e(a) = (1, 1/2, 1/3, \dots, 1/n, \dots)$. The economy \mathcal{E} is strong-uniformly proper, it satisfies Assumptions 3.1–3.3 but not Assumption 3.4. It is proved in Zame [30] that \mathcal{E} has no competitive equilibrium.

Counterexample 3.6. Consider the economy \mathcal{E} where $\Omega = [0, 1]$, \mathcal{A} is the Lebesgue σ -algebra and μ is the Lebesgue measure. The commodity space E is $L^p([0, 1], \mu)$ where $1 < p < +\infty$ ordered by the "smooth" pointwise positive cone $L^p([0, 1], \mu)_+$. For each trader $a \in [0, 1]$, the utility function u_a is defined by

$$u_a(x) = \int_{[0,1]} q_a(t)x(t)\mu(dt)$$

and the initial endowment is $e(a) : t \mapsto 1$. For each $a \in (0, 1]$, the function q_a is defined by

$$q_a(t) = \begin{cases} \frac{1}{2} + \frac{t}{2a} & \text{if } 0 \leq t \leq a \\ \frac{a-2}{2(a-1)} + \frac{t}{2(a-1)} & \text{if } a \leq t \leq 1. \end{cases}$$

The economy \mathcal{E} is strong-uniformly proper, it satisfies Assumptions 3.1–3.3 but not Assumption 3.4. It is proved in Zame [30] that \mathcal{E} has no competitive equilibrium.

3.2 The separable case

In this section, we consider economies with the space ℓ^p as the commodity space and for which preference relations are represented by separable utility functions. For each $1 < p < +\infty$, we denote by ℓ^p the real vector space of sequences $x = (x_k)_k$ in $\mathbb{R}^{\mathbb{N}}$ such that $\lim_n \sum_{k=0}^n |x_k|^p < \infty$ and we denote by $\|x\|_p = (\sum_{k \in \mathbb{N}} |x_k|^p)^{1/p}$. We denote by ℓ^p_+ the natural positive cone defined by $x \in \ell^p_+$ if and only if $x_k \geq 0$ for each $k \in \mathbb{N}$. The space ℓ^p endowed with the norm $\|\cdot\|_p$ and the positive cone ℓ^p_+ is a reflexive and separable Banach lattice whose dual is ℓ^q where $1 < q < +\infty$ is defined by $1/p + 1/q = 1$.

Definition 3.7. A utility function $u : \ell^p_+ \rightarrow \mathbb{R}$ is called separable if there exists for each n , a function $v_n : [0, +\infty) \rightarrow \mathbb{R}$ concave and strictly increasing such that

$$\forall x \in \ell^p_+, \quad u(x) = \sum_{n \in \mathbb{N}} v_n(x_n).$$

The function $v = (v_n)$ is called the kernel of u . The left derivative of v_n in $t > 0$ is noted $v_n^-(t)$ and the right derivative is denoted $v_n^+(t)$. If $x \in \ell_+^p$ then we note $v^-(x) := (v_n^-(x_n))_n$. For $x \in \ell_+^p$ we define $S(x) = \{h \in \ell^p : \exists t > 0, x+th \geq 0\}$ and $I(x) = S(x) \cap -S(x)$. We define $u'(x) \cdot h = \lim_{r \rightarrow 0} (1/r)[u(x+rh) - u(x)]$ for each $x \in \ell_+^p$ and each $h \in S(x)$. Note that if $h \geq 0$ then $u'(x) \cdot h = \sum_n v_n^+(x_n)h_n$.

For economies with separable utility functions, a weaker condition than the uniform properness will be sufficient to prove the existence of competitive equilibria.

Definition 3.8. An economy \mathcal{E} is said *separably proper* if for each agent $a \in \Omega$, the preference relation \succ_a is represented⁸ by a separable utility function u_a which kernel is denoted v_a and if there exists a measurable set $\Omega' \in \mathcal{A}$ of full measure,⁹ satisfying the following conditions.

(a) There exists $\beta \in \ell^q$ such that for each $a \in \Omega'$,

$$v_a^-(e(a)) \leq \beta.$$

(b) There exists $k \in \mathbb{N}$ and $\alpha_k > 0$ such that $\omega_k \alpha_k > 0$ and for each $a \in \Omega'$,

$$0 < \alpha_k \leq \inf\{v_{a,k}^+(t) : t \geq 0\} = \lim_{t \rightarrow +\infty} v_{a,k}^+(t).$$

We can now state our second existence result for economies with preference relations represented by utility functions.

Theorem 3.9. *If the economy \mathcal{E} is separably proper then there exists a competitive equilibrium.*

The two properness conditions are not comparable. Obviously, not all strong-uniformly proper economies are separably proper. Moreover, we provide hereafter an example of an economy which is separably proper but not strong-uniformly proper.

Example 3.10. Consider the economy \mathcal{E} where $\Omega = [0, 1]$, \mathcal{A} is the Lebesgue σ -algebra and μ is the Lebesgue measure. For each trader $a \in [0, 1]$, the consumption set coincide with ℓ_+^p , the utility function u_a is defined by

$$u_a(x) = x_0 + \sum_{n \geq 1} \frac{1 - \exp(-ax_n n^2)}{n^2}$$

and the initial endowment is defined by $e(a) = (1, 1/2, 1/3, \dots, 1/n, \dots)$. Following Example 4.6 the economy \mathcal{E} satisfies Assumptions 3.1–3.4. Moreover this economy is separably proper but not strong-uniformly proper.

4 Proper economies

We discuss in this section the notions of properness used in Theorem 3.3 and Theorem 3.9.

⁸ That is $x' \succ_a x$ if and only if $u_a(x') > u_a(x)$.

⁹ That is $\mu(\Omega \setminus \Omega') = 0$.

4.1 Strong-uniformly proper economies

We recall that the preference relations (\succeq_a) are said to be strong-uniformly proper, if there exist strictly positive prices α and β in E_+ with $\alpha \leq \beta$ and such that for every $a \in \Omega$, whenever $x, u, v \in E_+$ satisfy $v \leq x$ and $\langle \alpha, u \rangle > \langle \beta, v \rangle$ then $x - v + u \succ_a x$. This properness condition is borrowed from Zame [30]. Note that this is a requirement on preferences that is uniform over agents as well as over consumption. We refer to Zame [30] for a discussion of this condition as well as for corresponding examples. Following Podczeck [25], it may be seen that if for each $a \in \Omega$, $\{y \in X(a) : y \succeq_a x\}$ is convex then uniform properness is equivalent to the following statement: *There are strictly positive prices $\alpha, \beta \in E^*$, such that given any $a \in \Omega$ and $x \in X(a)$ there is a price p in the order interval $[\alpha, \beta]$ such that $\langle p, x \rangle \leq \langle p, y \rangle$ for all $y \in X(a)$ with $y \succeq_a x$.* Since supporting prices are measures of marginal rates of substitution, the strong-uniform properness assumption is a condition that puts strong bounds on these rates.

We recall the notion of uniform properness introduced by Yannelis–Zame [29] for economies with finitely many agents.

Definition 4.1. The preference relations (\succeq_a) are said to be *v-uniformly proper* with $v \in E$, if there exists a norm-open 0-neighborhood $V \subset E$ such that for each $a \in \Omega$, for each $x \in E_+$, $(x + \Gamma) \cap E_+ \subset P_a(x)$ where $\Gamma = \cup_{t>0} t(v + V)$.

Remark 4.2. The strong-uniform properness assumption on the preference relations implies that

$$\forall a \in \Omega, \quad \forall x \in E_+, \quad (x + \Gamma) \cap E_+ \subset P_a(x),$$

where Γ is the convex and norm-open cone defined by $\Gamma = \{x \in E : \alpha(x^+) > \beta(x^-)\}$.

Example 4.3. Consider the case of positive separable utility functions $u_a : \ell_+^p \rightarrow \mathbb{R}$, defined by the formula $u_a(x) = \sum_n v_{a,n}(x_n)$ where for each n , the function $v_{a,n} : [0, +\infty) \rightarrow \mathbb{R}$ is continuous, the derivative $v'_{a,n}(t)$ exists for each $t > 0$. Suppose that there exist α and β two strictly positive functionals in ℓ^q such that

$$\forall a \in \Omega, \quad \forall t > 0, \quad \alpha_n \leq v'_{a,n}(t) \leq \beta_n.$$

Then the preference relations defined by the utility functions $(u_a)_{a \in \Omega}$ are strong-uniformly proper. Indeed, let $x, y, z \in \ell_+^p$ satisfying $y \leq x$ and $\alpha(z) > \beta(y)$. Using the mean value theorem, we see that for each n there exists $t_n > 0$ such that

$$v_{a,n}(x_n - y_n + z_n) - v_{a,n}(x_n) = v'_{a,n}(t_n)[z_n - y_n].$$

But $v'_{a,n}(t_n)[z_n - y_n] \geq \alpha_n z_n - \beta_n y_n$, in particular

$$u(x - y + z) - u(x) \geq \alpha(z) - \beta(y) > 0.$$

We refer to Araujo–Monteiro [3], Le Van [17] and Aliprantis [1] for precisions about proper conditions for separable utility functions.

4.2 Separably proper economies

Following Aliprantis [1], we introduce the following notion of separable utility function.

Definition 4.4. A separable utility function $u : \ell_+^p \rightarrow \mathbb{R}$, where $u(x) = \sum_n v_n(x_n)$, is said to be *rational* if for each $n \in \mathbb{N}$,

- (a) $v_n(0) = 0$;
- (b) v_n is positive, continuous and concave on $[0, +\infty)$; and
- (c) v_n is differentiable on $(0, +\infty)$ with $v'_n(t) > 0$ for each $t > 0$.

Now let u be a rational separable utility function. The components of the lower and upper gradient sequences $\underline{v}' = (\underline{v}'_1, \underline{v}'_2, \dots)$ and $\bar{v}' = (\bar{v}'_1, \bar{v}'_2, \dots)$ are given by

$$\underline{v}'_n = \lim_{t \rightarrow +\infty} v'_n(t) \quad \text{and} \quad \bar{v}'_n = \lim_{t \rightarrow 0} v'_n(t).$$

Following Aliprantis [1, Theorem 6.7], we have the following result.

Proposition 4.5. Let $u : \ell_+^p \rightarrow \mathbb{R}$ be rational utility function given by $u(x) = \sum_n v_n(x_n)$. If the preference relations represented by u are ω -uniformly proper for some $\omega \in \ell^p$ strictly positive, then

- (a) the lower gradient \underline{v}' is non-zero and belongs to ℓ_+^q ; and
- (b) there exists some $k \in \mathbb{N}$ such that

$$(0, 0, \dots, 0, \bar{v}'_k, \bar{v}'_{k+1}, \dots) \in \ell_+^q.$$

It follows that if \mathcal{E} is an economy with rational separable utility function such that \mathcal{E} is ω -uniformly proper and ω is strictly positive, then \mathcal{E} is separably proper. We provide hereafter an example of a rational separable utility function which is separably proper but which is not uniformly proper.

Example 4.6. Consider the rational separable utility function $u : \ell_+^p \rightarrow \mathbb{R}$ defined by

$$v_0(t) = t \quad \text{and} \quad \forall n \geq 1, \quad v_n(t) = \frac{1 - \exp(-tn^2)}{n^2}.$$

For each $n \geq 1$, $v'_n(t) = \exp(-n^2t)$. It follows that

$$\underline{v}' = (1, 0, 0, \dots) \quad \text{and} \quad \bar{v}' = (1, 1, 1, \dots).$$

It follows that u is not uniformly proper. However if $e = (e_n)_n$ is defined by $e_0 = 1$ and for each $n \geq 1$ by $e_n = 1/n$ then

$$v'(e) = (e^{-n})_n \in \ell_+^p.$$

Hence u is separably proper.

5 Proof of Theorem 3.3 and Theorem 3.9

Since E is a separable and reflexive Banach space, it follows that E^* is norm-separable. Let $(p_i)_{i \in \mathbb{N}}$ be a norm-dense sequence in the closed unit ball of E^* and define for each x, y in E ,

$$d(x, y) = \sum_{i \in \mathbb{N}} \frac{|\langle p_i, x - y \rangle|}{2^i}.$$

The topology defined by this distance coincide with the w -topology on norm-bounded subsets of E . Moreover, the d -topology is separable and the Borel σ -algebra generated by d coincide with the Borel σ -algebra \mathcal{B} generated by the norm-topology and the w -topology.

Let \mathcal{E} be an economy satisfying Assumptions 3.1–3.4. Suppose that \mathcal{E} is either strong-uniformly proper or separably proper. The correspondence X is graph measurable. Applying Theorem B.1, there exists a sequence $(f_k)_k$ of measurable selections of X such that for each $a \in \Omega$, $X(a) = s\text{-cl}\{f_k(a) : k \in \mathbb{N}\}$. For every $k \in \mathbb{N}$, we let R_k be the correspondence from Ω into E , defined by $R_k(a) = \{x \in E_+ : x \succeq_a f_k(a)\}$. For each $\nu \in \mathbb{N}$, we let $X_\nu : a \mapsto X_\nu(a) := X(a) \cap \nu B$ and $R_{k,\nu} : a \mapsto R_{k,\nu}(a) := R_k(a) \cap \nu B$, where B is the closed unit ball in E .

Claim 5.1. *There exists¹⁰ a sequence $(\sigma^n)_n$ of measurable partitions $\sigma^n = (A_i^n)_{i \in S^n}$ of (Ω, \mathcal{A}) , and a sequence $(A^n)_n$ of finite sets $A^n = \{a_i^n : i \in S^n\}$ subordinated to the measurable partition σ^n , satisfying¹¹ for each $a \in \Omega$,*

- (i) $\lim_n \|e^n(a) - e(a)\| = 0$ and $\forall k \in \mathbb{N}, \lim_n \|f_k^n(a) - f_k(a)\| = 0$;
- (ii) for each $\nu \in \mathbb{N}$, for each sequence $(x^n)_n$ of E , d -converging to $x \in E$ and for every $k \in \mathbb{N}$,

$$\lim_n d(x^n, R_{k,\nu}^n(a)) = d(x, R_{k,\nu}(a));$$

- (iii) if we pose¹² $g(a) := \rho[e(a)]$ then g is an integrable function satisfying

$$\forall n \in \mathbb{N}, \quad \rho[e^n(a)] \leq 1 + g(a).$$

Proof. If f is a function from Ω to E , then we let $\{f(\cdot)\}$ be the correspondence from Ω into E defined for each $a \in \Omega$, by $\{f(\cdot)\}(a) := \{f(a)\}$. Note that if f is measurable then f is Bochner integrable if and only if $\|f(\cdot)\| : a \mapsto \|f(a)\|$ from Ω to \mathbb{R}_+ is integrable.

Let $Z := E \times E$ and consider the following distance δ on Z defined for each $x = (x_1, x_2)$ and $y = (y_1, y_2)$ in Z by

$$\delta(x, y) := \|x_1 - y_1\| + d(x_2, y_2).$$

¹⁰ We refer to Appendix A.2 for definitions and notations.

¹¹ Following notations of Section B.2, if f is function from Ω to E , then for each n , $\{f(\cdot)\}^n = \{f^n(\cdot)\}$.

¹² The functional ρ is given by Assumption 3.4.

The metric space (Z, δ) is complete and separable. Let $\mathcal{G} := \{\rho[e(\cdot)]\}$ and¹³

$$\mathcal{F} := \{\{e(\cdot)\} \times \{0\}, \{f_k(\cdot)\} \times \{0\}, \{0\} \times R_k : k \in \mathbb{N}\}.$$

Now apply Theorem B.6.

We construct a sequence $(\mathcal{E}^n)_n$ of economies with finitely many consumers converging to \mathcal{E} . We let

$$\mathcal{E}^n := \left((I^n, 2^{I^n}, \sigma), E, X^n, \succ^n, e^n \right),$$

where $I^n = \{i \in S^n : \mu(A_i^n) \neq 0\}$ is the finite set of consumers; σ is the counting measure on I^n ; for each agent $i \in I^n$, the consumption set is defined by $X_i^n := \mu(A_i^n)X(a_i^n) = E_+$, the initial endowment is defined by $e_i^n := \mu(A_i^n)e(a_i^n)$ and the preference relation is defined by $x' \succ_i^n x$ if and only if $(x'/\mu(A_i^n)) \succ_{a_i^n}(x/\mu(A_i^n))$. In particular, the correspondence of strictly preferred bundles P_i^n from X_i^n to X_i^n is defined by $P_i^n(x) = \mu(A_i^n)P_{a_i^n}(x/\mu(A_i^n))$, for each $x \in X_i^n$.

Claim 5.2. *There exists a feasible allocation $(x_i^n)_{i \in I^n}$ for the finite economy \mathcal{E}^n , a non-zero price p^n and a w^* -compact set $K \subset E^*$ such that*

- (1) *for each $i \in I^n$, $\langle p^n, x_i^n \rangle = \langle p^n, e_i^n \rangle$, and $z \in P_i^n(x_i^n)$ implies $\langle p^n, z \rangle \geq \langle p^n, x_i^n \rangle$; and*
- (2) *$p^n \in K$ with $\langle p^n, \omega \rangle = 1$.*

Proof. If the economy \mathcal{E} is strong-uniformly proper then each economy \mathcal{E}^n satisfies the assumptions of Theorem A.1. In particular if we let $K := \{q \in E^* : \langle q, \omega \rangle = 1 \text{ and } \langle q, \Gamma \rangle > 0\}$, where $\Gamma = \{x \in E : \langle \alpha, x^+ \rangle > \langle \beta, x^- \rangle\}$, then Claim 5.2 is proved.

If the economy \mathcal{E} is separably proper then for each n large enough, $\alpha_k \omega_k^n > 0$. Hence the economy \mathcal{E}^n satisfies the assumptions of Theorem A.3. In particular for each n , $\|p^n\|_q \leq (1/\alpha_k \omega_k^n) \|\beta\|_q$. Since $(\omega_k^n)_n$ is norm-convergent to ω^k , it follows that the sequence $(p^n)_n$ lies in a norm-bounded set $K \subset \ell^q$. In particular, Claim 5.2 is proved. □

If we denote by x^n and e^n the Bochner integrable mappings defined by

$$x^n := \sum_{i \in I^n} x_i^n \chi_{A_i^n} \quad \text{and} \quad e^n := \sum_{i \in I^n} e_i^n \chi_{A_i^n}$$

then

$$(5.1) \quad \int_{\Omega} x^n d\mu = \int_{\Omega} e^n d\mu$$

$$\text{for a.e. } a \in \Omega, \quad \langle p^n, x^n(a) \rangle = \langle p^n, e^n(a) \rangle$$

$$(5.2) \quad \text{and } z \in P_a^n(x^n(a)) \Rightarrow \langle p^n, z \rangle \geq \langle p^n, e^n(a) \rangle$$

$$(5.3) \quad p^n \in K \quad \text{and} \quad \langle p^n, \omega \rangle = 1.$$

¹³ If F and G are two correspondences from Ω to E , then we let $F \times G$ be the correspondence from Ω to $E \times E$, defined for each $a \in \Omega$ by $(F \times G)(a) = F(a) \times G(a)$.

The set K is w^* -compact. Since E is norm-separable, passing to a subsequence if necessary, we can suppose that $(p^n)_n$ w^* -converge to a non-zero price $p \in E^*$ which satisfies $\langle p, \omega \rangle = 1$.

Now we want to apply a Fatou’s Lemma to the sequence (x^n) . It is only at this step that we need Assumption 3.4. For each $z \in E_+$, $\|z\| \leq \rho(z)$; and for each $y \in E_+$, $\rho(z + y) = \rho(z) + \rho(y)$. Hence from (5.1), we have

$$\int_{\Omega} \|x^n(a)\| d\mu(a) \leq \int_{\Omega} \rho[x^n(a)]d\mu(a) = \int_{\Omega} \rho[e^n(a)]d\mu(a).$$

Applying Claim 5.1, the sequence of Bochner integrable mappings $(x^n)_n$ is mean norm-bounded, i.e.

$$\sup_n \int_{\Omega} \|x^n(a)\| d\mu(a) < +\infty.$$

Passing to a subsequence if necessary, we can suppose that $w\text{-}\lim_n \int_{\Omega} x^n d\mu$ exists in E . Applying Fatou’s Lemma (Theorem C.1) of Cornet–Martins-da-Rocha [10], there exists a Bochner integrable mapping x from Ω to E such that

$$(5.4) \quad \int_{\Omega} x d\mu \leq w\text{-}\lim_n \int_{\Omega} x^n d\mu$$

$$(5.5) \quad x(a) \in \overline{\text{co}} w\text{-}\text{ls}_n \{x_n(a)\} \quad \text{a.e.}$$

Claim 5.3. *The Bochner integrable mapping x satisfies $\int_{\Omega} x d\mu \leq \int_{\Omega} e d\mu$.*

Proof. This is a direct consequence of Claim 5.1, (5.1) and (5.4). □

Claim 5.4. *For almost every $a \in \Omega$, $x(a) \in E_+$ and $z \in P_a(x(a))$ implies $\langle p, z \rangle \geq \langle p, e(a) \rangle$.*

Proof. Consider $\Omega_0 = \bigcup_{n \in \mathbb{N}} \Omega \setminus (\bigcup_{i \in I^n} A_i^n)$, then $\mu(\Omega_0) = 0$. Let Ω' be a measurable subset of $\Omega \setminus \Omega_0$ with $\mu(\Omega \setminus \Omega') = 0$ and such that all *almost everywhere* assumptions and properties are satisfied for each $a \in \Omega'$.

Since $X(a) = E_+$ is closed convex, we have that for each $a \in \Omega'$, $x(a) \in X(a)$. We will now prove that for each $a \in \Omega'$, if $z \in P_a(x(a))$ then $\langle p, z \rangle \geq \langle p, e(a) \rangle$. Let $a \in \Omega'$ and let $z \in P_a(x(a))$. Since $E_+ = s\text{-cl}\{f_k(a) : k \in \mathbb{N}\}$, we can suppose (extracting a subsequence if necessary) that $(f_k(a))_k$ is norm-convergent to z . But $P_a(x(a))$ is norm-open in E_+ , thus there exists $k_0 \in \mathbb{N}$, such that for each $k \geq k_0$, $f_k(a) \in P_a(x(a))$. To prove that $\langle p, z \rangle \geq \langle p, e(a) \rangle$, it is sufficient to prove that for each k large enough, $\langle p, f_k(a) \rangle \geq \langle p, e(a) \rangle$. Now, let $k \geq k_0$.

Claim 5.5. *There exists an increasing function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ such that*

$$\forall n \in \mathbb{N}, \quad f_k^{\varphi(n)}(a) \in P_a^{\varphi(n)} \left(x^{\varphi(n)}(a) \right).$$

Proof. Suppose that for each increasing function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$, there exists an increasing function $\phi : \mathbb{N} \rightarrow \mathbb{N}$, such that:

$$\forall n \in \mathbb{N}, \quad x^{\varphi \circ \phi(n)}(a) \in R_k^{\varphi \circ \phi(n)}(a).$$

Let $\ell \in w\text{-ls}\{x^n(a) : n \in \mathbb{N}\}$, then there exists a subsequence $(x^{\varphi(n)}(a))_n$ w -converging to ℓ . In particular $(x^{\varphi \circ \phi(n)}(a))_n$ is w -convergent to ℓ . It follows that there exists $\nu > 0$ such that for each n , $x^{\varphi \circ \phi(n)}(a)$ belongs to νB . In particular, $d(x^{\varphi \circ \phi(n)}(a), R_k^{\varphi \circ \phi(n)}(a)) = 0$. Applying Claim 5.1, it follows that $d(\ell, R_{k,\nu}(a)) = 0$. Since $R_{k,\nu}(a)$ is w -closed and d coincide with w on νB , we have that $\ell \in R_k(a)$. Thus $w\text{-ls}\{x^n(a)\} \subset R_k(a)$, and under Assumption 3.2, this implies that $\overline{w\text{-ls}_n\{x_n(a)\}} \subset R_k(a)$. It follows that $x(a) \in R_k(a)$, i.e. $f_k(a) \notin P_a(x(a))$: contradiction. \square

With Claim 5.5 and (5.2), for each n , $\langle p^{\varphi(n)}, f_k^{\varphi(n)}(a) \rangle \geq \langle p^{\varphi(n)}, e^{\varphi(n)}(a) \rangle$. Passing to the limit, we get that $\langle p, f_k(a) \rangle \geq \langle p, e(a) \rangle$. \square

Now let \bar{x} be the Bochner integrable mapping from Ω to ℓ^p defined by

$$\forall a \in \Omega, \quad \bar{x}(a) = x(a) + (1/\mu(\Omega)) \int_{\Omega} (e - x) d\mu.$$

Claim 5.6. *The pair (\bar{x}, p) is an equilibrium of \mathcal{E} .*

Proof. Since $\int_{\Omega} (e - x) d\mu \geq 0$, Assumption 3.1 implies that $\bar{x}(a) \in E_+$ and $P_a(\bar{x}(a)) \subset P_a(x(a))$. In particular the allocation \bar{x} is feasible and for each $a \in \Omega'$, if $z \in P_a(\bar{x}(a))$ then $\langle p, z \rangle \geq \langle p, e(a) \rangle$. Since \succ_a is monotone, it follows that $\bar{x}(a)$ belongs to the norm-closure of $P_a(\bar{x}(a))$, in particular $\langle p, \bar{x}(a) \rangle \geq \langle p, e(a) \rangle$. But $\int_{\Omega} \bar{x} d\mu = \int_{\Omega} e d\mu$, it follows that for almost every $a \in \Omega$, $\langle p, \bar{x}(a) \rangle = \langle p, e(a) \rangle$. To prove that (\bar{x}, p) is an equilibrium, it is now sufficient to prove that for almost every $a \in \Omega$,

$$\inf\{\langle p, z \rangle : z \in E_+\} < \langle p, e(a) \rangle.$$

Let $B := \{a \in \Omega' : \langle p, e(a) \rangle > 0\}$. The set B is measurable and since $\langle p, \omega \rangle = 1$, $\mu(B) \neq 0$. Now for each $a \in B$, $\inf\{\langle p, z \rangle : z \in E_+\} < \langle p, e(a) \rangle$ and $z \in P_a(\bar{x}(a)) \Rightarrow \langle p, z \rangle \geq \langle p, \bar{x}(a) \rangle$. It follows that

$$\forall a \in B, \quad z \in P_a(\bar{x}(a)) \Rightarrow \langle p, z \rangle > \langle p, \bar{x}(a) \rangle.$$

The preference relation \succ_a is monotone, i.e. for each $z > 0$, $\bar{x}(a) + z \in P_a(\bar{x}(a))$. It follows that for each $z > 0$, $\langle p, z \rangle > 0$. Now from Assumption 3.1, for each $a \in \Omega$, $e(a) > 0$, hence

$$\inf\{\langle p, z \rangle : z \in E_+\} = 0 < \langle p, e(a) \rangle. \quad \square$$

Appendix A. Finitely many consumers

We suppose in this section that the economy is finite in the sense that the set of consumers $(\Omega, \mathcal{A}, \mu)$ is $(I, 2^I, \sigma)$ where I is a finite set, 2^I is the σ -algebra of all subsets of I and σ is the counting measure.

A.1 The general case

If \mathcal{E} is strong-uniformly proper then we denote by Γ the norm-open convex cone defined by $\Gamma = \{x \in E : \langle \alpha, x^+ \rangle > \langle \beta, x^- \rangle\}$.

Theorem A.1. *Let \mathcal{E} be a finite economy satisfying Assumptions 3.1–3.4. If \mathcal{E} is strong-uniformly proper then there exists a pair (x, p) consisting of a feasible allocation x and a non-zero price p such that*

1. for each $i \in I$, $\langle p, x_i \rangle = \langle p, e_i \rangle$, and $z \in P_i(x_i)$ implies $\langle p, z \rangle \geq \langle p, x_i \rangle$;
2. $\langle p, \omega \rangle = 1$ and $\langle p, \Gamma \rangle > 0$.

Proof. Since order intervals $[0, x] = \{y \in E : 0 \leq y \leq x\}$ are w -compact,¹⁴ following Florenzano [13], there exists a feasible allocation $x = (x_i)_i$ such that¹⁵

$$0 \notin G(x) := \text{co} \bigcup_{i \in I} [P_i(x_i) - e_i].$$

Lemma A.2. $G(x) \cap -\Gamma = \emptyset$.

Proof. To see this,¹⁶ assume by way of contradiction that $G(x) \cap -\Gamma \neq \emptyset$. Then there exist $\gamma \in \Gamma$, $(\lambda_i)_i$ with $\lambda_i \geq 0$, $\sum_i \lambda_i = 1$ and $(z_i)_i$ with $z_i \in P_i(x_i)$ such that

$$\sum_i \lambda_i z_i + \gamma = \sum_i \lambda_i e_i.$$

Suppose first that $\gamma \geq 0$. For each $i \in I$, we set $y_i := z_i + \gamma$. Then $y_i \succ_i z_i$ for each i since preference relations are strictly monotone, whence $y_i \succ_i x_i$ by transitivity. One the other hand,

$$\sum_i \lambda_i y_i = \sum_i \lambda_i e_i$$

and we have thus got a contradiction.

Thus suppose that $\gamma^- \neq 0$. We must have $\gamma^- \leq \sum_i \lambda_i z_i$, so by the Riesz decomposition theorem there exist elements $u_i \geq 0$ such that $u_i \leq z_i$ and $\sum_i \lambda_i u_i = \gamma^-$. Set for each i ,

$$v_i = \frac{\langle \beta, u_i \rangle}{\langle \beta, \gamma^- \rangle} \gamma^+.$$

Since $\langle \alpha, \gamma^+ \rangle > \langle \beta, \gamma^- \rangle$ by definition of Γ ,

$$\langle \alpha, v_i \rangle = \frac{\langle \beta, u_i \rangle}{\langle \beta, \gamma^- \rangle} \langle \alpha, \gamma^+ \rangle \geq \langle \beta, u_i \rangle,$$

¹⁴ Since E is a Banach lattice, the order interval $[0, x]$ is a subset of $\|x\| B$. Since E is separable and reflexive then B is w -compact.

¹⁵ In fact x is an Edgeworth equilibrium of \mathcal{E} .

¹⁶ The argument given in the sequel to establish this lemma is taken from Zame [30] and Podczeck [25].

with strict inequality if $u_i \neq 0$. Hence because $u_i \leq z_i$ and $v_i \geq 0$, we have $z_i - u_i + v_i \succeq_i z_i$ for each i (in fact, $z_i - u_i + v_i \succ_i z_i$ in case $u_i \neq 0$), and therefore by transitivity of preference relations, $z_i - u_i + v_i \succ_i x_i$. Also

$$\sum_i \lambda_i(z_i - u_i + v_i) = \sum_i \lambda_i z_i - \gamma^- + \frac{\langle \beta, \sum_i \lambda_i u_i \rangle}{\langle \beta, \gamma^- \rangle} \gamma^+ = \sum_i e_i,$$

again we get a contradiction. □

Following Lemma A.2, since Γ is norm-open, it follows from the separation theorem that there exists some non-zero linear functional $p \in E^*$ with $\langle p, g \rangle \geq -\langle p, \gamma \rangle$ for each $g \in G(x)$ and $\gamma \in \Gamma$. It is now routine to prove that (x, p) satisfies properties (1) and (2) of Theorem A.1. □

A.2 The separable case

We recall that a utility function $u : \ell^p_+ \rightarrow \mathbb{R}$ is called separable if there exists for each n , a function $v_n : [0, +\infty) \rightarrow \mathbb{R}$ concave and strictly increasing such that

$$\forall x \in \ell^p_+, \quad u(x) = \sum_{n \in \mathbb{N}} v_n(x_n).$$

Theorem A.3. *Let \mathcal{E} be a finite economy satisfying Assumptions 3.1–3.4. If \mathcal{E} is separably proper then there exists a pair (x, p) consisting of a feasible allocation x and a non-zero price p such that*

- (1) *for each $i \in I$, $\langle p, x_i \rangle = \langle p, e_i \rangle$, and $z \in P_i(x_i)$ implies $\langle p, z \rangle \geq \langle p, x_i \rangle$;*
- (2) *$\langle p, \omega \rangle = 1$ and $\|p\|_q \leq (1/\alpha_k \omega_k) \|\beta\|_q$.*

The proof of Theorem A.3 is mostly inspired by the proof of Theorem 3 in Araujo–Monteiro [4].

Proof. We prove Theorem A.3 in two steps. For the first step, we suppose that the economy satisfies an additional assumption on the initial endowments.

Step 1: Strictly positive initial endowments. Suppose that for each i , e_i is strictly positive. Let E^ω be the vector space of all $z \in \ell^p$ such that there exists $r > 0$ satisfying $-r\omega \leq z \leq r\omega$. From Lemma 1 in Araujo–Monteiro [4], there exists a pair (x, p) consisting of a feasible allocation¹⁷ x and a non-zero linear functional $p : E^\omega \rightarrow \mathbb{R}$ such that p is positive, i.e. $\langle p, z \rangle \geq 0$ for each $z \in E^\omega_+$; $\langle p, \omega \rangle = 1$ and such that

$$\forall i \in I, \quad \langle p, x_i \rangle = \langle p, e_i \rangle \quad \text{and} \quad z \in P_i(x_i) \cap E^\omega \Rightarrow \langle p, z \rangle \geq \langle p, e_i \rangle.$$

Now there exists i with $\langle p, e_i \rangle > 0$, and since u_i is strictly monotone, p is strictly positive, i.e. $\langle p, z \rangle > 0$ for each $0 \neq z \in E^\omega_+$. In particular $\langle p, e_i \rangle > 0$ for each

¹⁷ Note that if x is a feasible allocation then $x_i \in E^\omega$.

$i \in I$. By the *concave alternative* (see Lemma 5 in [4]), for each i there exists $\lambda_i > 0$ such that

$$(A.1) \quad \forall z \in E_+^\omega, \quad u_i(z) - u_i(x_i) \leq \lambda_i \langle p, z - x_i \rangle.$$

For $z \in \ell_+^p$ we define $S(z) = \{h \in \ell^p : \exists t > 0, z + th \geq 0\}$ and $I(z) = S(z) \cap -S(z)$. Using (A.1) like in [4], we have

$$(A.2) \quad \forall h \in I(x_i) \cap E_+^\omega, \quad \lambda_i \langle p, h \rangle \leq \sum_n v_{i,n}^-(x_{i,n}) h_n.$$

Since e_i is strictly positive for each i , we have that $b := \inf\{e_i : i \in I\}$ is strictly positive. Hence E^b is norm-dense in ℓ^p . From this we conclude that if p is norm-continuous on E^b then we can extend it to a linear functional still noted p in ℓ^q , such that (x, p) satisfies

$$\forall i \in I, \quad \langle p, x_i \rangle = \langle p, e_i \rangle \quad \text{and} \quad z \in P_i(x_i) \Rightarrow \langle p, z \rangle \geq \langle p, e_i \rangle.$$

So let us prove that p is norm-continuous on E^b .

We define for each n , $I_n := \{i \in I : x_{i,n} \geq e_{i,n}\}$ and for each i we define $N_i = \{n \in \mathbb{N} : i = \min I_n\}$. Since x is a feasible allocation, we have $I_n \neq \emptyset$ for every n and $(N_i)_i$ is a partition of \mathbb{N} . Take $h \in E^b$, there exists $r > 0$ such that $-re_i \leq h \leq re_i$ for each i . Now let $h^i \in E^b$ be defined by

$$h^i = (h_n^i)_n \quad \text{where} \quad h_n^i = \begin{cases} |h_n| & \text{if } n \in N_i \\ 0 & \text{if } n \notin N_i. \end{cases}$$

As h^i belongs to $I(x_i)$, it follows from (A.2) that¹⁸

$$\lambda_i \langle p, h^i \rangle \leq \sum_{n \in N_i} v_{i,n}^-(x_{i,n}) |h_n| \leq \sum_{n \in N_i} v_{i,n}^-(e_{i,n}) |h_n|.$$

Since p is positive, we have that $|\langle p, h \rangle| \leq \sum_i \langle p, h^i \rangle$. It follows that

$$|\langle p, h \rangle| \leq \sum_i (1/\lambda_i) \sum_{n \in N_i} v_{i,n}^-(e_{i,n}) |h_n|.$$

We define $u'_i(z) \cdot h = \lim_{r \rightarrow 0} (1/r)(u_i(z + rh) - u_i(z))$ for each $z \in \ell_+^p$ and each $h \in S(z)$. It follows from (A.1) and separable properness that

$$0 < \alpha_k \omega_k < u'_i(x_i) \cdot \omega \leq \lambda_i.$$

In particular

$$\begin{aligned} |\langle p, h \rangle| &\leq (1/\alpha_k \omega_k) \sum_i \sum_{n \in N_i} v_{i,n}^-(e_{i,n}) |h_n| \\ &\leq (1/\alpha_k \omega_k) \sum_{n \in \mathbb{N}} \beta_n |h_n| \leq (1/\alpha_k \omega_k) \|\beta\|_q \|h\|_p. \end{aligned}$$

¹⁸ Note that $v_{i,n}^-$ is a decreasing function.

From separable properness we have that $\|\beta\|_q < +\infty$. This proves the norm-continuity of p on E^b .

Step 2: Positive initial endowments. Let \mathcal{E} be a separably proper finite economy satisfying Assumptions 3.1–3.4. Let v be a strictly positive vector of ℓ^p and consider \mathcal{E}^n the economy defined by $\mathcal{E} = (I, \ell^p, X, \succ, e^n)$ where $e_i^n := e_i + (1/n)v$. Since $v_i^-(e_i^n) \leq v_i^-(e_i)$, the economy \mathcal{E}^n is separably proper and satisfies Assumptions 3.1–3.4. Applying Step 1, there exists a pair (x^n, p^n) consisting of a feasible allocation x^n and a non-zero price p^n such that for each $i \in I$, $\langle p, x_i^n \rangle = \langle p, e_i^n \rangle$, $z \in P_i(x_i^n)$ implies $\langle p^n, z \rangle \geq \langle p^n, x_i^n \rangle$, $\langle p^n, \omega^n \rangle = 1$ and $\|p^n\|_q \leq (1/\alpha_k \omega_k^n) \|\beta\|_q$.

Since the sequence $(\omega_k^n)_n$ is norm-convergent to ω_k , it follows that the sequence $(p^n)_n$ is norm-bounded, and passing to a subsequence if necessary, we can suppose that the sequence $(p^n)_n$ is w^* -convergent to a price $p \in \ell^q$ with $\|p\|_q \leq (1/\alpha_k \omega_k) \|\beta\|_q$. Moreover, since $\langle p^n, \omega^n \rangle = 1$ it follows that $\langle p, \omega \rangle = 1$.

For each i , x_i^n belongs to the interval $[0, \omega + v]$, in particular, passing to a subsequence if necessary, we can suppose that $(x_i^n)_n$ is w -convergent to $x_i \in \ell^p_+$. Moreover, since $\sum_i x_i^n = \omega^n$, we have that x is a feasible allocation for the economy \mathcal{E} . It is now routine to prove that for each $i \in I$, $\langle p, x_i \rangle = \langle p, e_i \rangle$, and $z \in P_i(x_i)$ implies $\langle p, z \rangle \geq \langle p, x_i \rangle$. □

Appendix B. Measurable correspondences

We consider $(\Omega, \mathcal{A}, \mu)$ a finite complete measure space and (D, d) a separable metric space. We recall that a function $f : \Omega \rightarrow D$ is *measurable* if for each open set $G \subset D$, $f^{-1}(G) \in \mathcal{A}$ where $f^{-1}(G) := \{a \in \Omega : f(a) \in G\}$. A correspondence $F : \Omega \rightarrow D$ is *graph measurable* if $G_F := \{(a, x) \in \Omega \times D : x \in F(a)\} \in \mathcal{A} \otimes \mathcal{B}(D)$, where $\mathcal{B}(D)$ is the σ -algebra of Borelian subsets of D .

B.1 Measurable selections

Following Aumann [8], graph measurable correspondences have measurable selections.

Theorem B.1. *Consider F a graph measurable correspondence from Ω into D with non-empty values. If (D, d) is complete then there exists a sequence $(z_n)_n$ of measurable selections of F , such that for each $a \in \Omega$, $(z_n(a))_n$ is d -dense in $F(a)$.*

B.2 Discretization of measurable correspondences

Definition B.2. A partition $\sigma = (A_i)_{i \in I}$ of Ω is a *measurable partition* if for each $i \in I$, the set A_i is non-empty and belongs to \mathcal{A} . A finite subset A^σ of Ω is *subordinated to the partition σ* if there exists a family $(a_i)_{i \in I} \in \prod_{i \in I} A_i$ such that $A^\sigma = \{a_i : i \in I\}$.

Given a couple (σ, A^σ) where $\sigma = (A_i)_{i \in I}$ is a measurable partition of Ω , and $A^\sigma = \{a_i : i \in I\}$ is a finite set subordinated to σ , we consider $\phi(\sigma, A^\sigma)$ the application which maps each measurable function f to a simple measurable function $\phi(\sigma, A^\sigma)(f)$, defined by

$$\phi(\sigma, A^\sigma)(f) := \sum_{i \in I} f(a_i)\chi_{A_i},$$

where χ_{A_i} is the characteristic¹⁹ function associated to A_i .

Definition B.3. A function $s : \Omega \rightarrow D$ is called a *simple function subordinated* to f if there exists a couple (σ, A^σ) where σ is a measurable partition of Ω , and A^σ is a finite set subordinated to σ , such that $s = \phi(\sigma, A^\sigma)(f)$.

Given a couple (σ, A^σ) where $\sigma = (A_i)_{i \in I}$ is a measurable partition of Ω , and $A^\sigma = \{a_i : i \in I\}$ is a finite set subordinated to σ , we consider $\psi(\sigma, A^\sigma)$, the application which maps each measurable correspondence $F : \Omega \rightarrow D$ to a simple measurable correspondence $\psi(\sigma, A^\sigma)(F)$, defined by

$$\psi(\sigma, A^\sigma)(F) := \sum_{i \in I} F(a_i)\chi_{A_i}.$$

Definition B.4. A correspondence $S : \Omega \rightarrow D$ is called a *simple correspondence subordinated* to a correspondence F if there exists a couple (σ, A^σ) where σ is a measurable partition of Ω , and A^σ is a finite set subordinated to σ , such that $S = \psi(\sigma, A^\sigma)(F)$.

Remark B.5. If f is a function from Ω to D , let $\{f\}$ be the correspondence from Ω into D , defined for each $a \in \Omega$ by $\{f\}(a) := \{f(a)\}$. We check that

$$\psi(\sigma, A^\sigma)(F) = \{\phi(\sigma, A^\sigma)(f)\}.$$

The space of all non-empty subsets of D is noted $\mathcal{P}^*(D)$. We let τ_{W_d} be the Wijsman topology on $\mathcal{P}^*(D)$, that is the weak topology on $\mathcal{P}^*(D)$ generated by the family of distance functions $(d(x, \cdot))_{x \in D}$.

Hereafter we assert that for a countable set of graph measurable correspondences, there exists a sequence of measurable partitions *approximating* each correspondence. The proof of the following theorem is given in Martins-da-Rocha [18].

Theorem B.6. *Let \mathcal{F} be a countable set of graph measurable correspondences with non-empty values from Ω into D and let \mathcal{G} be a finite set of integrable functions from Ω into \mathbb{R} . There exists a sequence $(\sigma^n)_n$ of finer and finer measurable partitions $\sigma^n = (A_i^n)_{i \in I^n}$ of Ω , satisfying the following properties.*

- (a) *Let $(A^n)_n$ be a sequence of finite sets A^n subordinated to the measurable partition σ^n and let $F \in \mathcal{F}$. For each $n \in \mathbb{N}$, we define the simple correspondence $F^n := \psi(\sigma^n, A^n)(F)$ subordinated to F . Then for each $a \in \Omega$, $F(a)$ is the Wijsman limit of the sequence $(F^n(a))_n$, i.e.,*

$$\forall a \in \Omega, \quad \forall x \in D, \quad \lim_n d(x, F^n(a)) = d(x, F(a)).$$

¹⁹ That is, for each $a \in \Omega$, $\chi_{A_i}(a) = 1$ if $a \in A_i$ and $\chi_{A_i}(a) = 0$ elsewhere.

(b) *There exists a sequence $(A^n)_n$ of finite sets A^n subordinated to the measurable partition σ^n , such that for each n , if we let $f^n := \phi(\sigma^n, A^n)(f)$ be the simple function subordinated to each $f \in \mathcal{G}$, then*

$$\forall f \in \mathcal{G}, \quad \forall a \in \Omega, \quad |f^n(a)| \leq 1 + \sum_{g \in \mathcal{G}} |g(a)|.$$

In particular, for each $f \in \mathcal{G}$,

$$\lim_{n \rightarrow \infty} \int_{\Omega} |f^n(a) - f(a)| d\mu(a) = 0.$$

Remark B.7. The property (a) implies in particular that, if $(x^n)_n$ is a sequence of D , d -converging to $x \in D$, then

$$\forall a \in \Omega, \quad \lim_n d(x^n, F^n(a)) = d(x, F(a)).$$

It follows that if F is non-empty closed valued, then property (a) implies that

$$\forall a \in \Omega, \quad \text{ls}_n F^n(a) \subset F(a).$$

Appendix C. Fatou’s Lemma

The proof of the following theorem is given in Cornet–Martins-da-Rocha [10].

Theorem C.1. *Let $(\Omega, \mathcal{A}, \mu)$ be a finite positive complete measure space. Let $(f_n)_n$ be a sequence of Bochner integrable mappings from Ω to E_+ , which is mean norm-bounded, i.e.*

$$\sup_n \int_{\Omega} \|f_n(a)\| d\mu(a) < +\infty.$$

Suppose that $w\text{-}\lim_n \int_{\Omega} f_n d\mu$ exists in E then there exists a Bochner integrable mapping f such that

$$\int_{\Omega} f d\mu \leq w\text{-}\lim_n \int_{\Omega} f_n d\mu$$

and

$$f(a) \in \overline{\text{co}} w\text{-}\text{ls}_n \{f_n(a)\} \quad \text{a.e.}$$

Moreover

$$\int_{\Omega} \|f(a)\| d\mu(a) \leq \sup_n \int_{\Omega} \|f_n(a)\| d\mu(a).$$

Appendix D. Fatou’s cone

Proposition D.1. *Let (T, \mathcal{T}, σ) be an atomless measure space. For $1 < p < +\infty$ we consider the space $E = L^p(T, \mathcal{T}, \sigma)$ ordered by the cone E_+ defined by*

$$E_+ = \{f \in L^p(T, \mathcal{T}, \sigma) : \forall t \in T, f(t) \geq 0\}.$$

The space E ordered by the cone E_+ is a Banach lattice. But for every Bochner integrable mapping $e : \Omega \rightarrow E_+$ with $\int_{\Omega} e d\mu > 0$, the cone E_+ is not a Fatou’s cone relatively to e .

Proof. Consider a Bochner integrable mapping $e : \Omega \rightarrow E_+$ with $\int_{\Omega} e d\mu > 0$. Suppose that E_+ is a Fatou’s cone relatively to e , then there exists a positive extended linear functional ρ satisfying conditions (a) and (b) of Definition 2.2. Since $a \mapsto \rho[e(a)]$ is integrable, there exists Ω' in \mathcal{A} such that $\mu(\Omega \setminus \Omega') = 1$ and for all $a \in \Omega'$, $\rho[e(a)] < +\infty$. Moreover, since $\int_{\Omega} e d\mu > 0$, there exists $b \in \Omega'$ such that $e(b) > 0$. We let $\omega = e(b)$, without any loss of generality, we may assume that $\|\omega\| = 1$. We denote by δ the probability on (T, \mathcal{T}) defined by

$$\forall B \in \mathcal{T}, \quad \delta(B) = \int_B [\omega(t)]^p \sigma(dt).$$

Since (T, \mathcal{T}, σ) is atomless, the measure space (T, \mathcal{T}, δ) is also atomless. Applying Lyapunov Convexity Theorem (see Aliprantis–Border [2, Theorem 12.33]), $\{\delta(B) : B \in \mathcal{T}\} = [0, 1]$. Fix $n > 0$, then there exists $T_1 \in \mathcal{T}$ such that

$$\int_{T_1} \omega^p d\sigma = \frac{1}{n}.$$

Now the restriction of δ to $T \setminus T_1$ is a finite positive measure, such that

$$\{\delta(B) : B \in \mathcal{T} \text{ and } B \cap T_1 = \emptyset\} = [0, 1 - 1/n].$$

Hence there exists $T_2 \in \mathcal{T}$ with $T_2 \cap T_1 = \emptyset$ such that

$$\int_{T_2} \omega^p d\sigma = \frac{1}{n}.$$

By induction, there exists a measurable partition (T_1, \dots, T_n) of T such that

$$\forall k \in \{1, \dots, n\}, \quad \int_{T_k} \omega^p d\sigma = \frac{1}{n}.$$

For each $k \in \{1, \dots, n\}$, let χ_k be the characteristic function of T_k , i.e. $\chi_k(t) = 1$ if $t \in T_k$ and $\chi_k(t) = 0$ elsewhere. Then

$$n^{1-1/p} = \sum_{k=1}^n \left(\frac{1}{n}\right)^{1/p} = \sum_{k=1}^n \|\omega\chi_k\| \leq \sum_{k=1}^n \rho[\omega\chi_k] = \rho(\omega),$$

contradiction. □

References

1. Aliprantis, C.D.: Separable utility functions. *Journal of Mathematical Economics* **28**(4), 415–444 (1997)
2. Aliprantis, C.D., Border, K.C.: *Infinite dimensional analysis*. Berlin Heidelberg New York: Springer 1999
3. Araujo, A., Monteiro, P.K.: *Equilibrium and dynamics: Essays in honor of David Gale*, ch. General equilibrium with infinitely many goods: the case of separable utilities. London: Macmillan 1989
4. Araujo, A., Monteiro, P.K.: Equilibrium without uniform conditions. *Journal of Economic Theory* **48**, 416–427 (1989)
5. Arrow, K.J., Debreu, G.: Existence of an equilibrium for a competitive economy. *Econometrica* **22**, 265–290 (1954)
6. Aumann, R.J.: Markets with a continuum of agents. *Econometrica* **32**, 39–50 (1964)
7. Aumann, R.J.: Existence of a competitive equilibrium in markets with a continuum of traders. *Econometrica* **34**, 1–17 (1966)
8. Aumann, R.J.: Measurable utility and measurable choice theorem. *La Décision Centre National de la Recherche Scientifique*, Paris, pp. 15–26 (1969)
9. Bewley, T.F.: *Equilibrium theory in infinite dimensional spaces*, chapt. A very weak theorem on the existence of equilibria in atomless economies with infinitely many commodities. New York: Springer 1991
10. Cornet, B., Martins-da-Rocha, V.F.: Fatou’s lemma for unbounded Gelfand integrable mappings. Working Paper, Université Paris-I Panthéon Sorbonne (2002)
11. Debreu, G.: *Theory of value: An axiomatic analysis of economic equilibrium*. New York: Wiley 1959
12. Diestel, J., Uhl, J.: *Vector measure*. Mathematical Surveys, No. 15, American Mathematical Society, Providence, RI (1977)
13. Florenzano, M.: On the non-emptiness of the core of a coalitional production economy without ordered preferences. *Journal of Mathematical Analysis and Application* **141**, 484–490 (1989)
14. Hildenbrand, W.: Existence of equilibria for economies with production and a measure space of consumers. *Econometrica* **38**, 608–623 (1970)
15. Jones, L.E.: Existence of equilibrium with infinitely many consumers and infinitely many commodities. *Journal of Mathematical Economics* **12**, 119–138 (1983)
16. Khan, M.A., Yannelis, N.C.: Equilibrium in markets with a continuum of agents and commodities. In: Khan, M.A., Yannelis, N.C. (eds.) *Equilibrium theory in infinite dimensional spaces*. Berlin Heidelberg New York: Springer 1991
17. Le Van, C.: Complete characterization of Yannelis–Zame and Chichilnisky–Kalman–Mas-Colell properness for separable concave functions defined in L^p_+ and L^p . *Economic Theory* **8**(1), 155–166 (1996)
18. Martins-da-Rocha, V.F.: Existence of equilibria for economies with a measurable space of agents and non-ordered preferences. Working paper, Université Paris-I Panthéon Sorbonne (2001)
19. Martins-da-Rocha, V.F.: Equilibria in large economies with a separable Banach commodity space and non-ordered preferences. *Journal of Mathematical Economics* **39**(8), 863–889 (2003)
20. Martins-da-Rocha, V.F.: Equilibria in large economies with differentiated commodities and non-ordered preferences. *Economic Theory* **23**(3), 529–552 (2004)
21. Mas-Colell, A.: A model of equilibrium with differentiated commodities. *Journal of Mathematical Economics* **2**, 263–295 (1975)
22. McKenzie, L.W.: On equilibrium in Graham’s model of world trade and other competitive systems. *Econometrica* **22**, 147–161 (1954)
23. Ostroy, J., Zame, W.R.: Nonatomic economies and the boundaries of perfect competition. *Econometrica* **62**, 593–633 (1994)
24. Podczeck, K.: Markets with infinitely many commodities and a continuum of agents with non-convex preferences. *Economic Theory* **9**, 585–629 (1997)
25. Podczeck, K.: Core and Walrasian equilibria when agents’ characteristics are extremely dispersed. *Economic Theory* **22**(4), 699–725 (2003)
26. Podczeck, K.: Note on the Core-Walras equivalence problem when the commodity space is a Banach lattice. Working Paper # 307, University of Vienna (2003)

27. Rustichini, A., Yannelis, N.C.: Edgeworth's conjecture in economies with a continuum of agents and commodities. *Journal of Mathematical Economics* **20**, 307–326 (1991)
28. Tourky, R., Yannelis, N.C.: Markets with many more agents than commodities: Aumann's "hidden" assumption. *Journal of Economic Theory* **101**, 189–221 (2001)
29. Yannelis, N.C., Zame, W.R.: Equilibria in Banach lattices without ordered preferences. *Journal of Mathematical Economics* **15**, 75–110 (1986)
30. Zame, W.R.: Markets with a continuum of traders and infinitely many commodities. Working paper, SUNY at Buffalo (1986)