
THE ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF THE FORCED BURGERS EQUATION ON THE CIRCLE

BY PATRICK BERNARD

ABSTRACT. — We describe the asymptotic behavior of entropy solutions of inviscid Burgers equation on the circle with time-periodic forcing. These solutions converge to periodic states, but the period of these limit states may be greater than the period of the forcing. A Corollary is that the only solutions without shocks are periodic. This result can be interpreted in terms of the invariant curves of the associated dynamics.

RÉSUMÉ. — On décrit les comportements asymptotiques des solutions entropiques d'équations de Burgers sans viscosité avec un potentiel dépendant périodiquement du temps. Ces solutions convergent vers des états périodiques, dont la période est un multiple de celle du potentiel. Un corollaire est que les seules solutions sans choc, c'est à dire les seules solutions continues, sont périodiques. Ce résultat a une interprétation en terme des courbes invariantes de la dynamique associée.

We study the quasi-linear first order partial differential equation

$$(B) \quad \partial_t y + \partial_x (H(t, x, y)) = 0,$$

where t is a real time, x is a periodic real variable, and the unknown y is a real valued function of t and x . The hypotheses on H will be specified later. Equations of this kind have been extensively studied as a very simple example of turbulence. Here is a physical model, which helps the intuition. The Hamiltonian H gives rise to a classical dynamics for particles on the configuration space, which is here the circle of x variable. On each point x of this circle, put a particle with momentum $y(0, x)$. Assume that each of these particles moves, independently of each other, driven by the classical dynamics associated to H . The solution $y(t, x)$ is then the momentum of the particle which is at point x at time t . The function $y(t, x)$ may be seen as the Eulerian description of this very simple fluid. Of course it might happen that after a certain time several different particles with different momenta have the same position. This is the reason why the equation do not have classical solutions in general. However, generalized solution can be defined by the following additional requirement: when a collision occurs between particles, these particles glue together and form what will be called a shock. Shocks then absorb the particles which hit them. Note that these shocks are not the same kind of objects as particles, their evolution is not driven by the Hamiltonian. The solutions describing this evolution are called the entropy solutions of (B).

The main goal of the paper is to describe the asymptotic behavior of entropy solutions of (B). We prove that, if the Hamiltonian is time-periodic, each solution converges to a periodic solution. The period of the limit is a multiple of the period of the Hamiltonian. We also prove that all solutions undergo shocks, except perhaps time-periodic solutions. This does not mean, however, that periodic solutions are always free of shocks. For many Hamiltonians, there do not exist any solution without shocks.

In order to obtain this asymptotic behavior, we introduce another equation, satisfied by the primitive of entropy solutions, the Hamilton-Jacobi equation, whose viscosity solutions are by

First version summer 2003, version of septembre 2004

PATRICK BERNARD, Institut Fourier BP 74, 38402, Saint Martin d'Hères Cedex, France

E-mail : Patrick.Bernard@ujf-grenoble.fr • *Url* : <http://www-fourier.ujf-grenoble.fr/~pbernard/>

now well understood. On the way, we have to understand how knowledge on entropy solutions of Burgers equation can be gained from knowledge on viscosity solutions of the associated Hamilton-Jacobi equation. Our second aim is to describe with some details these relations, and to show how most of what is known on the Burgers equation can be recovered from this point of view. As a consequence, a large part of the paper is devoted to the presentation of known results both on the Burgers equation and on the Hamilton-Jacobi equation. It is worth noticing that the trajectories of the particles, that is the Lagrangian description of our elementary fluid, play a major role in this approach.

Recently, it has been understood that the study of entropy solutions of equation (B) provides interesting insights in the dynamics of the Hamiltonian H . This aspect is one of our motivations, and our work also imply a new observation concerning the dynamics of the time-one flow of H . We prove that if a curve in phase space is a graph, and is such that infinitely many of its successive images by the time-one flow are also graphs, then the curve has to be periodic, that is mapped into itself by some power of the flow. Similar results had previously been obtained by Albert Fathi in the case of autonomous Hamiltonian systems.

Most of this work was written while I was staying for one year in EPFL, Swizerland, with a grant of the Swiss National Fund for research. This period was very pleasant and fruitful. I would like to thank Boris Buffoni for proposing me this opportunity, and for his interest to the present work.

1. INTRODUCTION

1.1. The standard circle \mathbb{R}/\mathbb{Z} is denoted by \mathbb{T} . The cotangent bundle $T^*\mathbb{T}$ is identified with $\mathbb{T} \times \mathbb{R}$. Given a function $f(t, x)$ of two variables, we will denote by f_t the function $x \mapsto f(t, x)$. The partial derivative with respect to the variable t will be denoted by $\partial_t f$. Throughout this paper, we will consider a time-periodic Hamiltonian $H(t, x, p) : \mathbb{R} \times T^*\mathbb{T} = \mathbb{R} \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$, the associated time-periodic vector-field of $\mathbb{T} \times \mathbb{R}$ is noted X . We have

$$X(t, x, p) = (\partial_p H(t, x, p), -\partial_x H(t, x, p)).$$

1.2. The following standard hypotheses will be assumed :

- i.* The Hamiltonian H is C^2 and 1-periodic in t .
- ii.* The Hamiltonian H is convex in p , and $\partial_{pp}^2 H > 0$.
- iii.* The Hamiltonian has superlinear growth in p , *i.e.* $\lim_{|p| \rightarrow \infty} H(t, x, p)/|p| = \infty$ for each (t, x) .
- iv.* The Hamiltonian flow is complete. More precisely for all (t_0, x_0, p_0) , there exists a C^1 curve $\gamma(t) = (x(t), p(t)) : \mathbb{R} \rightarrow \mathbb{T} \times \mathbb{R}$ such that $(x(t_0), p(t_0)) = (x_0, p_0)$ and $\dot{\gamma}(t) = X(t, \gamma(t))$ for all $t \in \mathbb{R}$.

The mapping $\gamma(t_0) \mapsto \gamma(t)$ is a diffeomorphism of $\mathbb{T} \times \mathbb{R}$, denoted $\phi_{t_0, t}$. We will pay special attention to the diffeomorphism $\phi = \phi_{0, 1}$. Note that the completeness hypothesis is satisfied if there exists a constant C such that $|\partial_t H| \leq C(1 + H)$.

1.3. A typical example of Hamiltonian satisfying our hypotheses is

$$H(t, x, p) = \frac{1}{2}p^2 + V(t, x)$$

with a C^2 potential V periodic in t .

1.4. We consider the equation

$$(B) \quad \partial_t y + \partial_x (H(t, x, y)) = 0$$

of the unknown function $y(t, x) : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$. This equation will be called the Burgers equation in the sequel. Note that in case $H = p^2/2 + V(t, x)$, we have the standard forced inviscid Burgers

equation

$$\partial_t y + y \partial_x y = -\partial_x V(t, x).$$

1.5. The Burgers equation is quasi-linear, and its characteristics (see [A], chapter 2) are the trajectories of X . This means that if $y(t, x) : [a, b] \times \mathbb{T} \rightarrow \mathbb{R}$ is a C^1 solution of the Burgers equation, and if $\gamma(t) = (x(t), p(t)) : [a, b] \rightarrow \mathbb{T} \times \mathbb{R}$ is a trajectory of the Hamiltonian vectorfield X , such that $p(a) = y(a, x(a))$, then the equality $p(t) = y(t, x(t))$ holds for all $t \in [a, b]$. In other words, for each $a \leq t_0 \leq t_1 \leq b$, the graph of the function $x \mapsto y(t_1, x)$ is the image by the diffeomorphism ϕ_{t_0, t_1} of the graph of the function $x \mapsto y(t_0, x)$.

1.6. Still assuming that y is a C^1 solution of the Burgers equation, we obtain that

$$c(t) = \int_{\mathbb{T}} y(t, x) dx$$

is a constant, that we denote c . The function y can be written $y(t, x) = c + \partial_x u(t, x)$, where $u(t, x) : I \times \mathbb{T} \rightarrow \mathbb{R}$ satisfies the Hamilton-Jacobi equation

$$(HJc) \quad \partial_t u + H(t, x, c + \partial_x u) = 0.$$

1.7. It is known that there exists in general no classical solution of the Burgers equation defined on $\mathbb{R}^+ \times \mathbb{T}$ satisfying a given initial condition $y(0, x) = y_0(x)$, see [A] or [Se]. Note that a consequence of our results is that the only classical solutions defined for all positive times are periodic in time. However, the Cauchy problem is well-posed in the sense of entropy solutions (we will define entropy solutions in the sequel). More precisely, to all $t_0 \in \mathbb{R}$ and all function $y_0 \in L^1(\mathbb{T})$, we associate a unique entropy solution $y(t, x) \in C([t_0, \infty), L^1(\mathbb{T}))$ such that $y_t \rightarrow y_0$ as $t \rightarrow t_0$ in a weak sense, see 3.2. This result is classical, see [Se] for example.

1.8. We want to describe the asymptotic behavior of entropy solutions. Let us first recall that for each c there exists a 1-periodic solution of average c . More precisely, there exists a continuous and increasing function $c \mapsto y^c$ from \mathbb{R} to $C(\mathbb{R}, L^1(\mathbb{T}))$ which, to each c , associates a 1-periodic Entropy solution of average c , see [KO] or [JKM]. Note however that there may exist more than one 1-periodic solution of a given average c . It is natural to ask whether all solutions are attracted by these 1-periodic solutions. The answer is negative, there are examples where there exist periodic Entropy solutions of minimal period greater than 1, see [FM]. These subharmonic solutions in turn attract all other solutions, as we now state.

1.9. THEOREM *Let $y(t, x) :]t_0, \infty[\times \mathbb{T} \rightarrow \mathbb{R}$ be an entropy solution of the Burgers equation. There exists an integer T and an entropy solution $\omega(t, x) : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ which is T -periodic in t and such that*

$$\|y_t - \omega_t\| \rightarrow 0$$

in $L^1(\mathbb{T})$ as t goes to infinity.

If H is a function of p only, then $\omega(t, x)$ is the constant $\int y(t, x) dx$. The result in this special case has been obtained by Lax [L]. If H does not depend on t , then the asymptotic solution ω does not depend on t either, the result in this case follows from works of Roquejoffre [Ro] and Fathi [Fa3]. The theorem will be proved in section 3 as a consequence of a similar result for viscosity solutions of (HJc) obtained in [Be], see also [BR]. We will also prove the following Corollary, which was obtained by Fathi [Fa3] in the Autonomous case.

COROLLARY *If $y(t, x) :]t_0, \infty[\times \mathbb{T} \rightarrow \mathbb{R}$ is a continuous solution of Burgers equation, then the function y is the restriction of a Lipschitz solution $\omega : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ which satisfies $\omega(t + T, x) = \omega(t, x)$ for some $T \in \mathbb{N}$.*

1.10. One can compare the situation with the viscous case. If one considers the parabolic equation

$$(B_\mu) \quad \partial_t y + \partial_x (H(t, x, y)) = \mu \partial_{xx} y$$

with $\mu > 0$, the behavior is much simpler. One can prove along the lines of [JKM], see also [B], that for each c , there exists a unique solution y^c of average c which is 1-periodic in time. This solution attracts all the solutions of average c . More precisely, if $y : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ is a solution of (B_μ) , and if $\int y_t dx = c$, then $y_t - y_t^c \rightarrow 0$ uniformly as $t \rightarrow \infty$.

1.11. The result in the inviscid case can be used to study the dynamics of the diffeomorphism $\phi = \phi_{0,1}$. Note that this diffeomorphism is a finite composition of area preserving right twist maps, and that any finite composition of right twist maps can be obtained this way. This correspondence between the twist property and the convexity of the Hamiltonian has been described by Moser, see [Mo].

1.12. In order to give a more geometrical meaning to Theorem 1.9, we consider the set \mathcal{E} of functions $f : \mathbb{T} \rightarrow \mathbb{R}$ which can be locally written as the sum of a continuous and a decreasing function. A function $f \in \mathcal{E}$ has a right limit $f^-(x)$ and a left limit $f^+(x)$ at each point x . These limits satisfy $f^-(x) \leq f^+(x)$, with a strict inequality on an at most countable set. Let $G^-(f)$ and $G^+(f)$ be the graphs, in $\mathbb{T} \times \mathbb{R}$ of the functions f^- and f^+ . We define the graph $G(f) = G^-(f) \cap G^+(f)$ of f . Note that $\overline{G(f)} = G^-(f) \cup G^+(f)$. It is also useful to consider the set

$$H(f) = \bigcup_x \{x\} \times [f^-(x), f^+(x)] \subset \mathbb{T} \times \mathbb{R},$$

which is a Jordan curve containing $\overline{G(f)}$. The Hausdorff distance $d_H(f, g)$ between the compact sets $H(f)$ and $H(g)$ defines a distance d_H on \mathcal{E} (one should take the quotient of \mathcal{E} by the relation of almost everywhere equality). The convergence in L^1 implies convergence for the distance d_H , see 3.3.

1.13. The link between entropy solutions and the dynamics of ϕ can now be detailed, see 3.6. It is convenient to denote by $E : L^1 \rightarrow L^1$ the operator which associates to each initial condition y_0 the solution y_1 at time 1. The operator E is continuous. In fact, as is well known, it is a contraction. We will prove this fact in section 4. For each $y \in L^1$, we have $E(y) \in \mathcal{E}$. If in addition $y \in \mathcal{E}$, then

$$\overline{G(E(y))} \subset \phi(G(y)).$$

This property has striking consequences. For example, if y^c is a fixed point of E , then $y^c \in \mathcal{E}$ and $\overline{G(y^c)}$ is negatively invariant. As a consequence, the α -limit of $\phi|_{G(y^c)}$ is a non-empty compact set which is fully invariant by ϕ . It is an Aubry-Mather set. The rotation number $\rho(c) \in \mathbb{R}$ of the orbits of this set depends only on c , and the function $c \mapsto \rho(c)$ is non-decreasing and continuous, see 2.7.

1.14. Having defined the rotation number $\rho(c)$ allows us to complement Theorem 1.9. The asymptotic behavior of solutions depend strongly on their space average c and on the associated rotation number $\rho(c)$. If $\rho(c)$ is irrational, then there exists a single fixed point of E of average c , as was proved in [E] and [So]. We will prove that it attracts all the trajectories of average c , that is one can take $T = 1$ in Theorem 1.9. If $\rho(c)$ is rational, $\rho(c) = p/q$ in lowest terms, then one can take $T = q$ in Theorem 1.9. It is thus natural to define the integer $T(c)$ by $T(c) = 1$ if $\rho(c)$ is irrational and $T(c) = q$ if $\rho(c) = p/q$ in lowest terms, and we have the following refinement of Theorem 1.9 :

THEOREM *For each entropy solution $y :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of (B) , there exists an entropy solution $\omega(t, x) : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ which is $T(c)$ -periodic in time, with $c = \int_{\mathbb{T}} y_t(x) dx$, and such that $y_t - \omega_t \rightarrow 0$ in $L^1(\mathbb{T})$ as t goes to infinity. As a consequence, we also have $d_H(y_t, \omega_t) \rightarrow 0$.*

1.15. Corollary 1.9 can be restated as a new result on the dynamics of ϕ , which may be seen as a converse to the celebrated result of Birkhoff (see [Ma2], [HF], [Si]) stating that a rotational (not homotopic to a constant) Jordan curve in $\mathbb{T} \times \mathbb{R}$ which is invariant under ϕ is the graph of a Lipschitz function $y : \mathbb{T} \rightarrow \mathbb{R}$. This Corollary was obtained by Fathi [Fa3], [Fa2] in the autonomous case.

COROLLARY *Let $G \subset \mathbb{T} \times \mathbb{R}$ be the graph of a continuous function $y : \mathbb{T} \rightarrow \mathbb{R}$. Assume that there exists an increasing sequence n_k of positive integers such that $\phi^{n_k}(G)$ is the graph of a continuous function. Then there exists a positive integer T such that $\phi^T(G) = G$, and the function y is Lipschitz. In addition, if $\phi(G) \neq G$, then the rotation number of $\phi|_G^T$ is an integer, hence G contains a T -periodic point of ϕ .*

Let us mention that it is certainly possible to give a more direct proof of this Corollary. One could use a topological approach, as suggested to me by P. Le Calvez or a variational approach, as suggested by J. Xia. The proof presented here in 3.9 is however extremely short.

1.16. In the rest of the paper, we will detail the outline given above. We will obtain all the important properties of entropy solutions of (B) as consequences of properties of the viscosity solutions of (HJc). Hence we first describe these viscosity solutions in section 2, and draw our conclusions in section 3. We explain in section 4 how to understand in our framework the important fact that entropy solutions form a contraction in L^1 .

2. CALCULUS OF VARIATIONS AND THE HAMILTON-JACOBI EQUATION

In the present section, we describe the main properties of viscosity solutions of the equation (HJc). These properties follow from the study of extremals via the Hopf-Lax-Oleinik formula, a global reference is the work of Fathi, [Fa1] and [Fa2]. We also state a result analogous to Theorem 1.9 for these solutions. The results in this section are standard or easy extensions of standard ones, however some of the proofs are original.

2.1. It is useful to introduce the Lagrangian $L : \mathbb{R} \times \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ associated to H . It is defined by

$$L(t, x, v) = \sup_p pv - H(t, x, p),$$

and has the following properties, which follow easily from the analogous properties 1.2 of H :

- i.* The Lagrangian L is C^2 and 1-periodic in t .
- ii.* The Lagrangian L is convex in v , and $\partial_{vv}^2 L > 0$.
- iii.* The Lagrangian has superlinear growth in v , *i.e.* $\lim_{|v| \rightarrow \infty} L(t, x, v)/|v| = \infty$ for each (t, x) .

The Lagrangian associated to the modified Hamiltonian $H(t, x, p + c)$ is $L(t, x, v) - cv$, it satisfies the three properties above.

2.2. For each c and each $t_0 \leq t$, we have the Hopf-Lax-Oleinik operator $V_{t_0, t}^c : C(\mathbb{T}, \mathbb{R}) \rightarrow C(\mathbb{T}, \mathbb{R})$ defined by

$$V_{t_0, t}^c(u)(x) = \min \left(u(x(t_0)) + \int_{t_0}^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds \right)$$

where the minimum is taken on the set of absolutely continuous curves $x : [t_0, t] \rightarrow \mathbb{T}$ such that $x(t) = x$. Any curve realizing the above minimum is C^2 and is the projection of a trajectory of X . More precisely, the curve $(x(s), \partial_v L(s, x(s), \dot{x}(s)))$ is a trajectory of X . These operators clearly satisfy the Markov property

$$V_{t_1, t_2}^c \circ V_{t_0, t_1}^c = V_{t_0, t_2}^c.$$

Note that these operators has been used in the study of viscosity solutions for quite a long time, see for instance [Fl]. We need the following:

DEFINITION A function $\tilde{u} : \mathbb{R}^n \rightarrow \mathbb{R}$ is called K -semi-concave if the function $\tilde{u} - K\|x\|^2/2$ is concave. The notion is extended to functions defined on convex subsets. A function $u : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ (resp. $\mathbb{T} \rightarrow \mathbb{R}$) is called K -semi-concave if it is of the form $u(t, x) = \tilde{u}(t, x \bmod 1)$ (resp. $u(x) = \tilde{u}(x \bmod 1)$), with a K -semi-concave function $\tilde{u} : [t_0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ (resp. $\tilde{u} : \mathbb{R} \rightarrow \mathbb{R}$). A function is called semi-concave if it can be written locally as the sum of a concave and a C^2 function.

PROPERTIES For each $c \in \mathbb{R}$ and each $t_0 < t_1$ we have :

- i. The operators V are equivariant under addition of a constant, more precisely, $V_{t_0, t_1}^c(U+u) = U + V_{t_0, t_1}^c(u)$ for all real constant U .
- ii. The operators V are non-decreasing, the inequality $V_{t_0, t_1}^c(u) \leq V_{t_0, t_1}^c(v)$ holds whenever the continuous functions u and v satisfy $u \leq v$.
- iii. The operators V are contractions, we have

$$\|V_{t_0, t_1}^c(u) - V_{t_0, t_1}^c(v)\|_\infty \leq \|u - v\|_\infty$$

for all continuous functions $u, v : \mathbb{T} \rightarrow \mathbb{R}$.

- iv. For each positive ϵ , there exists a constant K such that, for $|c| \leq 1/\epsilon$ and any continuous initial condition u_0 , the function

$$u(t, x) = V_{t_0, t}^c u_0(x)$$

is K -Lipschitz and K -semi-concave on $[t_0 + \epsilon, \infty) \times \mathbb{T}$.

2.3. PROPOSITION The following properties are equivalent for a function $u \in C([t_0, t_1] \times \mathbb{T}, \mathbb{R})$:

- i. The function u is a viscosity solution of (HJc) (see [Ba] or [Fa2] for the definition, which will not be used in the present paper).
- ii. The function u is locally Lipschitz on $]t_0, t_1[\times \mathbb{T}$ and it satisfies (HJc) almost everywhere. In addition, there exists a non-increasing function $K :]t_0, t_1[\rightarrow]0, \infty[$ such that the functions u_t is $K(t)$ -semi-concave for $t_0 < t \leq t_1$.
- iii. For each t and t' such that $t_0 \leq t \leq t' \leq t_1$, we have $u_{t'} = V_{t, t'}^c(u_t)$.

COROLLARY For each continuous initial condition u_{t_0} , there exists one and only one viscosity solution $u : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of (HJc) , given by

$$u(t, x) = V_{t_0, t}^c u_{t_0}(x).$$

There exists a non-increasing function $K(t) :]t_0, \infty[\rightarrow]0, \infty[$ and such that u is $K(t)$ -Lipschitz and $K(t)$ -semi-concave on $[t, \infty) \times \mathbb{T}$ for each $t > t_0$. In addition, given a constant $C > 0$, the function $K(t)$ can be chosen uniformly for all equations (HJc) with $|c| \leq C$.

PROOF. It is standard that $i. \iff iii.$, see a good exposition in [FMa]. Let us recall a possible sketch of the proof. One can first prove using variations around the maximum principle that there is at most one function satisfying $i.$ with a given initial condition u_{t_0} (see [Ba], 2.4.). On the other hand, it is obvious that there exists one and only one function satisfying $iii.$, namely $(t, x) \mapsto V_{t_0, t}^c u_{t_0}(x)$. One can prove, see [FMa], that this function also satisfies $i.$ It is then the only one to do so, by uniqueness. That $iii. \implies ii.$ results from properties 2.2. We shall prove more carefully that $ii. \implies i.$, which seems less classical. Let us fix $(S, Q) \in]t_0, \infty) \times \mathbb{T}$, it is enough to prove (see [Ba], 5.3) that all C^1 functions ϕ such that $u - \phi$ has a local minimum at (S, Q) , satisfy the equation at (S, Q) . If such a function ϕ exists, then $\partial_x u(S, Q)$ exists and is equal to $\partial_x \phi(S, Q)$. It follows from the Lemma below that u is differentiable at (S, Q) , and satisfies the equation at this point, which implies that ϕ also satisfies the equation at (S, Q) . \square

LEMMA Let $u(t, x)$ be a function satisfying condition $ii.$ of the proposition. If $(S, Q) \in]t_0, \infty) \times \mathbb{T}$ is a point where $\partial_x u$ exists, then the function u is differentiable and satisfies (HJc) at (S, Q) .

PROOF. Let us fix a time $t_2 \in]t_0, S[$. In view of the fact that all the functions $u_t, t \geq t_2$ are $K(t_2)$ -semi-concave, it is not hard to prove that $\partial_x u(s_n, q_n) \rightarrow \partial_x u(S, Q)$ when (s_n, q_n) is a sequence of points of differentiability of u converging to (S, Q) . If we assume in addition that (HJc) holds at (s_n, q_n) , we obtain that $\partial_t u(s_n, q_n)$ has a limit $H(t, x, c + \partial_x u(t, x))$. Let us denote L the linear form $(s, q) \mapsto q \partial_x u(S, Q) + s H(S, Q, \partial_x u(S, Q))$. We have proved that there exists a modulus of continuity δ and a set $K \subset \mathbb{R} \times \mathbb{T}$ of full measure in a neighborhood of (S, Q) such that, for each $(S + s, Q + q) \in K$, the function u is differentiable at $(S + s, Q + q)$ and $\|du(S + s, Q + q) - L\| \leq \delta(\|(s, q)\|)$. It follows that we have the estimate

$|u(S + s, Q + q) - u(S, Q) - L(s, q)| \leq \|(s, q)\| \delta(\|(s, q)\|)$ for all (s, q) small enough, hence u is differentiable at (S, Q) , and $du(S, Q) = L$. \square

2.4. Let $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) , and let $t_0 \leq t < t'$. An absolutely continuous curve $x(s) : [t, t'] \rightarrow \mathbb{T}$ is said to be calibrated by u if

$$u(t', x(t')) = u(t, x(t)) + \int_t^{t'} L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds.$$

If $x(s)$ is a calibrated curve, then it is C^2 and the curve $(x(s), \partial_v L(s, x(s), \dot{x}(s)))$ is a trajectory of the Hamiltonian vector field X . By extension, we say that a curve $\gamma(s) = (x(s), p(s)) : [t, t'] \rightarrow \mathbb{T} \times \mathbb{R}$ is calibrated by u if $x(s)$ is calibrated by u and if $p(s) = \partial_v L(s, x(s), \dot{x}(s))$. It is then a trajectory of X . A curve is said to be calibrated by u on an interval I if it is calibrated by u on $[t, t']$ for all $[t, t'] \subset I$. The following theorem is due to Albert Fathi.

THEOREM (CALIBRATED CURVES)

- i.* Let $x(s) :]t_0, \infty) \supset [t, t'] \rightarrow \mathbb{T}$ be a calibrated curve. For each $s \in]t, t'[$, the function u is differentiable at $(s, x(s))$, and $\partial_x u(s, x(s)) + c = p(s) = \partial_v L(s, x(s), \dot{x}(s))$
- ii.* For every $(t, x) \in]t_0, \infty) \times \mathbb{T}$ there exists a calibrated curve $x(s) : [t_0, t] \rightarrow \mathbb{T}$ such that $x(t) = x$. If u is differentiable at (t, x) , then there exists only one such calibrated curve, and it satisfies $\partial_x u(t, x(t)) + c = p(t) = \partial_v L(t, x(t), \dot{x}(t))$.
- iii.* If in addition the function u_{t_0} is semi-concave, and if $x(s) : [t_0, t] \rightarrow \mathbb{T}$ is a calibrated curve, then the function u_{t_0} is differentiable at $x(t_0)$, and $\partial_x u(t_0, x(t_0)) + c = p(t_0)$.

PROOF. We only sketch the proof. Details can be found in [Fa2] or [Fa1]. The calibrated curve of *ii.* is obtained as a minimizer of the functional

$$u(t_0, x(t_0)) + \int_{t_0}^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds$$

among the absolutely continuous curves $x(s) : [t_0, t] \rightarrow \mathbb{T}$ satisfying $x(t) = x$. The second point is Fathi's key regularity result. Let us mention the following Lemma from [Fa2].

LEMMA If $\gamma(s) = (x(s), p(s)) : [t, t'] \rightarrow \mathbb{T} \times \mathbb{R}$ is calibrated by u , then $p(t') - c$ is a proximal super-differential of $u_{t'}$ at $x(t')$, i.e.

$$u_{t'}(x(t') + x) \leq u_{t'}(x(t')) + (p(t') - c)x + O(x^2)$$

near $x = 0$, and $p(t) - c$ is a proximal sub-differential of u_t at $x(t)$, i.e.

$$u_t(x(t) + x) \geq u_t(x(t)) + (p(t) - c)x + O(x^2).$$

Using this Lemma, we complete the proof of the Theorem. If $\gamma(s) = (x(s), p(s)) : [t, t'] \rightarrow \mathbb{T} \times \mathbb{R}$ is calibrated by u , and if $s \in]t, t'[$, then the restrictions of γ to $[t, s]$ and $[s, t']$ are calibrated, hence $p(s) - c$ is both a sub and a super-differential of u_s at $x(s)$, hence it is the differential. This proves *i.* In addition, if $u_{t'}$ is differentiable at $x(t')$, then its only possible subdifferential at this point is $\partial_x u(t', x(t'))$, hence the calibrated curve has to satisfy the equation $\partial_x u(t', x(t')) + c = p(t') = \partial_v L(t', x(t'), \dot{x}(t'))$. By Cauchy-Lipschitz uniqueness, only one characteristic can satisfy this equation. This proves the second point in *ii.* The proof of *iii.* is similar: the existence of a calibrated curve emanating from a point provides a sub-differential, while semi-concavity provides a super-differential. \square

2.5. We are interested mainly in the asymptotic behavior of solutions. A first description is given by the following result. The function $\alpha(c)$ introduced below has been given several names, it is the α function of Mather, the averaged Hamiltonian, or the eigenvalue in terms of idempotent algebra.

PROPOSITION For each c , there exists a real number $\alpha(c)$ with the following property. For every solution $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of (HJc), the function $u(t, x) + t\alpha(c)$ is bounded on $[t_0, \infty) \times \mathbb{T}$.

From now on, we shall mainly consider the corrected equation

$$(Hc) \quad \partial_t u + H(t, x, c + \partial_x u) = \alpha(c).$$

PROOF. We work with a fixed parameter c . Let us define the sequences

$$M_n := \max_{x \in \mathbb{T}} V_{0,n}^c(0)(x) \text{ and } m_n := \min_{x \in \mathbb{T}} V_{0,n}^c(0)(x).$$

Since $V_{0,n}^c(0), n \geq 1$, are K -Lipschitz, we have

$$0 \leq M_n - m_n \leq K$$

for $n \geq 1$. We claim that $M_{n+m} \leq M_n + M_m$. This follows from the inequalities

$$V_{0,m+n}^c(0)(x) = V_{0,m}^c(V_{0,n}^c(0))(x) \leq V_{0,m}^c(M_n)(x) \leq M_n + V_{0,m}^c(0)(x).$$

Hence by a classical result on subadditive sequences, we have $\lim M_n/n = \inf M_n/n$. We denote by $-\alpha$ this limit. In the same way, the sequence $-m_n$ is subadditive, hence $m_n/n \rightarrow \sup m_n/n$. This limit is also $-\alpha$ because $0 \leq M_n - m_n \leq K$. As a consequence, we have, for all $n \geq 1$,

$$-K - n\alpha \leq m_n \leq -n\alpha \leq M_n \leq K - n\alpha.$$

Now let $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be any solution, and let t_1 be an integer greater than t_0 , then for all integer $n \in \mathbb{N}$ we obtain, for all x ,

$$\min_{\mathbb{T}} u_{t_1} - K \leq \min_{\mathbb{T}} u_{t_1} + m_n + n\alpha \leq u(n + t_1, x) + n\alpha \leq \max_{\mathbb{T}} u_{t_1} + M_n + n\alpha \leq \max_{\mathbb{T}} u_{t_1} + K.$$

Recalling that u is Lipschitz on $[t_1, \infty) \times \mathbb{T}$, we obtain the conclusion of the Proposition with $\alpha(c) = \alpha$. \square

2.6. The following Lemma is due to Jensen, the proof is from [BR]. The Corollary is fundamental, and by now classical.

LEMMA If $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ is a viscosity solution of (HJc), the function $v(\theta, x) : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$ defined by

$$v(\theta, x) = \liminf_{t \bmod 1 = \theta} (u(t, x) - t\alpha(c))$$

is a viscosity solution of the Hamilton-Jacobi equation

$$\partial_\theta v + H(\theta, x, c + \partial_x v) = \alpha(c),$$

where θ denotes $t \bmod 1$.

COROLLARY The Hamilton-Jacobi equation

$$\partial_\theta u + H(\theta, x, c + \partial_x u) = a$$

has a 1-periodic viscosity solution if and only if $a = \alpha(c)$.

PROOF OF THE LEMMA. We have to prove that $V_{s,t}^c(v_s) = v_t$ for all $s \leq t$. Let us first prove that $V_{s,t}^c(v_s) \leq v_t$. In order to do so, we fix (t, x) and consider an increasing sequence n_k of integers such that $u(t + n_k, x) \rightarrow v(t, x)$. There exists a sequence of curves $x_k : [s, t] \rightarrow \mathbb{T}$ which are calibrated by the $u(t + n_k, x)$ and satisfy $x_k(t) = x$. We have

$$u(t + n_k, x) = u(s + n_k, x_k(0)) + \int_s^t L(\sigma, x_k(\sigma + t), \dot{x}_k(\sigma + t)) d\sigma.$$

Let us assume, taking a subsequence is necessary, that the sequence x_k is converging to $x : [s, t] \rightarrow \mathbb{T}$ uniformly on compact sets. Taking the lim inf in the equality above gives

$$v(t, x) \geq v(s, x(0)) + \int_s^t L(\sigma, x(\sigma + t), \dot{x}(\sigma + t)) d\sigma \geq V_{s,t}^c(v_s)$$

We have used that the functions u_{s+n_k} have a common Lipschitz constant to conclude that $\liminf u(s + n_k, x_k(0)) = \liminf u(s + n_k, x(0)) \geq v(s, x(0))$.

In order to prove the reversed inequality, note that for all curves $x : [s, t] \rightarrow \mathbb{T}$, we have

$$u(t + n, x(t)) \leq u(s + n, x(s)) + \int_s^t L(\sigma, x(\sigma), \dot{x}(\sigma)) d\sigma.$$

Taking the lim inf, we obtain

$$v(t, x(t)) \leq v(s, x(s)) + \int_s^t L(\sigma, x(\sigma), \dot{x}(\sigma)) d\sigma$$

for each curve x , hence $V_{s,t}^c(v_s) \geq v_t$ which is the desired inequality. \square

2.7. PROPOSITION *The function $\alpha(c) : \mathbb{R} \rightarrow \mathbb{R}$ is convex, C^1 , and superlinear. For each viscosity solution $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of (HJc), there exist curves $\gamma : [t_0, \infty) \rightarrow \mathbb{T}$ which are calibrated by u . These curves all have the same rotation number*

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_{t_0}^t \dot{x} = \rho(c) := \alpha'(c).$$

PROOF. We begin with superlinearity. Let $v(\theta, x)$ be a periodic solution of

$$\partial_\theta v + H(\theta, x, c + \partial_x v) = \alpha(c).$$

Let $(\theta(c), x(c))$ be a minimum of v . The function v is semi-concave, hence differentiable at $(\theta(c), x(c))$, where $\partial_t v = 0 = \partial_x v$. It follows that

$$\alpha(c) = H(\theta(c), x(c), c)$$

and the superlinearity of α follows from the superlinearity of H .

Let us continue with convexity. For fixed t and x , the function $c \mapsto V_{0,t}^c(0)(x)$ is clearly concave, since it is defined as a minimum of linear functions. It follows that the function $c \mapsto -\alpha(c) = \lim_{t \rightarrow \infty} V_{0,t}^c(0)(x)/t$ is concave as a limit of concave functions, hence $\alpha(c)$ is a convex function.

LEMMA *Let $u_c : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) and let $x(t) : [t_0, \infty) \rightarrow \mathbb{T}$, be calibrated by u_c , then*

$$\alpha'(c-) \leq \liminf_{t \rightarrow \infty} \frac{1}{t} \int_{t_0}^t \dot{x}(s) ds \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \int_{t_0}^t \dot{x}(s) ds \leq \alpha'(c+).$$

PROOF. See [G] for related material. Let $u_c : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) and let $x(t) : [t_0, \infty) \rightarrow \mathbb{T}$, be calibrated by u_c . We have

$$u_c(t, x(t)) - u_c(t_0, x(t_0)) = \int_{t_0}^t L(s, x(s), \dot{x}(s)) ds - c \int_{t_0}^t \dot{x}(s) ds.$$

In view of the definition of $\alpha(c)$ in 2.5, the function $u_c(t, x) + t\alpha(c)$ is bounded; hence

$$\lim_{t \rightarrow \infty} \left(\frac{c}{t} \int_{t_0}^t \dot{x}(s) ds - \frac{1}{t} \int_{t_0}^t L(s, x(s), \dot{x}(s)) ds \right) = \alpha(c).$$

Let us now consider a viscosity solution $u_e : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of (HJe) . With the same curve $x(s)$, we have

$$u_e(t, x(t)) - u_e(t_0, x(t_0)) \leq \int_{t_0}^t L(s, x(s), \dot{x}(s)) ds - e \int_{t_0}^t \dot{x}(s) ds$$

hence

$$\alpha(e) \geq \limsup \left(\frac{e}{t} \int_{t_0}^t \dot{x}(s) ds - \frac{1}{t} \int_{t_0}^t L(s, x(s), \dot{x}(s)) ds \right),$$

which implies

$$\alpha(e) - \alpha(c) \geq \limsup \left(\frac{e-c}{t} \int_{t_0}^t \dot{x}(s) ds \right).$$

The desired inequalities follow at the limit. \square

We still have to prove that the function α is differentiable. We need some preliminaries.

2.8. Let \mathcal{C} be the set of continuous curves $x(t) : \mathbb{R} \rightarrow \mathbb{T}$ such that the mapping $t \mapsto \tilde{x}(t) = (t \bmod 1, x(t))$ with values in \mathbb{T}^2 is either one to one or periodic. Let us endow \mathcal{C} with the topology of uniform convergence on compact sets. The following result of Poincaré is very standard, and will not be proved here.

PROPOSITION *Each curve $x \in \mathcal{C}$ has a well defined real rotation number*

$$\rho(x) := \lim \frac{1}{t} \int_0^t \dot{x}(s) ds.$$

In addition, the mapping $x \mapsto \rho(x)$ is continuous on \mathcal{C} . If $x(t)$ and $y(t)$ are two curves of \mathcal{C} such that $\tilde{x}(\mathbb{R})$ and $\tilde{y}(\mathbb{R})$ are disjoint subsets of \mathbb{T}^2 , then they have the same rotation number.

2.9. LEMMA *Let $u(\theta, x)$ and $v(\theta, x)$ be two viscosity solutions of (Hc) of time period 1. Then there exists a curve $x(t) : \mathbb{R} \rightarrow \mathbb{T}$ which is calibrated both by u and v .*

It is usual in Aubry-Mather theory to define the Aubry set as the unions of all curves that are calibrated by all 1-periodic viscosity solutions of (Hc) . It is known that the Aubry set is not empty. This result of course implies the lemma, but we provide a simple self-contained proof.

PROOF. Let us define the number $M(t) := \max(v_t - u_t)$. The function $M(t)$ is non-increasing, because the inequality $v_t \leq M + u_t$ implies $v_s \leq M + u_s$ for all $s \geq t$, by Properties 2.2 *i* and *ii*. On the other hand the functions u and v are periodic, hence $M(t)$ is periodic, hence constant. We denote by M this constant. Let us fix a time t and a point x_t such that $v(t, x_t) = M(t) + u(t, x_t)$. Let $x(s) : (-\infty, 0] \rightarrow \mathbb{T}$ be a curve calibrated by u and such $x(t) = x_t$. Since

$$u(t, x(t)) = u(s, x(s)) + \int_s^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds$$

and

$$v(t, x(t)) \leq v(s, x(s)) + \int_s^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds$$

for all $s < t \leq 0$, we obtain

$$M \geq v(s, x(s)) - u(s, x(s)) \geq v(t, x(t)) - u(t, x(t)) \geq M,$$

hence the inequalities are all equalities, which imply that the curve $x(s)$ is also calibrated by v . We have found a curve $x(s) : (-\infty, 0] \rightarrow \mathbb{T}$ which is calibrated by u and v . We now consider a curve $\tilde{x}(s) : \mathbb{R} \rightarrow \mathbb{T}$ obtained as an accumulation point of the sequence $x(s - n), n \in \mathbb{N}$. This curve is calibrated by u and by v . \square

COROLLARY *There exists a real number $\rho(c) \in [\alpha'(c-), \alpha'(c+)]$ such that each curve $x : \mathbb{R} \rightarrow \mathbb{T}$ calibrated by a 1-periodic viscosity solution $u(\theta, x)$ of (HJc) has rotation number $\rho(c)$.*

PROOF. It is easy to see that, if x is calibrated by a periodic solution, then $x \in \mathcal{C}$; hence x has a rotation number. Let us consider two curves $x(t)$ and $q(t)$, such that x is calibrated by the 1-periodic solution $u(\theta, x)$ of (Hc) and q is calibrated by the 1-periodic solution $v(\theta, x)$ of (Hc) . In view of the lemma, there exists a third curve $f(t)$ which is calibrated both by u and v . Since x and f are both calibrated by u , the sets $\tilde{x}(\mathbb{R})$ and $\tilde{f}(\mathbb{R})$ are either disjoint or equal. In each case, we have $\rho(x) = \rho(f)$. In the same way, f and q are both calibrated by v hence $\rho(q) = \rho(f)$. As a consequence, we have $\rho(q) = \rho(x)$. Hence all the curves that are calibrated by some 1-periodic solution of (Hc) have the same rotation number, and we can denote by $\rho(c)$ this rotation number. It follows from Lemma 2.7 that $\rho(c) \in [\alpha'(c-), \alpha'(c+)]$. \square

2.10. We are going to prove that the function $\rho(c)$ is continuous. This implies the differentiability of α , and completes the proof of Proposition 2.7. Let us fix a value of c , and consider a sequence c_n converging to c . Let $v_n(\theta, x)$ be a time-periodic viscosity solution of (Hc_n) and let $x_n(t) : \mathbb{R} \rightarrow \mathbb{R}$ be a curve calibrated by v_n . We extract a subsequence k of n in such a way that the sequence v_k converges, uniformly on compact subsets of $\mathbb{R} \times \mathbb{T}$, to a viscosity solution v of the equation (Hc) . In addition, we can suppose that the curves x_k are converging uniformly on compact sets to a curve $x(t)$ which is calibrated by v . By the definition of $\rho(c)$, we have $\rho(x_k) = \rho(c_k)$ and $\rho(x) = \rho(c)$. By continuity of the rotation number on \mathcal{C} , we have $\rho(x_k) \rightarrow \rho(x)$, hence $\rho(c_k) \rightarrow \rho(c)$. This is the expected continuity of ρ . We have proved Proposition 2.7. \square

2.11. The asymptotic behavior of viscosity solutions is described by the following theorem, obtained in [Be] (see also [BR] for another proof, and see [Fa3] and [Ro] for related results). Let $T(c) \in \mathbb{N}$ be defined by:

$T(c)=1$ if $\rho(c)$ is irrational,

$T(c)=q$ if $\rho(c)$ is the rational p/q in lowest terms.

THEOREM *Let $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) . There exists a viscosity solution $v(t, x) : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ of (HJc) which satisfies $v_{t+T(c)} = v_t - T(c)\alpha(c)$ for each $t \in \mathbb{R}$, and*

$$\lim_{t \rightarrow \infty} \|u_t - v_t\|_{\infty} = 0.$$

3. ENTROPY SOLUTIONS AND CHARACTERISTICS

The relation between classical solutions of Burgers equation and the Hamiltonian dynamics is quite well understood from 1.5. We shall now describe the main properties of entropy solutions, with emphasis on their relation with dynamics. We will also prove Theorem 1.14 and Corollary 1.15.

3.1. A function $y :]t_0, t_1[\rightarrow \mathbb{R}$ is called an entropy solution of Burgers equation (B) if :

- i. The functions y and $H(t, x, y(t, x))$ are locally integrable and the equation holds in the sense of distributions :

$$\int_{[t_0, \infty) \times \mathbb{T}} y(t, x) \partial_t \phi(t, x) + H(t, x, y(t, x)) \partial_x \phi(t, x) dt dx = 0$$

for all smooth function $\phi :]t_0, t_1[\times \mathbb{T} \rightarrow \mathbb{R}$ with compact support (see [Se] for details). Note that the space average $\int_{\mathbb{T}} y(t, x) dx$ is then a constant c .

- ii. The Oleinik inequalities

$$y_t(x + \delta) - y_t(x) \leq K(t)\delta$$

hold for all $t \in]t_0, t_1[$, $x \in \mathbb{T}$ and $\delta > 0$, with a positive and decreasing function $K(t)$.

3.2. Let y_n be a sequence of functions of $L^1(\mathbb{T})$. Considering these functions as periodic functions on \mathbb{R} , we can define the primitives $Y_n(x) := \int_0^x y_n$. We say that the sequence y_n converges *very*

weakly to y if the primitives Y_n converge to Y , uniformly on compact sets.

PROPOSITION For each $t_0 \in \mathbb{R}$ and each $y_0 \in L^1(\mathbb{T})$, there exists a unique entropy solution $y :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of Burgers equation such that y_t converges very weakly⁽¹⁾ to y_0 when t converges to t_0 . This solution satisfies

$$y \in L^\infty([t_1, \infty) \times \mathbb{T}) \cap C(]t_0, \infty), L^1(\mathbb{T})),$$

for all $t_1 > t_0$, and it is given by $y(t, x) = c + \partial_x u(t, x)$, where $c = \int_{\mathbb{T}} y_0 dx$ and $u(t, x)$ is the viscosity solution of (HJc) of initial condition $u_{t_0}(x) = \int_0^x (y_0 - c)$.

PROOF. Let us first deal with uniqueness. The standard method to prove uniqueness is to use the Oleinik inequalities 3.1 *ii.*, via a duality method, see [H], Theorem 2.2.1, or [Se], 2.8. We shall use the Hamilton-Jacobi equation. Indeed, let $y(t, x) :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be an entropy solution. Note that this function is locally bounded in view of the Oleinik inequalities. Define

$$\tilde{u}(t, x) = \int_0^x (y(t, q) - c) dq - \int_0^1 \int_0^x (y(t, q) - c) dq dx,$$

where $c = \int_{\mathbb{T}} y_t dx$. We have, in the sense of distributions, $\partial_{tx} \tilde{u} = \partial_t y = -\partial_x (H(t, x, y))$. Hence the distribution $\partial_t \tilde{u}(t, x) + H(t, x, y(t, x))$ does not depend on x , and is the locally integrable function $f(t) = \int_{\mathbb{T}} H(t, x, y(t, x)) dx$. The function $u(t, x) = \tilde{u}(t, x) - \int_{t_0}^t f(s) ds$, satisfies $\partial_t u + H(t, x, y) = 0$ and $\partial_x u = y - c$ in the sense of distributions, hence both $\partial_x u$ and $\partial_t u$ are locally bounded functions on $]t_0, \infty) \times \mathbb{T}$ hence (see for example [EG], 4.2.3) the function u is locally Lipschitz on $]t_0, \infty) \times \mathbb{T}$. In addition, the Hamilton-Jacobi equation $\partial_t u + H(t, x, c + \partial_x u) = 0$ holds almost everywhere. The function u satisfies condition *ii.* of Proposition 2.3 as a consequence of the Oleinik inequalities *ii.* above. Since the functions y_t converge very weakly to y_0 as $t \rightarrow t_0$, we have $\tilde{u}_t \rightarrow u_{t_0}$ uniformly as $t \rightarrow t_0$, which implies that $u_t \rightarrow u_{t_0}$ uniformly. As a consequence, the function u is continuous on $[t_0, \infty) \times \mathbb{T}$. In view of 2.3, the function u has to be *the* viscosity solution of (HJc) of initial condition u_{t_0} . So the only candidate to be an entropy solution of (B) is $y(t, x) = c + \partial_x u(t, x)$.

It is classical to obtain the existence of entropy solutions as limits of regular solutions of the viscous equation (B_μ) . However we shall use once more the Hamilton Jacobi equation, *i.e.* we shall prove that the function $y(t, x) = c + \partial_x u(t, x)$ introduced in the discussion on uniqueness is indeed an entropy solution. Recall that u is Lipschitz on $[t_1, \infty) \times \mathbb{T}$ for all $t_1 \geq t_0$, hence $\partial_t u$ and $y = c + \partial_x u$ are well defined in $L^\infty([t_1, \infty) \times \mathbb{T})$, as well as $H(t, x, y(t, x))$. It is straightforward that *i.* is satisfied, and *ii.* follows from the property *ii.* of Proposition 2.3.

3.3. In order to prove the continuity of the mapping $t \mapsto y_t$ from $]t_0, \infty)$ to $L^1(\mathbb{T})$, let us introduce, for each positive number K , the set \mathcal{E}_K of functions which satisfy both

$$|y(x)| \leq K \text{ for all } x$$

and

$$y(x + \delta) \leq K\delta + y(x) \text{ for all } x \in \mathbb{T} \text{ and } \delta > 0.$$

LEMMA The set \mathcal{E}_K is relatively compact in $L^1(\mathbb{T})$ (for the strong topology). In addition, there exists a modulus of continuity $C_K(\epsilon)$ such that

$$d_H(y, z) \leq C_K(\|y - z\|_{L^1})$$

for $y, z \in \mathcal{E}_K$, where d_H is the distance defined in 1.12.

PROOF. Let us denote by $\tau_\delta y$ the function $x \mapsto y(x + \delta)$. We claim that, for each $\delta \in \mathbb{R}$, we have $\|\tau_\delta y - y\|_{L^1} \leq 2K|\delta|$. In view of Riesz-Fréchet-Kolmogorov compactness criterion (see [Br], IV.5) this implies the first part of the Lemma. In order to prove the claim, let us first suppose that $\delta > 0$. Using Oleinik inequality, we get $\tau_\delta y - y \leq K\delta$. Let us set $z^+(x) = \max\{0, \tau_\delta y(x) - y(x)\}$ and

⁽¹⁾The nature of the convergence of entropy solutions to their initial condition can be described more precisely depending on the regularity of the initial condition. This question, however, is not very relevant to our discussion.

$z^- = \max\{0, y(x) - \tau_\delta y(x)\}$, so that $\tau_\delta y - y = z^+ - z^-$. We have $z^+ \leq K\delta$, hence $\int_{\mathbb{T}} z^+ dx \leq K\delta$. Noticing that $\int_{\mathbb{T}} (\tau_\delta y - y) dx = 0$, we obtain that $\int_{\mathbb{T}} z^- dx = \int_{\mathbb{T}} z^+ dx$ hence

$$\|\tau_\delta y - y\|_{L^1} = \int_{\mathbb{T}} z^+ dx + \int_{\mathbb{T}} z^- dx \leq 2K\delta.$$

The inequality for $\delta < 0$ follows in the same way from the fact that $\tau_\delta y - y \geq K\delta$ in this case.

We now prove the second part of the Lemma. See [KO] for related material. Let us consider two functions y and z of \mathcal{E}_K and assume that $d_H(y, z) \geq 2\epsilon > 0$. There exists a point $(x_0, y_0) \in H(y)$ (see 1.12 for the notation) such that the ball B of radius 2ϵ centered at (x_0, y_0) in $\mathbb{T} \times \mathbb{R}$ does not contain any point of $H(z)$. Hence either $z(x) \geq y_0 + \epsilon$ for almost all $x \in [x_0 - \epsilon, x_0 + \epsilon]$ or $z(x) \leq y(x_0) - \epsilon$ for almost all $x \in [x_0 - \epsilon, x_0 + \epsilon]$. We will treat the first case, the second being similar. Using Oleinik inequality, we obtain, for each $\delta \in [0, \epsilon]$,

$$z(x_0 + \delta) \geq y_0 + \epsilon \geq y(x_0 + \delta) - K\delta + \epsilon.$$

Hence

$$\|y - z\|_{L^1} \geq \int_{x_0}^{x_0 + \min\{\epsilon, \epsilon/K\}} z(x) - y(x) dx \geq \int_0^{\min\{\epsilon, \epsilon/K\}} \epsilon - Kx dx \geq \epsilon^2 \max\left(\frac{1}{2}, \frac{1}{2K}\right).$$

□

COROLLARY *Let $\mathcal{V}_K \subset C(\mathbb{T}, \mathbb{R})$ be the set of K -semi-concave (hence K -Lipschitz) functions. Then every function $u \in \mathcal{V}_K$ has a derivative $\partial_x u$ in \mathcal{E}_K , and the operator $\partial_x : \mathcal{V}_K \rightarrow \mathcal{E}_K$ is uniformly continuous, when \mathcal{V}_K is endowed with the topology of uniform convergence and \mathcal{E}_K with the strong L^1 topology.*

PROOF. In order to prove that the operator $\partial_x : \mathcal{V}_K \rightarrow \mathcal{E}_K$ is uniformly continuous, let us consider two sequences u_n and v_n of \mathcal{V}_K , such that $\|u_n - v_n\|_\infty \rightarrow 0$. We have to prove that $\|\partial_x u_n - \partial_x v_n\|_{L^1} \rightarrow 0$. If not, there would exist subsequences u_k and v_k such that $\partial_x u_k$ and $\partial_x v_k$ have different limits y and z in L^1 . This implies that the sequences u_k and v_k have limits u and v in $C(M, \mathbb{R})$, and that these limits are different. But this is obviously in contradiction with the assumption that $\|u_n - v_n\|_\infty \rightarrow 0$. □

In view of this corollary, the continuity of $t \mapsto y_t$ follows from the continuity of $t \mapsto u_t$. □

3.4. It is clear from what has just been written that Theorem 2.11 implies our main new results, Theorems 1.9 and 1.14. Indeed, if $y(t, x)$ is an entropy solution of (B) , we write it $y = c + \partial_x u$, where u is a viscosity solution of (HJc) . Theorem 2.11 gives the existence of a viscosity solution v of (HJc) such that $v(t + T(c), x) = v(t, x) - T(c)\alpha(c)$ and $\|v_t - u_t\|_\infty \rightarrow 0$ as $t \rightarrow \infty$. It follows from the considerations above that the function $\omega(t, x) = c + \partial_x v(t, x)$ is an entropy solution of (B) , and it clearly satisfies $\omega(t + T(c), x) = \omega(t, x)$. In addition, for $t_1 > t_0$, all the functions $u_t, t \geq t_1$ and $v_t, t \in \mathbb{R}$ belong to the same set \mathcal{V}_K . We obtain that $\|y_{t_1} - \omega_{t_1}\|_{L^1} = \|\partial_x u_{t_1} - \partial_x v_{t_1}\|_{L^1} \rightarrow 0$ as $t \rightarrow \infty$ in view of the uniform continuity of the operator $\partial_x : \mathcal{V}_K \rightarrow \mathcal{E}_K$. The second part of Lemma 3.3, then implies that $d_H(y_t, \omega_t) \rightarrow 0$. □

3.5. Consider an entropy solution $y(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of Burgers equation, let $c = \int_{\mathbb{T}} y dx$ and let $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) such that $y = c + \partial_x u$. A trajectory $\gamma(s) : [t_0, \infty) \supset [t, t'] \rightarrow \mathbb{T} \times \mathbb{R}$ of the Hamiltonian vector-field X is called a y -characteristic if $\gamma(s) \in \overline{G(y_s)}$ for each $s \in [t, t']$. A curve $x(s) : [t_0, \infty) \supset [t, t'] \rightarrow \mathbb{T}$ is also called a y -characteristic if it is the projection of a y -characteristic $\gamma(s)$. The following theorem extends the method of characteristics to entropy solutions :

THEOREM (CHARACTERISTICS)

- i. A curve $\gamma(s) : [t_0, \infty) \supset [t, t'] \rightarrow \mathbb{T} \times \mathbb{R}$ is a y -characteristic if and only if it is calibrated by u .*

- ii. Let $\gamma(s) :]t_0, \infty) \supset]t, t' [\longrightarrow \mathbb{T} \times \mathbb{R}$ be a y -characteristic, then for each $s \in]t, t' [$, the function y_s is continuous at $x(s)$ and $y(s, x(s)) = \overline{p(s)}$
- iii. For every $t > t_0$ and every $(x, p) \in \overline{G(y_t)}$, there exists a unique y -characteristic $\gamma(s) :]t_0, t] \longrightarrow \mathbb{T} \times \mathbb{R}$ such that $\gamma(t) = (x, p)$.
- iv. If in addition $y_{t_0} \in \mathcal{E}$, then for every $t > t_0$ and every characteristic $x :]t_0, t[\longrightarrow \mathbb{T}$, we have $p(t_0) = y(t_0, x(t_0))$. Where $(x(t_0), p(t_0))$ is naturally defined by prolongation of the Hamiltonian trajectory $(x(t), p(t))$.
- v. There exists a characteristic $x(t) :]t_0, \infty) \longrightarrow \mathbb{T}$.

PROOF. The proposition will be deduced from the properties of calibrated curves mentioned in 2.4. If $\gamma(s) = (x(s), p(s)) : [t, t'] \longrightarrow \mathbb{T} \times \mathbb{R}$ is a calibrated curve, then we have seen that y is continuous at $(s, x(s))$ for each s of $]t, t' [$, and that $y(s, x(s)) = p(s)$. As a consequence, calibrated curves are characteristics, and *ii.* holds for these curves.

Let us now prove *iii.* Let us fix a point $(x, p) \in \overline{G(y_t)}$, and let (x_n, p_n) be a sequence of points of $G(y_t)$ converging to (x, p) . For each n , there exists a calibrated curves $(x_n(s), p_n(s)) : [t_0, t] \longrightarrow \mathbb{T} \times \mathbb{R}$ such that $x_n(t) = x_n$. Since (t, x_n) is a point of differentiability of u , this calibrated curve satisfies $p_n(t) = \partial_x u(t, x_n) + c = p_n$. These calibrated curves converge to a calibrated curve which terminates at (t, x, p) . Uniqueness in *iii.* is a consequence of Cauchy-Lipschitz Theorem for the Hamiltonian vectorfield.

We now prove that all characteristics are calibrated curves. Let $\gamma(s) = (x(s), p(s)) : [t, t'] \longrightarrow \mathbb{T} \times \mathbb{R}$ be a characteristic. Let $\tilde{\gamma}(s) : [t_0, t'] \longrightarrow \mathbb{T} \times \mathbb{R}$ be the unique characteristic terminating at $(t', x(t'), p(t'))$ which was obtained above (and which is a calibrated curve). Clearly, γ has to be the restriction of $\tilde{\gamma}$, hence it is a calibrated curve. \square

3.6. In terms of the dynamics, using the notation of 1.12, this theorem implies that

$$\overline{G(y_{t'})} \subset \phi_{t,t'}(G(y_t))$$

when $t_0 < t < t'$, and that

$$\overline{G(E(y))} \subset \phi(G(y))$$

when $y \in \mathcal{E}$.

3.7. A function $y :]t_0, t_1[\times \mathbb{T} \longrightarrow \mathbb{R}$ is called a backward entropy solution of (B) if the function $\check{y}(s, x) := -y(t_1 - s)$ is an entropy solution of the reversed equation

$$(\check{B}) \quad \partial_t \check{y} + \partial_x (\check{H}(t, x, y)) = 0,$$

where $\check{H}(t, x, y) = H(t_1 - t, x, -y)$. If y is a C^1 solution of Burgers equation, then it is an entropy solution and a backward entropy solution. However, entropy solutions and backward entropy solutions are different in general. More precisely, the function y is a backward entropy solution if and only if it is a solution in the sense of distributions, and satisfies the reversed Oleinik inequalities

$$y_t(x + \delta) \geq y_t(x) - K(t)\delta$$

when $\delta > 0$, with an increasing function $K :]t_0, t_1[\longrightarrow]0, \infty)$. As a consequence, if y is both an entropy solution and a backward entropy solution, then y_t is Lipschitz for each $t \in]t_0, t_1[$. More precisely, we have:

PROPOSITION *Let $y :]t_0, \infty) \times \mathbb{T} \longrightarrow \mathbb{R}$ be an entropy solution of (B) . The function y is a backward entropy solution of (B) on $]t_0, t_1[\times \mathbb{T}$ if and only if the function y_{t_1} is continuous. The function y is then locally Lipschitz on $]t_0, t_1[\times \mathbb{T}$.*

PROOF. Let us first assume that y is both an entropy solution and a backward entropy solution. Then for each compact interval of time $I \subset]t_0, t_1[$, the functions $y_t, t \in I$, are equiLipschitz. In view of the Lemma in 2.3, we conclude that y is Lipschitz on I .

Assume now that y_{t_1} is continuous. Then each point $(x, y_{t_1}(x))$ is the endpoint of a single characteristic $\Gamma_x(t) = (Q(t, x), P(t, x)) :]t_0, t_1] \longrightarrow \mathbb{T}$. Clearly, we have $y(t, Q(t, x)) = P(t, x)$ for all $(t, x) \in]t_0, t_1] \times \mathbb{T}$. Let $z(t, x) : (-\infty, t_1] \times \mathbb{T} \longrightarrow \mathbb{R}$ be the unique backward entropy solution

with final condition y_{t_1} given by Proposition 3.2. Let us fix a point $(t, x) \in [t_0, t_1] \times \mathbb{T}$, and a z -characteristic $\gamma(s) = (x(s), p(s)) : [t, t_1] \rightarrow \mathbb{T} \times \mathbb{R}$ satisfying $x(t) = x$. Since $-y_{t_1} \in \mathcal{E}$, this characteristic furthermore satisfies $p(t_1) = z(t_1, x(t_1))$ (this results from *iv.* of the theorem on characteristics). As a consequence, the characteristic $\gamma(s)$ coincides with $\Gamma_{x(t_1)}(s)$. This implies that z_t is continuous at x and satisfies

$$z(t, x) = p(t) = P(t, x(t_1)) = y(t, Q(t, x(t_1))) = y(t, x).$$

We have proved that $y_t = z_t$ for $t \in]t_0, t_1[$. □

3.8. PROPOSITION *Let $y, z :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be two continuous entropy solutions of (B). Then the function $t \mapsto \|y_t - z_t\|_{L^1}$ is constant.*

This result will be extended in section 4, where we give a proof of the well known fact that the function $t \mapsto \|y_t - z_t\|_{L^1}$ is non-increasing if y and z are entropy solutions. Since continuous solutions are both entropy solutions and backward entropy solutions, this implies the proposition. However, we give an independent proof here.

PROOF. Let us consider the graphs Γ_{y_t} and Γ_{z_t} of y and z in $\mathbb{T} \times \mathbb{R}$. Let U_t be the domain enclosed by these two curves, that is the union of all bounded connected components of $\mathbb{T} \times \mathbb{R} - (\Gamma_{y_t} \cup \Gamma_{z_t})$. Clearly, $\|y_t - z_t\|_{L^1}$ is the area of U_t . It follows from the theorem on characteristics that $U_t' = \phi_{t,t'}(U_t)$ for $t' > t > t_0$, where ϕ is the Hamiltonian flow. The area of U_t does not depend on t because the flow is area-preserving. □

3.9. We are now in a position to prove Corollary 1.9. Let us first state it in slightly greater generality.

COROLLARY *Let $y(t, x) :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a solution of Burgers equation. If there exists an unbounded sequence $t_n > t_0$ of times such that the function y_{t_n} is continuous for all n , then the function y is the restriction of a Lipschitz solution $\omega : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ which satisfies $\omega(t + T, x) = \omega(t, x)$ for some $T \in \mathbb{N}$.*

PROOF. It follows from Proposition 3.7 that the function y is both a forward and a backward entropy solution on $]t_0, t_n[\times \mathbb{T}$ for each n . It follows from the Oleinik inequality and the reversed Oleinik inequality that the function y is Lipschitz on $[t'_0, \infty) \times \mathbb{T}$ for all $t'_0 > t_0$. On the other hand, there exists a periodic solution $\omega(t, x)$ such that $\|y_t - \omega_t\|_{L^1} \rightarrow 0$ as $t \rightarrow \infty$. The functions ω_t are equilipshitz because the functions $y_t, t \geq t'_0$ are equilipshitz. As a consequence, both entropy solutions ω and y are continuous on $[t'_0, \infty) \times \mathbb{T}$, hence $\|y_t - \omega_t\|_{L^1}$ is constant, hence it is zero. We have proved that $y = \omega$ on $[t_1, \infty) \times \mathbb{T}$. □

3.10. The fact that backward characteristics always exist, but that forward characteristics don't, is one of the key features of entropy solutions. Particle may be absorbed by shocks, but they can't be created by shocks. In order to understand the full future of a given particle, it is useful to introduce the notion of weak characteristic. Consider an entropy solution $y(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ of Burgers equation, let $c = \int_{\mathbb{T}} y dx$ and let $u(t, x) : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solution of (HJc) such that $y = c + \partial_x u$. We say that an absolutely continuous curve $x(s) : [t_0, \infty) \supset [t, t'] \rightarrow \mathbb{T}$ is a weak y -characteristic if it satisfies the equation

$$\dot{x}(s) \in [\partial_p H(s, x(s), y_s^-(x(s))), \partial_p H(s, x(s), y_s^+(x(s)))]$$

almost everywhere. It is clear that y -characteristics are weak y -characteristics, and that the notions of weak y characteristics and of y -characteristics coincide if y is a continuous solution.

PROPOSITION *For each $t > t_0$ and $x_0 \in \mathbb{T}$, there exists one and only one weak characteristic $x(s) : [t, \infty) \rightarrow \mathbb{T}$ satisfying $x(t) = x_0$.*

PROOF. In view of the convexity of H in the p variable, there exists a constant K such that the

function $Y(s, x) = \partial_p H(s, x, y(s, x))$ satisfies $|Y(s, x)| \leq K$ and the Oleinik inequality

$$Y(s, x + \delta) \leq Y(s, x) + K\delta$$

for $\delta > 0$, $s \geq t$. Since $\partial_p H(s, x, y^\pm(s, x)) = Y^\pm(s, x)$, the equation for weak characteristics can be written as

$$\dot{x}(s) \in [Y^-(x(s)), Y^+(x(s))].$$

Now the Proposition follows from a result of Filippov, see [H], Theorem 1.4.2. \square

3.11. PROPOSITION *All weak y -characteristics $x(s) : [t, \infty) \rightarrow \mathbb{T}$ have the same rotation number*

$$\lim \frac{1}{t} \int_{t_0}^t \dot{x} = \rho(c) = \alpha'(c).$$

PROOF. Let $x(t) :]t_0, \infty) \rightarrow \mathbb{T}$ and $q(t) :]t_0, \infty) \rightarrow \mathbb{T}$ be respectively a genuine y -characteristic and a weak y -characteristic. Let $\bar{x}(t) :]t_0, \infty) \rightarrow \mathbb{R}$ and $\bar{q}(t) :]t_0, \infty) \rightarrow \mathbb{R}$ be continuous functions such that $\bar{x}(t) \bmod 1 = x(t)$ and $\bar{q}(t) \bmod 1 = q(t)$, and such that $\bar{x}(t_0) \leq \bar{q}(t_0) \leq \bar{x}(t_0) + 1$. Note that $\bar{q}(t) = \bar{q}(t_0) + \int_{t_0}^t \dot{q}$, with a similar property for \bar{x} . The uniqueness of weak y -characteristics implies that $\bar{x}(t) \leq \bar{q}(t) \leq \bar{x}(t) + 1$ for all t . Since x is a genuine y -characteristic, it is calibrated by u , so that Proposition 2.7 implies

$$\frac{\bar{x}(t)}{t} \rightarrow \rho(c) \text{ and hence } \frac{\bar{q}(t)}{t} \rightarrow \rho(c).$$

\square

4. PAIRS OF SOLUTIONS AND DISSIPATION

It is well known that if y and z are entropy solutions of (B) , then the norm $\|y_t - z_t\|_{L^1}$ is a non-increasing function of t . We shall give a proof of this fact. In short, it is a consequence of the fact that values of local maxima of differences of viscosity solutions are decreasing. We now give the details.

4.1. Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be a continuous function. We say that $x_0 \in \mathbb{T}$ is a point of local maximum of f if there exists an interval $]a, b[\subset \mathbb{T}$ containing x_0 and such that $f(x_0) = \max_{]a, b[} f$ and $f(x_0) > \max(f(a), f(b))$. The value $f(x_0)$ is then called a value of local maximum. The following proposition states that the values of local maxima of a difference $u_t - v_t$ of viscosity solutions are non-increasing functions of t .

PROPOSITION *Let $u, v : [t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a pair of viscosity solutions of (HJc) , let $t_1 > t_0$, and let $d = u - v$. Let x_1 be a point of local maximum for d_{t_1} , and let $]a, b[$ be an open interval containing x_1 , and such that $\max(d_{t_1}(a), d_{t_1}(b)) < d(t_1, x_1) = \max_{]a, b[} d_{t_1}$. Let $a(t)$ and $b(t)$ be curves calibrated by u and satisfying $a(t_1) = a$, $b(t_1) = b$. Then for each t in $]t_0, t_1]$, we have $M(t) = \max_{[a(t), b(t)]} d_t > \max(d(t, a(t)), d(t, b(t)))$, hence $M(t)$ is a value of local maximum. In addition, the value $M(t)$ is a continuous and non-increasing function of t on $]t_0, t_1]$.*

PROOF. The functions $d(t, a(t))$ and $d(t, b(t))$ are non-decreasing on $[t_0, t_1]$; hence $d(t, a(t)) \leq d(t_1, a(t_1)) < M(t_1)$ and $d(t, b(t)) \leq d(t_1, b(t_1)) < M(t_1)$. On the other hand, consider a curve $q(t)$ which is calibrated by v and satisfies the final condition $q(t_1) = x_1$. Since the function $d(t, q(t))$ is non-increasing, we have $d(t, q(t)) \geq d(t_1, q(t_1)) = M(t_1)$. It follows that $d(t, q(t)) > d(t, a(t))$ and $d(t, q(t)) > d(t, b(t))$ for each t , hence the curve $q(t)$ can't cross the curves $a(t)$ and $b(t)$, and $q(t) \in]a(t), b(t)[$ for each t . As a consequence, we have $M(t) \geq d(t, q(t)) > \max(d(t, a(t)), d(t, b(t)))$ hence $M(t)$ is a value of local maximum. In addition, we have $M(t) \geq d(t, q(t)) \geq M(t_1)$. The same construction can be performed with a smaller final time t'_1 satisfying $t < t'_1 \leq t_1$, hence we have the inequality $M(t) \geq M(t'_1)$ for $t < t'_1 \leq t_1$. As a consequence, the function $M(t)$ is non-increasing on $]t_0, t_1]$. \square

4.2. In order to apply the proposition to pairs of entropy solution of (B) with possibly different averages, we need some refinements. From now on, we sometimes consider functions on \mathbb{T} as periodic functions on \mathbb{R} without changing their names.

COROLLARY *Let c and c' be two real numbers, let $u :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solutions of (HJc) , let $v :]t_0, \infty) \times \mathbb{T} \rightarrow \mathbb{R}$ be a viscosity solutions of (HJc') , and let $t_1 > t_0$. We denote by d_t the difference $d_t(x) = u_t(x) - v_t(x) + (c - c')x : \mathbb{R} \rightarrow \mathbb{R}$. Let $x_1 \in \mathbb{R}$ be a point of local maximum for d_{t_1} , and let $]a, b[\subset \mathbb{R}$ be an open interval of length less than 1 containing x_1 , and such that $\max(d_{t_1}(a), d_{t_1}(b)) < d(t_1, x_1) = \max_{]a, b[} d_{t_1}$. Let $a(t)$ and $b(t)$ be curves calibrated by u (for (HJc)) and satisfying $a(t_1) = a$, $b(t_1) = b$. Then there exists $t'_0 < t_1$ such that, for $t \in [t'_0, t_1]$, we have $M(t) = \max_{]a(t), b(t)[} d_t > \max(d(t, a(t)), d(t, b(t)))$. The value $M(t)$ is a continuous and non-increasing function of t on $[t'_0, t_1]$. The functions $t \mapsto d(t, a(t))$ and $t \mapsto d(t, b(t))$ are non-decreasing on $[t'_0, t_1]$.*

ADDENDUM *If $]a', b'[\subset]a, b[$ is a smaller interval and if $M' = d(t_1, x'_1) = \max_{]a', b'[} d_{t_1} > \max(d_{t_1}(a'), d_{t_1}(b'))$, then the Corollary can be applied to the value M' in the interval $]a', b'[$ with the same time t'_0 .*

PROOF. Let us fix intervals I and J of length less than 1 such that $[a, b] \subset I \subset \bar{I} \subset J$. We also call I and J the images of I and J in \mathbb{T} , which are proper subsets of \mathbb{T} . Let K' be a common Lipschitz constant of the functions $v_t, t \geq (t_0 + t_1)/2$, and let $K = 2K'/(1 - |J|)$. Let us fix a time $t'_0 < t_1$ and a 1-periodic K -Lipschitz function $w_{t'_0} : \mathbb{R} \rightarrow \mathbb{R}$ such that $w_{t'_0}(x) = v(t'_0, x) + (c' - c)x$ for $x \in \mathbb{Z} + J$. We can see $w_{t'_0}$ as a function on \mathbb{T} , and define in a natural way $w(t, x) = V_{t'_0, t}^c(w_{t'_0})(x)$, for $t \geq t'_0$. We claim that $w(t, x) = v(t, x) + (c' - c)x$ for all $x \in I$ and $t \in [t'_0, t_1]$, if $t'_0 < t_1$ is large enough. Assuming the claim, and assuming in addition that t'_0 is sufficiently large that the curves $a(t)$ and $b(t)$ remain in I on $[t'_0, t_1]$, the corollary follows from the proposition above applied to u and w , which both solve (HJc) . In order to prove the claim, recall that

$$w(t, x) = \min \left(w_{t'_0}(x(t'_0)) + \int_{t'_0}^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds \right),$$

and

$$v(t, x) = \min \left(v_{t'_0}(x(t'_0)) + \int_{t'_0}^t L(s, x(s), \dot{x}(s)) - c'\dot{x}(s) ds \right),$$

where the minima are taken on the set of C^1 curves $x(s) : [t'_0, t] \rightarrow \mathbb{R}$ terminating at x . By superlinearity of L , it is possible to choose t'_0 so close to t_1 that, if $t'_0 \leq t \leq t' \leq t_1$ and $x(s) : [t, t'] \rightarrow \mathbb{R}$ is a curve starting outside of J and ending inside I , then

$$\int_t^{t'} L(s, x(s), \dot{x}(s)) ds \geq 2K + \int_t^{t'} L(s, x(t'), 0) ds + (|c| + |c'|) \int_t^{t'} |\dot{x}(s)| ds.$$

If t'_0 is chosen sufficiently close to t_1 , then for each $x \in I$, the curves reaching the minima in the expressions of w and v above satisfy $x(s) \in J$ for all s . We have proved that, if t'_0 is large enough, then for each $x \in I$ it is possible to restrict the minima in the expression for w and v above to curves $x(s)$ contained in J . For such a curve, we have

$$\begin{aligned} & w_{t'_0}(x(t'_0)) + \int_{t'_0}^t L(s, x(s), \dot{x}(s)) - c\dot{x}(s) ds \\ &= v_{t'_0}(x(t'_0)) + (c' - c)x(t'_0) + c(x(t'_0) - x) + \int_{t'_0}^t L(s, x(s), \dot{x}(s)) ds \\ &= v_{t'_0}(x(t'_0)) + (c' - c)x + \int_{t'_0}^t L(s, x(s), \dot{x}(s)) - c'\dot{x}(s) ds. \end{aligned}$$

As a consequence, the equality $w(t, x) = (c' - c)x + v(t, x)$ holds for all $t \in [t'_0, t_1]$ and all $x \in J$. This ends the proof of the claim, and the proof of the corollary. \square

4.3. THEOREM *Let $y(t, x)$ and $z(t, x) :]t_0, \infty) \times \mathbb{T} \longrightarrow \mathbb{R}$ be two entropy solutions of (B). Then the function*

$$t \longmapsto \|y_t - z_t\|_1$$

is non-increasing. In addition, let $x_0(t)$ and $x_1(t)$ be two curves which are characteristics both for y and z , and let $J(t)$ and $I(t)$ be the two continuously varying intervals of \mathbb{T} such that $I(t) \cup J(t) = \mathbb{T}$ and $I(t) \cap J(t) = \{x_0(t), x_1(t)\}$. Then each of the functions

$$t \longmapsto \|(y_t - z_t)|_{I(t)}\|_1 \quad \text{and} \quad t \longmapsto \|(y_t - z_t)|_{J(t)}\|_1$$

is non-increasing.

PROOF. We shall prove the first part, leaving the easy modifications needed to prove the second part to the reader. Let $y = c + \partial_x u$ and $z = c' + \partial_x v$ be two entropy solutions of (B), and let $d(t, x) = u_t(x) - v_t(x) + (c - c')x$. The idea of the proof is very simple. The L^1 norm of $y_t - z_t$ is the total variation of the function d_t . Since the local maxima of this function do not increase, and since its local minima do not decrease, the total variation of d_t can't increase. The details are, however, a bit tedious.

We consider u and v as periodic functions $]t_0, \infty) \times \mathbb{R} \longrightarrow \mathbb{R}$, and define $d(t, x) = u_t(x) - v_t(x) + (c - c')x$. We denote by $\text{Var}(d_t)$ the total variation of d_t on any interval of length 1. This quantity does not depend on the interval, and

$$\text{Var}(d_t) = \|y_t - z_t\|_{L^1(\mathbb{T})}.$$

We shall prove that, for each $t_1 > t_0$, there exists a time $t'_0 \in]t_0, t_1[$ such that, for each $t \in]t'_0, t_1[$, we have $\text{Var}(d_t) \geq \text{Var}(d_{t_1})$. Because it is also continuous, the function $t \longmapsto \text{Var}(d_t)$ is then non-increasing, which is the desired result.

Let us fix a time $t_1 > t_0$. Assume first that the function d_{t_1} does not have any local minimum. In this case, $\text{Var}(d_{t_1}) = |c' - c|$. On the other hand, the inequality $\text{Var}(d_t) \geq |c' - c|$ clearly holds for all t , hence we have proved that

$$\text{Var}(d_t) \geq \text{Var}(d_{t_1})$$

for all $t < t_1$.

Otherwise, there exists a point $a \in \mathbb{R}$ of local minimum of d_{t_1} . We consider a y -characteristic $a(t) :]t_0, t_1] \longrightarrow \mathbb{R}$ terminating at a . There exist non decreasing sequences \mathcal{M}_k^- and \mathcal{M}_k^+ of finite subsets of $]a, a + 1[$ such that, for each k :

- Each point of \mathcal{M}_k^- is a point of local minimum of d_{t_1} ;
- Each maximal interval in the complement $]a, a + 1[- \mathcal{M}_k^-$ of \mathcal{M}_k^- contains exactly one point of \mathcal{M}_k^+ ;
- Each point of \mathcal{M}_k^+ is a point of local maximum of d_{t_1} ;
- The sets \mathcal{M}_k^- and \mathcal{M}_k^+ are disjoint.
- We have, when $k \longrightarrow \infty$,

$$2 \sum_{x \in \mathcal{M}_k^+} d_{t_1}(x) - 2 \sum_{x \in \mathcal{M}_k^-} d_{t_1}(x) - d_{t_1}(a(t_1)) - d_{t_1}(a(t_1) + 1) \longrightarrow \text{Var}(d_{t_1}).$$

It is possible that \mathcal{M}_k^- is empty for all k . In this case, \mathcal{M}_k^+ is also independent of k and contains a single point x , and we have

$$\text{Var}(d_{t_1}) = 2d_{t_1}(x) - d_{t_1}(a(t_1)) - d_{t_1}(a(t_1) + 1).$$

In view of the corollary, there exists t'_0 such that the function $M(t) = \max_{]a(t), a(t+1)[} d_t$ is non-increasing on $[t'_0, t_1]$, and the functions $d(t, a(t))$ and $d(t, b(t))$ are non-decreasing on this interval. As a consequence, for $t \in]t_0, t_1[$, we have

$$\text{Var}(d_t) \geq 2M(t) - d_t(a(t)) - d_t(a(t) + 1) \geq 2M(t_1) - d_{t_1}(a(t_1)) - d_{t_1}(a(t_1) + 1) = \text{Var}(d_{t_1}).$$

Otherwise, the function d_{t_1} has oscillations and the sets \mathcal{M}_k^- are not empty. Let us choose, for each point $x \in \mathcal{M}_k^-$, a y -characteristic $x(t) : [t_0, t_1] \longrightarrow \mathbb{R}$ satisfying $x(t_1) = x$. Let $\mathcal{M}_k^-(t)$ be the union of all the points $x(t)$ obtained that way. The corollary above and its addendum allow to find a time $t'_0 \in]t_0, t_1[$ independent of k such that, if $]x, x'[\subset]a, a + 1[$ is a maximal interval of

the complement of \mathcal{M}_k^- , then the y -characteristics $x(t)$ and $x'(t)$ associated to x and x' satisfy $a(t) \leq x(t) < x'(t) \leq a(t) + 1$ for each $t \in]t'_0, t_1[$, and the interval $]x(t), x'(t)[$ contains a point of local maximum $q(t)$ of d_t , such that the associated value of local maximum $d(t, q(t))$ is non-increasing. Let $\mathcal{M}_k^+(t)$ be the union of all such points $q(t)$. Recall also, from the corollary, that the function $t \mapsto d(t, x(t))$ is non-decreasing on $]t'_0, t_1[$ for each of the chosen characteristics $x(t) \in \mathcal{M}_k^-(t)$. We have, for $t \in [t'_0, t_1]$

$$\begin{aligned} \text{Var}(d_t) &\geq 2 \sum_{q \in \mathcal{M}_k^+(t)} d_t(q) - 2 \sum_{x \in \mathcal{M}_k^-(t)} d_t(x) - d_t(a(t)) - d_t(a(t) + 1) \\ &\geq 2 \sum_{q \in \mathcal{M}_k^+(t_1)} d_{t_1}(q) - 2 \sum_{x \in \mathcal{M}_k^-(t_1)} d_{t_1}(x) - d_{t_1}(a(t_1)) - d_{t_1}(a(t_1) + 1) \longrightarrow \text{Var}(d_{t_1}), \end{aligned}$$

hence $\text{Var}(d_t) \geq \text{Var}(d_{t_1})$. □

BIBLIOGRAPHY

- [A] V. I. Arnold : *Chapitres supplémentaires de la théorie des équations différentielles ordinaires* (1980); or *Geometrical methods in the theory of ordinary differential equations* ; English transl. from the Russian. - 2nd ed. - Grundlehren der mathematischen Wissenschaften **250**, Springer (1988).
- [Ba] G. Barles : *Solutions de viscosité des équations de Hamilton-Jacobi*, *Mathematiques & Application* **17**, Springer (1994).
- [Be] P. Bernard: Connecting orbits of time dependent Lagrangian systems, *Ann. Inst. Fourier* **52**, no. 5, 1533–1568, (2002).
- [BR] P. Bernard, J.M. Roquejoffre : Convergence to time-periodic solutions in Hamilton-Jacobi equations on the circle, *Comm. in P.D.E.* **29**, no. 3-4, 457–469, (2004).
- [B] U. Bessi : Aubry-Mather theory and Hamilton-Jacobi equations, *Comm. Math. Phys.* **235**, no. 3, 495–511, (2003).
- [Br] H. Brezis : *Analyse Fonctionnelle*, Masson, (1983).
- [E] W. E: Aubry-Mather theory and periodic solutions of the forced Burgers equation, *Comm. Pure Appl. Math.* **52**, no. 7, 811–828, (1999).
- [EG] L. C. Evans, R. E. Gariepy : *Measure theory and fine properties of functions*, Studies in advanced mathematics, CRC Press, (1992).
- [Fa1] A. Fathi : Théorème KAM faible et théorie de Mather sur les systèmes lagrangiens. (French) [A weak KAM theorem and Mather’s theory of Lagrangian systems] *C. R. Acad. Sci. Sér. I Math.* **324**, no. 9, 1043–1046, (1997).
- [Fa2] A. Fathi : Book in preparation.
- [Fa3] A. Fathi : Sur la convergence du semi-groupe de Lax-Oleinik. (French) [Convergence of the Lax-Oleinik semigroup] *C. R. Acad. Sci. Sér. I Math.* **327** (1998), no. 3, 267–270, (1998).
- [FM] A. Fathi, J. N. Mather : Failure of convergence of the Lax-Oleinik semi-group in the time-periodic case. *Bull. Soc. Math. France* **128**, no. 3, 473–483, (2000).
- [FMa] A. Fathi, E. Maderna : Weak KAM theorem on non compact manifolds, preprint (2002).
- [Fl] W. H. Fleming : The Cauchy problem for a nonlinear first order partial differential equation, *J. Differential Equations* **5**, 515–530, (1969).
- [G] D. Gomes: Viscosity Solutions of Hamilton-Jacobi Equations, and Asymptotics for Hamiltonian Systems - *Calc. Var. and P. D. E.* **14.3**, 345–357, (2002).
- [HF] M.R. Herman : *Sur les courbes invariantes par les difféomorphismes de l’anneau*, Vol. 1. (French) [On the curves invariant under diffeomorphisms of the annulus. Vol. 1] With an appendix by Albert Fathi. *Astérisque* **103-104**, (1983).
- [H] L. Hörmander : *Lectures on nonlinear hyperbolic differential equations*, *Mathematiques & Application* **26**, Springer (1997).

- [JKM] H.R. Jauslin, O. Kreiss, J. Moser : On the forced Burgers equation with periodic boundary conditions, *Proceedings of symposia in pure Mathematics* **65**, 133–153, (1999).
- [KO] Y. Katznelson, D.S. Ornstein : Twist maps and Aubry-Mather sets, *Contemporary Mathematics* **211**, 343–357, (1997).
- [L] P. Lax : Hyperbolic systems of conservation laws II, *Comm. Pure App. Maths* **10**, 537–566, (1957).
- [Ma1] J.N. Mather: Action minimizing invariant measures for positive definite Lagrangian systems, *Math. Z.* **207**, 169–207, (1991)
- [Ma2] J. N. Mather : Variational construction of orbits of twist diffeomorphisms, *J. Amer. Math. Soc.* **4**, no. 2, 207–263, (1991).
- [Mo] J. Moser : Monotone twist mappings and the calculus of variations, *Ergodic Theory Dynam. Systems* **6**, no. 3, 401–413, (1986).
- [Ro] J-M. Roquejoffre : Convergence to steady states or periodic solutions in a class of Hamilton-Jacobi equations, *J. Math. Pures Appl.* **80,1**, 85–104 (2001).
- [So] A. N. Sobolevskii : Periodic solutions of the Hamilton-Jacobi equation with a periodic nonhomogeneity, and the Aubry-Mather theory. (Russian) *Mat. Sb.* **190**, no. 10, 87–104 (1999); translation in *Sb. Math.* **190**, no. 9-10, 1487–1504, (1999).
- [Se] D. Serre : *Systèmes de lois de conservation* , vol 1, Paris, New York, Amsterdam : Diderot, (1996).
- [Si] K. F. Siburg : A dynamical approach to Birkhoff’s theorem, *Ens. Math.* **44**, 291–303 (1998).