

A unified approach to *a priori* estimates for supersolutions of BSDEs in general filtrations

Bruno BOUCHARD ^{*} Dylan POSSAMAÏ [†] Xiaolu TAN[‡] Chao ZHOU[§]

September 1, 2015

Abstract

We provide a unified approach to *a priori* estimates for supersolutions of BSDEs in general filtrations, which may not be quasi left-continuous. As an example of application, we prove that reflected BSDEs are well-posed in a general framework.

Key words: Backward SDE, supersolution, Doob-Meyer decomposition, reflected backward SDE.

MSC Classification (2010): 60H99

1 Introduction

Supersolutions of backward stochastic differential equations (BSDEs from now on) were introduced by El Karoui et al. in their seminal paper [7], in order to study superhedging strategies in mathematical finance. In the simple context of a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F} := (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ where \mathbb{F} is the (augmented) natural filtration of a d -dimensional Brownian motion W , a supersolution of a BSDE with terminal condition ξ and generator g consists is a triple of \mathbb{F} -adapted processes (Y, Z, K) , living in appropriate spaces, with K predictable non-decreasing, such that

$$Y_t = \xi - \int_t^T g_s(Y_s, Z_s) ds - \int_t^T Z_s \cdot dW_s + \int_t^T dK_s, \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.} \quad (1.1)$$

These objects appeared later to be at the very heart of the study of reflected BSDEs, as introduced in El Karoui et al. [8], and more generally of BSDEs satisfying some constraint, see Cvitanić, Karatzas and Soner [4] for constraints on the Z -component and Peng et al. [17, 21, 22, 23, 24] for general restrictions. More recently, supersolutions of BSDEs have proved to provide the semimartingale decomposition of the so-called second order BSDEs, introduced by Soner, Touzi and Zhang [26] and generalized by Possamaï, Tan and Zhou [25], and of the weak BSDEs studied by Bouchard, Élie and Réveillac in [1].

^{*}CEREMADE, Université Paris Dauphine, bouchard@ceremade.dauphine.fr.

[†]CEREMADE, Université Paris Dauphine, possamai@ceremade.dauphine.fr.

[‡]CEREMADE, Université Paris Dauphine, tan@ceremade.dauphine.fr, the author gratefully acknowledges the financial support of the ERC 321111 Rofirm, the ANR Isotace, and the Chairs Financial Risks (Risk Foundation, sponsored by Société Générale) and Finance and Sustainable Development (IEF sponsored by EDF and CA).

[§]Department of Mathematics, National University of Singapore, Singapore, matzc@nus.edu.sg. Research supported by NUS Grant R-146-000-179-133.

When the generator g is equal to 0, the process Y defined above is nothing else but a supermartingale, and (1.1) is simply its Doob-Meyer decomposition. This was generalized by Peng [21] using the notion of non-linear supermartingales, see also [2, 9] and the references therein.

As seen through the above examples, supersolutions of BSDEs appear quite frequently in the literature, as natural semimartingale decompositions for various stochastic processes, and are often used to study their fine properties. Having at hand *a priori* estimates on the moments and on the stability of supersolutions is crucial in these contexts. Unfortunately, in almost all the previously cited works, with the exception of [25], such estimates have been written in, roughly speaking, the context of a Brownian filtration. This is rather limiting from the point of view of both the theory and the applications, and it has created a tendency in the recent literature to reproduce very similar proofs every time that the context was generalized.

In this paper, we propose a general approach which allows one to consider a quite sufficiently general setting. In particular, we do not assume that the underlying filtration is generated by a Brownian motion. In this case, one needs to introduce another component in the definition of a supersolution of a BSDE, namely a martingale M that is orthogonal to W :

$$Y_t = \xi - \int_t^T g_s(Y_s, Z_s) ds - \int_t^T Z_s \cdot dW_s - \int_t^T dM_s + \int_t^T dK_s, \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.} \quad (1.2)$$

When $K \equiv 0$, such objects were first introduced by El Karoui and Huang [6], and studied more recently by Kruse and Popier [13] to handle more general filtrations, in the context of \mathbb{L}^p -solutions, as in the seminal papers [7, 3]. Supersolutions in general filtrations play a crucial role for the class of reflected BSDEs studied by Klimsiak [12], which is, as far as we know, the most general reference to date. However, all [6], [12]¹, [13] still impose that the filtration is quasi left-continuous, a property which, for instance, is not satisfied for the second order BSDEs studied in [25].

To understand the simplifications induced by the quasi left-continuity assumption, let us give a brief sketch of the strategy of proof usually used to obtain estimates, say in \mathbb{L}^2 for simplicity:

(i) Apply Itô's formula to $e^{\alpha \cdot} Y^2$ to obtain

$$\begin{aligned} e^{\alpha t} Y_t^2 + \alpha \int_t^T e^{\alpha s} Y_s^2 ds + \int_t^T e^{\alpha s} \|Z_s\|^2 ds + \int_t^T e^{\alpha s} d[M]_s \\ = e^{\alpha T} \xi^2 - 2 \int_t^T e^{\alpha s} Y_s g_s(Y_s, Z_s) ds - 2 \int_t^T e^{\alpha s} Y_s Z_s \cdot dW_s - 2 \int_t^T e^{\alpha s} Y_s dM_s + 2 \int_t^T e^{\alpha s} Y_s dK_s. \end{aligned}$$

(ii) Take expectations on both sides, use classical inequalities (namely Young and Burkholder-Davis-Gundy) and some continuity assumptions on g (usually Lipschitz continuity) to control \mathbb{L}^2 -type norms of (Y, Z, M) by the norm of K times a small constant, when $\alpha > 0$ is large enough.

(iii) Use the definition of a supersolution to control the norm of K by the norms of (Y, Z, M) , and conclude.

What is actually hidden in this reasoning is that, because of the quasi left-continuity assumption, the martingale M cannot jump at predictable times, and thus the bracket $[M, K]$ is identically equal to 0. This is no longer true for general filtrations, in which case we have to deal with the term $[M, K]$. It turns out to be difficult to control, which makes this traditional approach not amenable to filtrations that are not quasi left-continuous.

¹Notice that in [12], when the generator does not depend on Z , there is no need for the quasi left-continuity assumption. But the general case requires it.

The main aim of our paper is to give a general proof of *a priori* estimates and stability for supersolutions of BSDEs in a possibly non-quasi-left-continuous filtration. The proof relies on the following property:

"It is sufficient to control the norm of Y to control the norm of (Y, Z, M, K) ."

This is the philosophy of the estimates of Meyer [19, Theorem 1] that apply to general supermartingales (see also the generalization in [16, Thm 3.1]). In Section 2, we show how it can be generalized to the non-linear context of BSDEs. Namely, Theorem 2.1 below provides the extension of [19, Theorem 1] to a supersolution, while Theorem 2.2 is a version that applies to the difference of two supersolutions. Both are valid for supersolutions that are only làdlàg . In Section 3, we use these results to provide a well-posedness result for reflected BSDEs with a càdlàg obstacle. When there is no quasi left-continuity assumption on the filtration, this result is not available in the existing literature.

Notations. For any $l \in \mathbb{N} \setminus \{0\}$, we denote the usual inner product of two vectors $(x, y) \in \mathbb{R}^l \times \mathbb{R}^l$ by $x \cdot y$. The Euclidean norm on \mathbb{R}^l is denoted by $\|\cdot\|$, and simplified to $|\cdot|$ when $l = 1$. Let $T > 0$ be fixed and let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space, equipped with a filtration $\mathbb{F} = (\mathcal{F}_t)_{0 \leq t \leq T}$ satisfying the usual conditions, and carrying a standard d -dimensional \mathbb{F} -Brownian motion W . Importantly, we do not assume that the filtration is quasi left-continuous. Given $p > 1$ and $\alpha > 0$, we introduce the classical spaces:

- \mathbb{L}^p is the space of \mathbb{R} -valued and \mathcal{F}_T -measurable random variables ξ such that

$$\|\xi\|_{\mathbb{L}^p}^p := \mathbb{E} [|\xi|^p] < +\infty.$$

- \mathbb{S}^p (resp. \mathbb{S}_r^p) denotes the space of \mathbb{R} -valued, \mathbb{F} -adapted processes Y , with \mathbb{P} -a.s. làdlàg (resp. càdlàg) paths, such that

$$\|Y\|_{\mathbb{S}^p}^p := \mathbb{E} \left[\sup_{0 \leq s \leq T} |Y_s|^p \right] < +\infty.$$

- $\mathbb{M}^{p,\alpha}$ is the space of \mathbb{R} -valued, \mathbb{F} -adapted martingales M , with \mathbb{P} -a.s. càdlàg paths, such that M is orthogonal to W and

$$\|M\|_{\mathbb{M}^{p,\alpha}}^p := \mathbb{E} \left[\left(\int_0^T e^{\alpha s} d[M]_s \right)^{\frac{p}{2}} \right] < +\infty.$$

- $\mathbb{H}^{p,\alpha}$ (resp. $\mathbb{H}_1^{p,\alpha}$) is the space of \mathbb{R}^d -valued (resp. \mathbb{R} -valued) and \mathbb{F} -predictable processes Z such that

$$\|Z\|_{\mathbb{H}^{p,\alpha}}^p := \mathbb{E} \left[\left(\int_0^T e^{\alpha s} \|Z_s\|^2 ds \right)^{\frac{p}{2}} \right] < +\infty.$$

- $\mathbb{I}^{p,\alpha}$ (resp. $\mathbb{I}_+^{p,\alpha}$, $\mathbb{I}_r^{p,\alpha}$, $\mathbb{I}_{+,r}^{p,\alpha}$) denotes the space of \mathbb{R} -valued, \mathbb{F} -predictable processes with bounded variations K , with \mathbb{P} -a.s. làdlàg (resp. non-decreasing làdlàg , càdlàg , non-decreasing càdlàg) paths, such that

$$\|K\|_{\mathbb{I}^{p,\alpha}}^p := \mathbb{E} \left[\left(\int_0^T e^{\frac{\alpha}{2}s} d\text{TV}(K)_s \right)^p \right] < +\infty$$

and $K_0 = 0$. In the above $\text{TV}(K)$ denotes the total variation of K .

- We also define \mathbb{S}_{loc}^p as the collection of processes Y such that, for an increasing sequence of stopping times $(\tau_n)_{n \geq 1}$ satisfying $\mathbb{P}(\lim_{n \rightarrow \infty} \tau_n = \infty) = 1$, the localized process $Y_{\tau_n \wedge \cdot}$ belongs to \mathbb{S}^p for each $n \geq 1$. The spaces $\mathbb{M}_{loc}^{p,\alpha}$, $\mathbb{H}_{loc}^{p,\alpha}$, $\mathbb{H}_{1,loc}^{p,\alpha}$, $\mathbb{I}_{loc}^{p,\alpha}$, and $\mathbb{I}_{+,loc}^{p,\alpha}$ are defined similarly.
- Finally, for $\alpha = 0$, we simplify the notation $\mathbb{M}^p := \mathbb{M}^{p,0}$, $\mathbb{H}^p := \mathbb{H}^{p,0}$, $\mathbb{H}_1^p := \mathbb{H}_1^{p,0}$, $\mathbb{I}^p := \mathbb{I}^{p,0}$, $\mathbb{I}_+^p := \mathbb{I}_+^{p,0}$, $\mathbb{I}_r^p := \mathbb{I}_r^{p,0}$ and $\mathbb{I}_{+,r}^p := \mathbb{I}_{+,r}^{p,0}$.

Note that the above spaces do not depend on the precise value of α as we work on the compact time interval $[0, T]$, two values of α actually provide equivalent norms. Still, we keep the parameter α which, as usual, will be very helpful for many of our arguments.

Given a l  dl  g optional process X , such that its right-limit process X^+ is a semimartingale, and a locally bounded predictable process ϕ , we define the stochastic integral as in [15]:

$$(\phi \star X)_t := \int_0^t \phi_s dX_s := \int_0^t \phi_s dX_s^+ - \phi_t(X_{t+} - X_t), \quad t \geq 0.$$

Moreover, we define $\int_t^T \phi_s dX_s := \int_0^T \phi_s dX_s - \int_0^t \phi_s dX_s$.

2 A priori estimates

Let us consider a BSDE with terminal condition ξ and generator $g : [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^d \longrightarrow \mathbb{R}$. For ease of notations, we denote $g^0(t, \omega) := g(t, \omega, 0, 0)$. Although, we will have to differentiate between possible values of $p > 1$, this parameter is fixed from now on. The following standing assumption is assumed throughout this section.

Assumption 2.1. (i) $\xi \in \mathbb{I}^p$, $g^0 \in \mathbb{H}_1^p$ and the process $(t, \omega) \longmapsto g_t(\omega, y, z)$ is \mathbb{F} -progressively measurable for all $(y, z) \in \mathbb{R} \times \mathbb{R}^d$.

(ii) There are two constants $(L_y, L_z) \in \mathbb{R}_+^2$ such that

$$|g_t(\omega, y, z) - g_t(\omega, y', z')| \leq L_y |y - y'| + L_z \|z - z'\|, \quad (2.1)$$

for all $(t, \omega, y, z, y', z') \in [0, T] \times \Omega \times (\mathbb{R} \times \mathbb{R}^d)^2$.

We recall here the definition of a supersolution.

Definition 2.1. We say that (Y, Z, M, K) is a solution (resp. local solution) of

$$Y_t = \xi - \int_t^T g_s(Y_s, Z_s) ds - \int_t^T Z_s \cdot dW_s - \int_t^T dM_s + K_T - K_t, \quad (2.2)$$

if the above holds for any $t \in [0, T]$, \mathbb{P} -a.s., and $(Y, Z, M, K) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}^p$ (resp. $(Y, Z, M, K) \in \mathbb{S}_{loc}^p \times \mathbb{H}_{loc}^p \times \mathbb{M}_{loc}^p \times \mathbb{I}_{loc}^p$). If moreover $K \in \mathbb{I}_+^p$ (resp. $\mathbb{I}_{+,loc}^p$), we say that (Y, Z, M) is a supersolution (resp. a local supersolution) of (2.2).

2.1 Estimates for the solution

Our main result says that it is enough, in order to control the \mathbb{S}^p -norm of Y of a solution (Y, Z, M, K) , to control the norms of the other terms (Z, M, K) . We emphasize that the general setting we consider here creates additional difficulties that have not been tackled so far in the literature, and which mainly

stems from the fact that it is possible for the processes K and M to jump at the same time, when the filtration is not quasi left-continuous. Therefore, the traditional approach which consists in applying Itô's formula to $|Y|^p$ to derive the desired estimates fails, as this makes the cross-variation between M and K appear, a term that has no particular sign and cannot be controlled easily. Our message here is that, in order to obtain such estimates in a general setting, one should rely on a deeper result from the general theory of processes, namely the estimates obtained in Meyer [19] for general supermartingales, a version of which we recall in the Appendix below, see Lemma A.1.

The following is an extension to the non-linear context.

Theorem 2.1. *Let $(Y, Z, M, K) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}_+^p$ be a solution of (2.2). Then, for any $\alpha \geq 0$, there is a constant $C_{2.1}^\alpha$ such that*

$$\|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|M\|_{\mathbb{M}^{p,\alpha}}^p + \|K\|_{\mathbb{I}^{p,\alpha}}^p \leq C_{2.1}^\alpha \left(\|\xi\|_{\mathbb{L}^p}^p + \|Y\|_{\mathbb{S}^p}^p + \|g^0\|_{\mathbb{H}_1^p}^p \right).$$

Before proving this result, we shall establish more general intermediate estimates, that will also be used to control the difference of solutions in Theorem 2.2 below. They use the notation

$$N := Z \star W + M - K.$$

We start with an easy remark.

Remark 2.1. (i) *First note that for any $\ell > 0$ and $(a_i)_{1 \leq i \leq n} \subset (0, +\infty)$,*

$$(1 \wedge n^{\ell-1}) \sum_{i=1}^n a_i^\ell \leq \left(\sum_{i=1}^n a_i \right)^\ell \leq (1 \vee n^{\ell-1}) \sum_{i=1}^n a_i^\ell. \quad (2.3)$$

Let us now consider a solution $(Y, Z, M, K) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}^p$ of (2.2). Since W and M are orthogonal, (2.3) implies that

$$(1 \wedge 2^{\frac{p}{2}-1}) (\|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|M - K\|_{\mathbb{M}^{p,\alpha}}^p) \leq \|N\|_{\mathbb{M}^{p,\alpha}}^p \leq (1 \vee 2^{\frac{p}{2}-1}) (\|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|M - K\|_{\mathbb{M}^{p,\alpha}}^p). \quad (2.4)$$

Moreover, if $K \in \mathbb{I}_+^{p,\alpha}$ then the Kunita-Watanabe inequality leads to

$$d[M + Z \star W] \leq 2(d[N] + d[K]) \leq 2(d[N] + 2K_- dK + d[K]) \leq 2(d[N] + dK^2),$$

so that, by (2.3) and the fact that W and M are orthogonal,

$$(1 \wedge 2^{\frac{p}{2}-1}) (\|M\|_{\mathbb{M}^{p,\alpha}}^p + \|Z\|_{\mathbb{H}^{p,\alpha}}^p) \leq \|M + Z \star W\|_{\mathbb{M}^{p,\alpha}}^p \leq (2^{\frac{p}{2}} \vee 2^{p-1}) \left(\|N\|_{\mathbb{M}^{p,\alpha}}^p + e^{\frac{\alpha p T}{2}} \|K\|_{\mathbb{I}^{p,\alpha}}^p \right), \quad (2.5)$$

in which the left-hand side inequality remains true even if K is not non-decreasing.

(ii) *In the following, we shall also use the standard Young's inequality*

$$ab \leq \beta a^p + \frac{b^q}{q(\beta p)^{q/p}}, \text{ for } a, b \in \mathbb{R}_+, \beta > 0, p, q > 1 \text{ and } \frac{1}{p} + \frac{1}{q} = 1. \quad (2.6)$$

(iii) *We also emphasize that for $(Y, Z, M) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p$, the process*

$$\int_0^\cdot e^{p \frac{\alpha}{2} s} \phi_p(Y_{s-}) d(M + Z \star W),$$

is a uniformly integrable martingale, where

$$\phi_p(y) = |y|^{p-1} \text{sgn}(y) \mathbf{1}_{y \neq 0}, \quad \text{for } y \in \mathbb{R}. \quad (2.7)$$

Indeed, Burkholder-Davis-Gundy and Hölder's inequalities imply

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} \left| \int_0^t e^{p \frac{\alpha}{2} s} \phi_p(Y_{s-}) d(M + Z \star W)_s \right| \right] \\ & \leq C \mathbb{E} \left[\sqrt{\int_0^T e^{p \alpha s} |Y_{s-}|^{2p-2} d[M]_s + \int_0^T e^{p \alpha s} |Y_s|^{2p-2} \|Z_s\|^2 ds} \right] \\ & \leq C e^{(p-1) \frac{\alpha}{2} T} \|Y\|_{\mathbb{S}^p}^{p-1} (\|M\|_{\mathbb{M}^{p,\alpha}} + \|Z\|_{\mathbb{H}^{p,\alpha}}), \end{aligned}$$

for some $C > 0$.

From now on, we use the generic notation C , combined with super- and subscripts, to denote constants in our estimates that only depend on L_y, L_z, p and α . If they depend on other parameters, this will be made clear. Although we do not provide their expressions explicitly, our proofs are written in such a way that the interested reader can easily keep track of them line after line.

In the following, the inequality (2.8) is the crucial one, this is the consequence of Meyer [19].

Lemma 2.1. *Let $(Y, Z, M, K) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}^p$ be a solution of (2.2).*

(i) *If $K \in \mathbb{I}_{+}^p$, then for all $\alpha \geq 0$ there exists a constant $C_{(2.8)}^\alpha$ such that*

$$\|K\|_{\mathbb{I}^{p,\alpha}}^p \leq C_{(2.8)}^\alpha \left(\|e^{\frac{\alpha}{2} \cdot} Y\|_{\mathbb{S}^p}^p + \|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p \right). \quad (2.8)$$

(ii) *If $p \geq 2$, then for all $\varepsilon > 0$ there exists $\alpha > 0$ and a constant $C_{(2.9)}^{\varepsilon,\alpha}$ such that*

$$\|Y\|_{\mathbb{H}_1^{p,\alpha}}^p + \|N\|_{\mathbb{M}^{p,\alpha}}^p \leq \varepsilon \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p + C_{(2.9)}^{\varepsilon,\alpha} \left(\|\xi\|_{\mathbb{L}^p}^p + \|(e^{\alpha \cdot} Y_- \star N)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \mathbf{1}_{p>2} + \mathbb{E}[(e^{\alpha \cdot} Y_- \star K)_T]^+ \mathbf{1}_{p=2} \right). \quad (2.9)$$

(iii) *If $p \in (1, 2)$, then for all $\varepsilon > 0$ there exists $\alpha > 0$ and a constant $C_{(2.10)}^{\varepsilon,\alpha}$ such that*

$$\|N\|_{\mathbb{M}^{p,\alpha}}^p \leq \varepsilon \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p + C_{(2.10)}^{\varepsilon,\alpha} \left(\|\xi\|_{\mathbb{L}^p}^p + \|e^{\frac{\alpha}{2} \cdot} Y\|_{\mathbb{S}^p}^p + \mathbb{E}[(e^{p \frac{\alpha}{2} \cdot} \phi_p(Y_-) \star K)_T]^+ \right), \quad (2.10)$$

where ϕ_p is defined in (2.7).

Proof. (i) Let us first prove (2.8). A simple application of Itô's formula implies that

$$e^{\frac{\alpha}{2} \cdot} Y_- - \int_0^\cdot e^{\frac{\alpha}{2} s} \left(g_s(Y_s, Z_s) + \frac{\alpha}{2} Y_s \right) ds$$

is a supermartingale. Moreover, the non-decreasing process in its Doob-Meyer decomposition is $\int_0^\cdot e^{\frac{\alpha}{2} s} dK_s$. Therefore, Lemma A.1, Assumption 2.1 and Jensen's inequality provide

$$\begin{aligned} (C_{A.1}^p)^{-p} \|K\|_{\mathbb{I}^{p,\alpha}}^p & \leq \left\| e^{\frac{\alpha}{2} \cdot} Y - \int_0^\cdot e^{\frac{\alpha}{2} s} \left(g_s(Y_s, Z_s) + \frac{\alpha}{2} Y_s \right) ds \right\|_{\mathbb{S}^p}^p \\ & \leq (1 \vee 2^{p-1}) \left(\|e^{\frac{\alpha}{2} \cdot} Y\|_{\mathbb{S}^p}^p + \mathbb{E} \left[\left(\int_0^T e^{\frac{\alpha}{2} s} \left(|g_s(Y_s, Z_s)| + \frac{\alpha}{2} |Y_s| \right) ds \right)^p \right] \right) \\ & \leq (1 \vee 2^{p-1}) \left(\left(1 + (1 \vee 3^{p-1}) T^p \left(L_y + \frac{\alpha}{2} \right)^p \right) \|e^{\frac{\alpha}{2} \cdot} Y\|_{\mathbb{S}^p}^p + (1 \vee 3^{p-1}) \left(L_z^p \|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p \right) \right), \end{aligned} \quad (2.11)$$

in which the constant $C_{A.1}^p$ is as in Lemma A.1.

(ii) We now turn to (2.9). As usual, we apply Itô's formula to $e^{\alpha \cdot} Y^2$, see [15, p.538] for the case of càdlàg processes, use Assumption 2.1 and (2.6), to obtain

$$\begin{aligned} e^{\alpha t} Y_t^2 + \left(\alpha - \frac{1}{\varepsilon} - 2L_y - \frac{L_z^2}{\eta} \right) \int_t^T e^{\alpha s} Y_s^2 ds + (1 - \eta) \int_t^T e^{\alpha s} \|Z_s\|^2 ds + \int_t^T e^{\alpha s} d[M - K]_s \\ \leq e^{\alpha T} |\xi|^2 + \varepsilon \int_t^T e^{\alpha s} |g_s^0|^2 ds - 2 \int_t^T e^{\alpha s} Y_{s-} dN_s, \end{aligned} \quad (2.12)$$

for any $(\varepsilon, \eta) \in (0, +\infty)^2$. Combined with (2.3), this implies that

$$\begin{aligned} C_1 \|Y\|_{\mathbb{H}_1^{p,\alpha}}^p + C_2 \|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|M - K\|_{\mathbb{M}^{p,\alpha}}^p &\leq 3^{\frac{p}{2}-1} \left(e^{\frac{p}{2}\alpha T} \|\xi\|_{\mathbb{L}^p}^p + \varepsilon^{\frac{p}{2}} \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p \right) \\ &\quad + 3^{\frac{p}{2}-1} 2^{\frac{p}{2}} \|(e^{\alpha \cdot} Y_- \star N)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \mathbf{1}_{p>2} \\ &\quad + 3^{\frac{p}{2}-1} 2^{\frac{p}{2}} \mathbb{E}[(e^{\alpha \cdot} Y_- \star K)_T]^+ \mathbf{1}_{p=2}, \end{aligned}$$

where $C_1 := (\alpha - \frac{1}{\varepsilon} - 2L_y - \frac{L_z^2}{\eta})^{\frac{p}{2}}$, $C_2 := (1 - \eta)^{\frac{p}{2}}$ and where we have used Remark 2.1(iii) in the case $p = 2$. Fix $\alpha > 0$ and $\eta \in (0, 1)$ such that $C_1, C_2 > 0$. We then deduce (2.9) from the right-hand side of (2.4) for α large enough.

(iii) It remains to prove (2.10). Since $p < 2$, we can not use the Burkholder-Davis-Gundy inequality with exponent $p/2$ to a martingale involving M , as it is only càdlàg. We then follow the approach proposed recently in [13]. We first appeal to Lemma A.2 below:

$$\begin{aligned} e^{p\frac{\alpha}{2}t} |Y_t|^p + \frac{p(p-1)}{2} \int_t^T e^{p\frac{\alpha}{2}s} |\phi_{p-1}(Y_s)| d[N]_s^c + \frac{\alpha p}{2} \int_t^T e^{p\frac{\alpha}{2}s} |Y_s|^p ds + A_t^T \\ \leq e^{p\frac{\alpha}{2}T} |\xi|^p + p \int_t^T e^{p\frac{\alpha}{2}s} |Y_s|^{p-1} |g_s(Y_s, Z_s)| ds - p \int_t^T e^{p\frac{\alpha}{2}s} \phi_p(Y_{s-}) dN_s, \end{aligned}$$

in which

$$A_t^T := \frac{p(p-1)}{2} \sum_{t < s \leq T} e^{p\frac{\alpha}{2}s} |\Delta N_s|^2 \left(|Y_{s-}|^2 \vee |Y_{s-} + \Delta N_s|^2 \right)^{\frac{p}{2}-1} \mathbf{1}_{|Y_{s-}| \vee |Y_{s-} + \Delta N_s| \neq 0},$$

with $\Delta N := N_+ - N_-$. Recalling Assumption 2.1 and using (2.6), this shows that, for any $\beta, \gamma > 0$,

$$\begin{aligned} e^{p\frac{\alpha}{2}t} |Y_t|^p + \left(\frac{p(p-1)}{2} - \beta \right) \int_t^T e^{p\frac{\alpha}{2}s} |\phi_{p-1}(Y_s)| d[N]_s^c + p \left(\frac{\alpha}{2} - L_y - \frac{pL_z^2}{4\beta} \right) \int_t^T e^{p\frac{\alpha}{2}s} |Y_s|^p ds + A_t^T \\ \leq e^{p\frac{\alpha}{2}T} |\xi|^p + p \int_t^T e^{p\frac{\alpha}{2}s} |Y_s|^{p-1} |g_s^0| ds - p \int_t^T e^{p\frac{\alpha}{2}s} \phi_p(Y_{s-}) dN_s \\ \leq e^{p\frac{\alpha}{2}T} |\xi|^p + \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p + \frac{p-1}{p^{\frac{p}{p-1}}} \sup_{0 \leq s \leq T} \left| e^{\frac{\alpha}{2}s} Y_s \right|^p - p \int_t^T e^{p\frac{\alpha}{2}s} \phi_p(Y_{s-}) dN_s. \end{aligned}$$

Let us take $\alpha \geq 2L_y + pL_z^2/(2\beta)$ with $\beta < p(p-1)/2$. Taking expectations on both sides, we obtain

$$\begin{aligned} \mathbb{E} \left[\int_t^T e^{p\frac{\alpha}{2}s} |\phi_{p-1}(Y_s)| d[N]_s^c + A_t^T \right] &\leq C_{(2.13)}^1 \|\xi\|_{\mathbb{L}^p}^p + C_{(2.13)}^2 \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p + C_{(2.13)}^3 \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p \\ &\quad + C_{(2.13)}^4 \mathbb{E}[(e^{p\frac{\alpha}{2} \cdot} \phi_p(Y_-) \star K)_T]^+, \end{aligned} \quad (2.13)$$

for some explicit constants $(C_{(2.13)}^i)_{1 \leq i \leq 4}$. We then argue as in [13, Step 2, Proof of Proposition 3] and use (2.6) again to obtain that

$$\|N\|_{\mathbb{M}^{p,\alpha}}^p \leq \left(\frac{2C_{(2.13)}^2}{\varepsilon p} \right)^{\frac{1}{p-1}} (2-p) \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p + \frac{\varepsilon}{C_{(2.13)}^2} \mathbb{E} \left[\int_0^T e^{p\frac{\alpha}{2}s} |\phi_{p-1}(Y_s)| d[N]_s^c + A_0^T \right],$$

for $\varepsilon > 0$. □

We are now in position to complete the proof of Theorem 2.1.

Proof of Theorem 2.1. 1. We first assume that $p \geq 2$. In the course of this proof, we will have to choose $\alpha > 0$ large to apply (2.9). However, since the norms in $\{\|\cdot\|_{\mathbb{H}^{p,\alpha}}^p, \alpha > 0\}$ (resp. $\{\|\cdot\|_{\mathbb{H}_1^{p,\alpha}}^p, \alpha > 0\}$ and $\{\|\cdot\|_{\mathbb{M}^{p,\alpha}}^p, \alpha > 0\}$) are equivalent for different values of α , this is enough to prove our general result. We first estimate the last term in (2.9):

$$\|(e^\alpha Y_- \star N)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \leq 2^{\frac{p}{2}-1} \left(\|(e^\alpha Y_- \star (M + Z \star W))_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} + \|(e^\alpha Y_- \star K)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \right),$$

in which, for any $\delta > 0$,

$$\|(e^\alpha Y_- \star K)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \leq \frac{1}{4\delta} \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p + \delta \|K\|_{\mathbb{I}^{p,\alpha}}^p,$$

recall (2.6), and

$$\begin{aligned} \|(e^\alpha Y_- \star (M + Z \star W))_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} &\leq C_p^* \mathbb{E} \left[\left(\int_0^T e^{2\alpha s} Y_{s-}^2 d[M + Z \star W]_s \right)^{\frac{p}{4}} \right] \\ &\leq \frac{(C_p^*)^2}{4\delta} \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p + \delta \|M + Z \star W\|_{\mathbb{M}^{p,\alpha}}^p, \end{aligned}$$

with

$$C_p^* := \left(\frac{p}{2} \vee \frac{p}{p-2} - 1 \right)^p \mathbf{1}_{p>2} + 2\mathbf{1}_{p=2}, \quad (2.14)$$

by Burkholder's inequality, see e.g. [20, Theorems 8.6 and 8.7]. Combining the above inequalities with (2.5) leads to

$$\|(e^\alpha Y_- \star N)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \mathbf{1}_{p>2} + \|(e^\alpha Y_- \star K)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \mathbf{1}_{p=2} \leq \frac{C_{(2.15)}^1}{\delta} \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p + \delta C_{(2.15)}^2 (\|N\|_{\mathbb{M}^{p,\alpha}}^p + \|K\|_{\mathbb{I}^{p,\alpha}}^p), \quad (2.15)$$

for some explicit constants $C_{(2.15)}^1$ and $C_{(2.15)}^2$ that do not depend on δ . By inserting the last inequality in (2.9), for e.g. $\varepsilon = 1$ and α chosen appropriately, we obtain for $\delta \in (0, 1)$:

$$(1 - \delta C_{(2.16)}^4) \|N\|_{\mathbb{M}^{p,\alpha}}^p \leq C_{(2.16)}^1 \|\xi\|_{\mathbb{L}^p}^p + C_{(2.16)}^2 \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p + \frac{C_{(2.16)}^3}{\delta} \left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p + \delta C_{(2.16)}^4 \|K\|_{\mathbb{I}^{p,\alpha}}^p, \quad (2.16)$$

in which the constants $(C_{(2.16)}^i)_{i \leq 4}$ are explicit and do not depend on $\delta \in (0, 1)$. In view of the left-hand side of (2.4) and (2.8), (2.16) provides the required bound on $\|Z\|_{\mathbb{H}^{p,\alpha}}^p$ by choosing $\delta > 0$ small enough, so that we can then use (2.8) again to deduce the corresponding bound on $\|K\|_{\mathbb{I}^{p,\alpha}}^p$:

$$\|Z\|_{\mathbb{H}^{p,\alpha}}^p + \|K\|_{\mathbb{I}^{p,\alpha}}^p \leq C_{(2.17)} \left(\|\xi\|_{\mathbb{L}^p}^p + \|Y\|_{\mathbb{S}^p}^p + \|g^0\|_{\mathbb{H}_1^{p,\alpha}}^p \right), \quad (2.17)$$

for some constant $C_{(2.17)}$. Finally, it remains to appeal to (2.5), (2.16) and (2.17) to obtain the required bound on $\|M\|_{\mathbb{M}^{p,\alpha}}^p$ and conclude the proof in the case $p \geq 2$.

2. We now consider the case $p \in (1, 2)$. We argue as above except that we now estimate the last term in (2.10) by using (2.6):

$$\mathbb{E}[(e^{p\frac{\alpha}{2}} \phi_p(Y_-) \star K)_T] \leq \frac{p-1}{(\delta p^p)^{\frac{1}{p-1}}} \|e^{\frac{\alpha}{2}} Y\|_{\mathbb{S}^p}^p + \delta \|K\|_{\mathbb{I}^{p,\alpha}}^p. \quad (2.18)$$

The latter combined with (2.4), (2.5), (2.8) and (2.10), as in the end of step 1, provides the required result after choosing $\delta > 0$ small enough. \square

When (Y, Z, M, K) is only a local solution of (2.2), all the arguments above hold true after a localization. Then, using Fatou's Lemma, it follows immediately that (Y, Z, M, K) is a solution. We formulate the following result but omit the proof.

Proposition 2.1. *Let (Y, Z, M, K) be a local solution of (2.2). Suppose in addition that $Y \in \mathbb{S}^p$. Then, (Y, Z, M, K) is a solution of (2.2).*

2.2 Difference of solutions and stability

In this section, we consider two terminal conditions ξ^1, ξ^2 , as well as two generators g^1 and g^2 , satisfying Assumption 2.1. We then denote by $(Y^i, Z^i, M^i, K^i) \in \mathbb{S}^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}_+^p$ a solution of (2.2) with terminal condition ξ^i and generator g^i , and set $N^i := Z^i \star W + M^i - K^i$, $i = 1, 2$. For notational simplicity, we also define

$$\begin{aligned} \delta Y &:= Y^1 - Y^2, \quad \delta Z := Z^1 - Z^2, \quad \delta M := M^1 - M^2, \quad \delta K := K^1 - K^2, \quad \delta N := N^1 - N^2, \\ \delta g(t, \omega, y, z) &:= g^1(t, \omega, y, z) - g^2(t, \omega, y, z), \quad \text{for all } (t, \omega, y, z) \in [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^d. \end{aligned}$$

By Assumption 2.1, we know that there is an \mathbb{R} -valued (resp. \mathbb{R}^d -valued), \mathbb{F} -progressively measurable process λ (resp. η), with $|\lambda| \leq L_y$ (resp. $\|\eta\| \leq L_z$) such that

$$\delta g_t := g^1(t, Y_t^1, Z_t^1) - g^2(t, Y_t^2, Z_t^2) = \delta g(t, Y_t^1, Z_t^1) + \lambda_t \delta Y_t + \eta_t \cdot \delta Z_t.$$

Then, $(\delta Y, \delta Z, \delta M, \delta K)$ satisfies (2.2) with driver δg and terminal condition $\delta \xi$. In particular, we can apply to it the results of Remark 2.1 and Lemma 2.1.

The main result of this section, Theorem 2.2 below, is in the spirit of Theorem 2.1: it suffices to control the norm of δY to control the norms of δZ and $\delta(M - K)$. Seemingly, it should just be an application of Theorem 2.1 to $(\delta Y, \delta Z, \delta M, \delta K)$ as it satisfies an equation of the form (2.2). However, it is not the case:

- (i) In Theorem 2.2, we will only control $\delta(M - K)$ and not δM and δK separately. Actually, as shown in Example A.1 below, there is no hope to control these two processes separately even in the seemingly benign case where $g^1 = g^2 = 0$.
- (ii) Actually, Theorem 2.2 can not be an immediate consequence of Theorem 2.1, because the process δK which appears in the dynamics of δY is no longer non-decreasing, and more importantly because the result of Lemma A.1 below does not hold for quasimartingales (instead of supermartingales). However, it is a direct consequence of the intermediate estimates of Lemma 2.1, which explains why they have been isolated.

(iii) If one has a more precise knowledge of the behavior of the non-decreasing processes K^1 and K^2 , then these estimates can actually be improved. We will make this point more clear when we will treat the special case of reflected BSDEs in Section 3.

Let us now state our result.

Theorem 2.2. *For any $\alpha \geq 0$, there is a constant $C_{2,2}^\alpha$ such that*

$$\|\delta Z\|_{\mathbb{H}^{p,\alpha}}^p + \|\delta(M - K)\|_{\mathbb{M}^{p,\alpha}}^p \leq C_{2,2}^\alpha \left(\|\delta\xi\|_{\mathbb{L}^p}^p + \|\delta Y\|_{\mathbb{S}^p}^p + \|\delta Y\|_{\mathbb{S}^p}^{\frac{p}{2} \wedge (p-1)} + \|\delta g(Y^1, Z^1)\|_{\mathbb{H}_1^{p,\alpha}}^p \right).$$

The constant $C_{2,2}^\alpha$ depends on L_y, L_z, p and α , as well as $(\|Y^i\|_{\mathbb{S}^p}^p, \|\xi^i\|_{\mathbb{L}^p}^p, \|g^i(0, 0)\|_{\mathbb{H}_1^{p,\alpha}}^p)_{i=1,2}$.

Proof. In this proof, we take α large enough so as to apply the estimates of Lemma 2.1. The general case is deduced by recalling that the different norms are equivalent for different values of α , since $[0, T]$ is compact.

1. We first assume that $p \geq 2$. We shall apply (2.9) to $(\delta Y, \delta Z, \delta M)$. Let us estimate the last term in this inequality. We first use (2.3) to obtain

$$\|(e^{\alpha \cdot} \delta Y_- \star \delta N)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \leq 2^{\frac{p}{2}-1} \left(\|(e^{\alpha \cdot} \delta Y_- \star (\delta M + \delta Z \star W))_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} + \|(e^{\alpha \cdot} \delta Y_- \star \delta K)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \right).$$

We then apply Burkholder inequality and use (2.3) again:

$$\|(e^{\alpha \cdot} \delta Y_- \star (\delta M + \delta Z \star W))_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \leq C_p^* \left\| e^{\frac{\alpha}{2} \cdot} \delta Y \right\|_{\mathbb{S}^p}^{\frac{p}{2}} \|\delta M + \delta Z \star W\|_{\mathbb{H}^{p,\alpha}}^{\frac{p}{2}}$$

where C_p^* is as in (2.14), while

$$\|(e^{\alpha \cdot} \delta Y_- \star \delta K)_T\|_{\mathbb{L}^{\frac{p}{2}}}^{\frac{p}{2}} \leq \left\| e^{\frac{\alpha}{2} \cdot} \delta Y \right\|_{\mathbb{S}^p}^{\frac{p}{2}} \|\delta K\|_{\mathbb{I}^{p,\alpha}}^{\frac{p}{2}}.$$

We can then conclude the proof in the case $p \geq 2$ by using (2.9), (2.4) and the bounds of Theorem 2.1 applied to $(Z^i, M^i, K^i)_{i=1,2}$.

2. We now assume that $p \in (1, 2)$ and proceed as above but use (2.10) in place of (2.9). Namely, since

$$\mathbb{E} \left[(e^{p \frac{\alpha}{2} \cdot} \phi_p(\delta Y_-) \star \delta K)_T \right] \leq \left\| e^{\frac{\alpha}{2} \cdot} \delta Y \right\|_{\mathbb{S}^p}^{p-1} \|\delta K\|_{\mathbb{I}^{p,\alpha}},$$

it suffices to use (2.10), (2.4) and the bound of Theorem 2.1 applied to K^1 and K^2 . \square

3 Application to reflected BSDEs: a general existence result

The results of the previous section show that it suffices to control the norm of Y (resp. δY) in order to control the norm of (Z, M, K) (resp. $(\delta Z, \delta M, \delta K)$), given a solution (Y, Z, M, K) (resp. two solutions (Y^1, Z^1, M^1, K^1) and (Y^2, Z^2, M^2, K^2)) of (2.2). In most examples of applications, we know how to control the norm of Y (and δY). This is in particular the case in the context of reflected BSDEs (see e.g. [8]), BSDEs with constraints (see e.g. [4, 23]), 2nd order BSDE (see e.g. [26, 25]), weak BSDEs (see [1]).

Let us exemplify this in the context of reflected BSDEs. In particular, the following results extend Klimsiak [10, 11, 12] to a filtration that only satisfies the usual conditions, and may not be quasi left-continuous. For sake of simplicity, we restrict to the case of a càdlàg obstacle, see [9] and the references therein for the additional specific arguments that could be used for irregular obstacles. Recall that \mathbb{S}_r^p (resp. $\mathbb{I}_{+,r}^p$) denote the set of elements of \mathbb{S}^p (resp. \mathbb{I}^p , \mathbb{I}_+^p) with càdlàg path, \mathbb{P} -a.s.

3.1 A priori estimates for reflected BSDEs

In this section, we assume that Assumption 2.1 holds for ξ and g .

Definition 3.1. Let S be a càdlàg process such that $S^+ := S \vee 0 \in \mathbb{S}_r^p$. We say that $(Y, Z, M, K) \in \mathbb{S}_r^p \times \mathbb{H}^p \times \mathbb{M}^p \times \mathbb{I}_{+,r}^p$ is a solution of the reflected BSDE with lower obstacle S if

$$Y_t = \xi - \int_t^T g_s(Y_s, Z_s) ds - \int_t^T Z_s \cdot dW_s - \int_t^T dM_s + K_T - K_t, \quad (3.1)$$

holds for any $t \in [0, T]$ \mathbb{P} -a.s., and if

$$\begin{cases} Y_t \geq S_t, \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.}, \\ \int_0^T (Y_{s-} - S_{s-}) dK_s = 0, \quad \mathbb{P} - \text{a.s.} \quad (\text{Skorokhod condition}) \end{cases}$$

In order to provide a first estimate on the component Y of a solution, we use the classical linearization procedure. By Assumption 2.1, there exists a \mathbb{R} -valued (resp. \mathbb{R}^d -valued), \mathbb{F} -progressively measurable (resp. \mathbb{F} -predictable) process λ (resp. η), with $|\lambda| \leq L_y$ (resp. $\|\eta\| \leq L_z$) such that

$$g_s(Y_s, Z_s) = g_s^0 + \lambda_s Y_s + \eta_s \cdot Z_s, \quad s \in [0, T].$$

Let us define

$$X_s := e^{-\int_0^s \lambda_s ds}, \quad \text{and} \quad \frac{d\mathbb{Q}}{d\mathbb{P}} := \mathcal{E} \left(- \int_0^\cdot \eta_s \cdot dW_s \right)_T, \quad W^\mathbb{Q} := W + \int_0^\cdot \eta_s ds, \quad (3.2)$$

in which \mathcal{E} denotes the Doléans-Dade exponential. Then, by Girsanov theorem, $W^\mathbb{Q}$ is a \mathbb{Q} -Brownian motion, M is still a \mathbb{Q} -martingale orthogonal to $W^\mathbb{Q}$, and we can re-write the solution of the reflected BSDE (3.1) as

$$\begin{cases} X_t Y_t = X_T \xi - \int_t^T X_s g_s^0 ds - \int_t^T X_s Z_s \cdot dW_s^\mathbb{Q} - \int_t^T X_{s-} dM_s + \int_t^T X_{s-} dK_s, \quad t \in [0, T], \\ X_t Y_t \geq X_t S_t, \quad t \in [0, T], \\ \int_0^T X_{s-} (Y_{s-} - S_{s-}) dK_s = 0. \end{cases}$$

One can now use the link between reflected BSDEs and optimal stopping problems. The proof is classical so that we omit it, see [8] for a proof in a Brownian filtration and for a continuous obstacle, or [14] for a càdlàg obstacle, and [5] for more results on optimal stopping. We denote by $\mathcal{T}_{t,T}$ the set of $[t, T]$ -valued \mathbb{F} -stopping times, while \mathbb{E}_t and $\mathbb{E}_t^\mathbb{Q}$ stands for the \mathcal{F}_t -conditional expectations under \mathbb{P} and \mathbb{Q} .

Proposition 3.1. Let (Y, Z, M, K) and S be as in Definition 3.1. Then,

$$X_t Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_{t,T}} \mathbb{E}_t^\mathbb{Q} \left[- \int_t^\tau X_s g_s^0 ds + X_\tau S_\tau \mathbf{1}_{\tau < T} + X_T \xi \mathbf{1}_{\tau = T} \right],$$

and

$$Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_{t,T}} \mathbb{E}_t \left[- \int_t^\tau g_s(Y_s, Z_s) ds + S_\tau \mathbf{1}_{\tau < T} + \xi \mathbf{1}_{\tau = T} \right],$$

for all $t \leq T$.

Before continuing, let us introduce the solution $(\mathcal{Y}, \mathcal{Z}, \mathcal{M}) \in \mathbb{S}_r^p \times \mathbb{H}^p \times \mathbb{M}^p$ of the following BSDE (well-posedness is a direct consequence of Theorem 4.1 below, see also [13]),

$$\mathcal{Y}_t = \xi - \int_t^T g_s(\mathcal{Y}_s, \mathcal{Z}_s) ds - \int_t^T \mathcal{Z}_s \cdot dW_s - \int_t^T d\mathcal{M}_s, \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.}$$

A simple application of the comparison result, which can be proved as in [13, Proposition 4], implies that

$$Y_t \geq \mathcal{Y}_t, \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.}$$

Let us first show that Proposition 3.1 is actually enough to control the Y -term of a solution (or the δY -term of the difference of two solutions) and therefore that Theorems 2.1 and 2.2 apply to reflected BSDEs.

Proposition 3.2. (i) Let (Y, Z, M, K) and S be as in Definition 3.1. Then, for any $\alpha \geq 0$,

$$\left\| e^{\frac{\alpha}{2} \cdot} Y \right\|_{\mathbb{S}^p}^p \leq C_{3.2}^\alpha \left(\|\xi\|_{\mathbb{L}^p}^p + \|e^{L_y \cdot} S^+\|_{\mathbb{S}^p}^p + \mathbb{E} \left[\left(\int_0^T e^{L_y s} |g_s^0| ds \right)^p \right] \right) + C_{\mathcal{Y}}^\alpha \left\| e^{\frac{\alpha}{2} \cdot} \mathcal{Y} \right\|_{\mathbb{S}^p}^p,$$

for some constants $C_{3.2}^\alpha$ and $C_{\mathcal{Y}}^\alpha$ that only depend on L_y, L_z and α . Moreover, if we replace S^+ by S in the above, we can take $C_{\mathcal{Y}}^\alpha = 0$.

(ii) For $i = 1, 2$, let (Y^i, Z^i, M^i, K^i) and S^i be as in Definition 3.1 for a generator g^i satisfying Assumption 2.1 and terminal condition $\xi^i \in \mathbb{L}^p$. Then, for any $\alpha \geq 0$,

$$\left\| e^{\frac{\alpha}{2} \cdot} \delta Y \right\|_{\mathbb{S}^p}^p \leq \bar{C}_{3.2}^\alpha \left(\|\delta \xi\|_{\mathbb{L}^p}^p + \|e^{L_y \cdot} \delta S\|_{\mathbb{S}^p}^p + \mathbb{E} \left[\left(\int_0^T e^{L_y s} |\delta g_s(Y_s^1, Z_s^1)| ds \right)^p \right] \right),$$

for some constant $\bar{C}_{3.2}^\alpha$ that only depends on L_y, L_z and α .²

Proof. (i) First of all, we recall that $Y \geq \mathcal{Y}$ on $[0, T]$. Next, we deduce from Proposition 3.1 that, for any $1 < \kappa < p$,

$$\begin{aligned} & \sup_{t \in [0, T]} \left\{ e^{\frac{\alpha}{2} t} |Y_t| \right\} \\ & \leq \sup_{t \in [0, T]} e^{(L_y + \frac{\alpha}{2})t} \mathbb{E}_t^{\mathbb{Q}} \left[\int_0^T e^{L_y s} |g_s^0| ds + \sup_{s \in [0, T]} \{e^{L_y s} S_s^+\} + e^{L_y T} |\xi| \right] + \sup_{t \in [0, T]} \left\{ e^{\frac{\alpha}{2} t} |\mathcal{Y}_t| \right\} \\ & = \sup_{t \in [0, T]} e^{(L_y + \frac{\alpha}{2})t} \mathbb{E}_t \left[\mathcal{E} \left(- \int_t^T \eta_s \cdot dW_s \right) \left(\int_0^T e^{L_y s} |g_s^0| ds + \sup_{s \in [0, T]} \{e^{L_y s} S_s^+\} + e^{L_y T} |\xi| \right) \right] \\ & \quad + \sup_{t \in [0, T]} \left\{ e^{\frac{\alpha}{2} t} |\mathcal{Y}_t| \right\} \\ & \leq e^{(L_y + \frac{\alpha}{2})T + \frac{\kappa}{2(\kappa-1)} L_z^2 T} 3^{\frac{\kappa-1}{\kappa}} \sup_{t \in [0, T]} \mathbb{E}_t \left[\left(\int_0^T e^{L_y s} |g_s^0| ds \right)^\kappa + \sup_{s \in [0, T]} \{e^{\kappa L_y s} (S_s^+)^\kappa\} + e^{\kappa L_y T} |\xi|^\kappa \right]^{\frac{1}{\kappa}} \\ & \quad + \sup_{t \in [0, T]} \left\{ e^{\frac{\alpha}{2} t} |\mathcal{Y}_t| \right\}, \end{aligned}$$

in which it is clear that we could have suppressed the term involving \mathcal{Y} if we had used S instead of S^+ .

²We are grateful to Marie-Claire Quenez for indicating us a technical problem in the proof in the first version.

Hence, we deduce from Doob's inequality that

$$\begin{aligned} \left\| e^{\frac{\alpha}{2} \cdot Y} \right\|_{\mathbb{S}^p}^p &\leq e^{p(L_y + \frac{\alpha}{2})T + p\frac{\kappa}{2(\kappa-1)}L_z^2 T} 6^{p-1} \left(\frac{p}{p-1} \right)^p \mathbb{E} \left[\left(\int_0^T e^{L_y s} |g_s^0| ds \right)^p \right] \\ &\quad + e^{p(L_y + \frac{\alpha}{2})T + p\frac{\kappa}{2(\kappa-1)}L_z^2 T} 6^{p-1} \left(\frac{p}{p-1} \right)^p \left(\|e^{L_y \cdot} S^+\|_{\mathbb{S}^p}^p + e^{pL_y T} \|\xi\|_{\mathbb{L}^p}^p \right) \\ &\quad + 2^{p-1} \left\| e^{\frac{\alpha}{2} \cdot \mathcal{Y}} \right\|_{\mathbb{S}^p}^p, \end{aligned}$$

where we have to replace the 6^{p-1} by 3^{p-1} if we use S instead of S^+ .

(ii) We first use a classical argument. We know that there exists an \mathbb{R} -valued (resp. \mathbb{R}^d -valued), \mathbb{F} -progressively measurable process $\tilde{\lambda}$ (resp. $\tilde{\eta}$), with $|\tilde{\lambda}| \leq L_y$ (resp. $\|\tilde{\eta}\| \leq L_z$) such that

$$g_s^1(Y_s^1, Z_s^1) - g_s^2(Y_s^2, Z_s^2) = \delta g_s(Y_s^1, Z_s^1) + \tilde{\lambda}_s \delta Y_s + \tilde{\eta}_s \cdot \delta Z_s, \quad s \in [0, T], \quad \mathbb{P} - a.s.$$

Therefore, we can define $\tilde{\mathbb{Q}} \sim \mathbb{P}$ and a bounded positive process \tilde{X} as in (3.2) above such that

$$\tilde{X}_t \delta Y_t = \mathbb{E}_t^{\tilde{\mathbb{Q}}} \left[\int_t^\tau \tilde{X}_s \delta g_s(Y_s^1, Z_s^1) ds + \tilde{X}_\tau (Y_\tau^1 - Y_\tau^2) + \int_t^\tau \tilde{X}_s d(K^1 - K^2)_s \right]$$

for all stopping time $\tau \geq t$. Set $\tau_\varepsilon := \inf\{s \geq t : Y_s^1 \leq S_s^1 + \varepsilon\} \wedge T$. Clearly, $Y_-^1 \geq S_-^1 + \varepsilon$ on $[t, \tau_\varepsilon]$. Hence, $K_{\tau_\varepsilon}^1 - K_t^1 = 0$ by the Skorokhod condition. Moreover, $Y_{\tau_\varepsilon}^1 \leq S_{\tau_\varepsilon}^1 + \varepsilon$ on $\{\tau_\varepsilon < T\}$. Then, the above leads to

$$\begin{aligned} \tilde{X}_t \delta Y_t &\leq \mathbb{E}_t^{\tilde{\mathbb{Q}}} \left[\int_t^{\tau_\varepsilon} \tilde{X}_s |\delta g_s(Y_s^1, Z_s^1)| ds + \tilde{X}_{\tau_\varepsilon} (Y_{\tau_\varepsilon}^1 - Y_{\tau_\varepsilon}^2) + 0 - \int_t^{\tau_\varepsilon} \tilde{X}_s dK_s^2 \right] \\ &\leq \text{ess sup}_{\tau \in \mathcal{T}_{t,T}} \mathbb{E}_t^{\tilde{\mathbb{Q}}} \left[\int_t^\tau \tilde{X}_s |\delta g_s(Y_s^1, Z_s^1)| ds + \tilde{X}_\tau |\delta S_\tau| \mathbf{1}_{\tau < T} + \tilde{X}_T |\delta \xi| \mathbf{1}_{\tau = T} + \varepsilon \tilde{X}_\tau \right]. \end{aligned}$$

Since the same applies to $Y^2 - Y^1$ in place of $\delta Y = Y^1 - Y^2$, it follows that

$$|\tilde{X}_t \delta Y_t| \leq \text{ess sup}_{\tau \in \mathcal{T}_{t,T}} \mathbb{E}_t^{\tilde{\mathbb{Q}}} \left[\int_t^\tau \tilde{X}_s |\delta g_s(Y_s^1, Z_s^1)| ds + \tilde{X}_\tau |\delta S_\tau| \mathbf{1}_{\tau < T} + \tilde{X}_T |\delta \xi| \mathbf{1}_{\tau = T} + \varepsilon \tilde{X}_\tau \right].$$

It remains to let ε go to 0 and then argue exactly as in (i). \square

We now show that one can actually take advantage of the Skorokhod condition satisfied by the solution of a reflected BSDE to improve the general stability result of Theorem 2.2. This result is crucial in order to prove existence of a solution when p is arbitrary, see the proof of Theorem 3.1 below. We only provide the result for $p = 2$. It could be extended to $p \in (1, 2)$, but this is not important as we can always reduce to $p = 2$ by localization, again see the proof of Theorem 3.1 below.

Proposition 3.3. *For $i = 1, 2$, let (Y^i, Z^i, M^i, K^i) be as in (ii) of Proposition 3.2. Then, for any $\varepsilon > 0$, there exists α large enough such that*

$$\|\delta Y\|_{\mathbb{H}_1^{2,\alpha}}^2 + \|\delta Z\|_{\mathbb{H}^{2,\alpha}}^2 + \|\delta(M - K)\|_{\mathbb{M}^{2,\alpha}}^2 \leq \varepsilon \|\delta g(Y^1, Z^1)\|_{\mathbb{H}_1^{2,\alpha}}^2 + C_{3.3}^\alpha \left(\|\delta \xi\|_{L^2}^2 + \left\| e^{\frac{\alpha}{2} \cdot} \delta S \right\|_{\mathbb{S}^2} \right),$$

for some constant $C_{3.3}^\alpha$ that only depends on $L_y, L_g, \alpha, \varepsilon$ and $(\|Y^i\|_{\mathbb{S}^2}^2, \|\xi^i\|_{\mathbb{L}^2}^2, \|g^i(0, 0)\|_{\mathbb{H}_1^{2,\alpha}}^2)_{i=1,2}$.

Proof. It suffices to apply (2.4), (2.9) to $(\delta Y, \delta Z, \delta M, \delta K)$ and use the Skorokhod condition to deduce the control

$$\mathbb{E} \left[\int_t^T e^{\alpha s} \delta Y_{s-} d(\delta K_s) \right] \leq \mathbb{E} \left[\int_t^T e^{\alpha s} \delta S_{s-} d(\delta K_s) \right] \leq \left\| e^{\frac{\alpha}{2} \cdot} \delta S \right\|_{\mathbb{S}^2} \|\delta K\|_{\mathbb{I}^{2,\alpha}},$$

in which $\|\delta K\|_{\mathbb{I}^{2,\alpha}}$ is bounded by Theorem 2.1 and Proposition 3.2. \square

3.2 Wellposedness of reflected BSDE under general filtration

We can finally prove the existence of a unique solution to the reflected BSDEs (3.1). Our proof is extremely close to the original one given in [8], but relies on the more general estimates given in this paper.

Theorem 3.1. *Let Assumption 2.1 hold true. Then, there is a unique solution (Y, Z, M, K) to the reflected BSDE (3.1).*

Proof. The uniqueness is an immediate consequence of Proposition 3.3, recall that the Doob-Meyer decomposition of the supermartingale $M - K$ is unique. We therefore concentrate on the problem of existence.

1. First, let us consider the case $p = 2$ and prove the existence in 3 steps.

(i) We assume first that the function g does not depend on (y, z) , that is, $g(t, \omega, y, z) = g(t, \omega)$. Existence of a solution is guaranteed by [12, Theorem 2.12]. Moreover, [12, Corollary 2.8] implies

$$Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau g_s ds + S_\tau \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}} \middle| \mathcal{F}_t \right], \quad t \in [0, T], \quad \mathbb{P} - \text{a.s.}$$

Hence, $Y \in \mathbb{S}^2$ since

$$|Y_t| \leq \mathbb{E} \left[|\xi| + \int_0^T |g_s| ds + \sup_{0 \leq s \leq T} |S_s| \middle| \mathcal{F}_t \right],$$

so that the Burkholder-Davis-Gundy inequality can be used to deduce that

$$\|Y\|_{\mathbb{S}^2}^2 \leq C \left(\|\xi\|_{\mathbb{L}^2}^2 + \|g\|_{\mathbb{H}_1^2}^2 + \|S\|_{\mathbb{S}^2}^2 \right),$$

for some $C > 0$. Then, the fact that (Z, M, K) lies in the good space follows by Theorem 2.1.

(ii) We now consider the general case. Given $\alpha > 0$, let us consider the space \mathcal{S} of processes (Y, Z, L) such that

- $(Y, Z) \in \mathbb{S}_r^2 \times \mathbb{H}^{2, \alpha}$,
- L is a càdlàg supermartingale such that $L =: M - K$ where M is orthogonal to W ,
- (Y, Z, K, L) satisfies

$$Y_t \geq S_t, \quad t \in [0, T] \quad \text{and} \quad \int_0^T (Y_{t-} - S_{t-}) dK_t = 0, \quad \mathbb{P} - \text{a.s.}, \quad (3.3)$$

and

$$\|(Y, Z, L)\|_{2, \alpha} := \|e^\alpha Y\|_{\mathbb{S}^2} + \|Z\|_{\mathbb{H}^{2, \alpha}} + \|L\|_{\mathbb{S}^2} < \infty.$$

We will show in Step 3 below that $(\mathcal{S}, \|\cdot\|_{2, \alpha})$ is a Banach space. Then, existence can be proved by using the classical fixed point argument. Let us define $(\bar{Y}^0, \bar{Z}^0, \bar{L}^0) := (0, 0, 0)$ and $(\bar{Y}^n, \bar{Z}^n, \bar{L}^n)_{n \geq 1}$ by

$$\bar{Y}_t^{n+1} = \xi - \int_t^T g_s(\bar{Y}_s^n, \bar{Z}_s^n) ds - \int_t^T \bar{Z}_s^{n+1} \cdot dW_s - \int_t^T d\bar{L}_s^{n+1}. \quad (3.4)$$

Using the estimates of Proposition 3.3, it follows that, for $\alpha > 0$ large enough, $(\bar{Y}^n, \bar{Z}^n)_{n \geq 1}$ is a Cauchy sequence in $\mathbb{H}_1^{2,\alpha} \times \mathbb{H}^{2,\alpha}$, and hence a Cauchy sequence in $\mathbb{S}_r^2 \times \mathbb{H}^{2,\alpha}$ by the estimates in Part (ii) of Proposition 3.2. Moreover, by (3.4), we have

$$\begin{aligned} \bar{L}_t^n - \bar{L}_t^m &= (\bar{Y}_t^n - \bar{Y}_t^m) - (\bar{Y}_0^n - \bar{Y}_0^m) - \int_0^t (\bar{Z}_s^n - \bar{Z}_s^m) \cdot dW_s \\ &\quad - \int_0^t (g_s(\bar{Y}_s^{n-1}, \bar{Z}_s^{n-1}) - g_s(\bar{Y}_s^{m-1}, \bar{Z}_s^{m-1})) ds. \end{aligned} \quad (3.5)$$

It follows that $(\bar{Y}^n, \bar{Z}^n, \bar{L}^n)_{n \geq 1}$ is a Cauchy sequence in $(\mathcal{S}, \|\cdot\|_{2,\alpha})$, from which existence follows.

(iii) \mathcal{S} is clearly a subspace of a Banach space. It is therefore enough to show that it is closed, in order to prove that \mathcal{S} is a Banach space. The only non trivial point to check is that the Skorokhod condition in (3.3) passes to the limit. Let $(Y^n, Z^n, L^n)_{n \geq 1}$ be a sequence in \mathcal{S} that converges to (Y, Z, L) for the norm $\|\cdot\|_{2,\alpha}$ defined above. Clearly, L is still a càdlàg supermartingale in \mathbb{S}^2 , by dominated convergence. Let $L = M - K$ be its Doob-Meyer decomposition. Notice that the brackets $[L^n, W]$ also converge a.s. to $[L, W]$ (at least along a subsequence) thanks to Burkholder-Davis-Gundy inequalities. It follows that the martingale part M of L is orthogonal to W . Finally, let us consider a sequence of stopping times $(\tau_m)_{m \geq 1}$ such that the process $\sup_{n \geq 1} (Y^n - S) \mathbf{1}_{[0, \tau_m]}$ is essentially bounded and $\tau_m \rightarrow \infty$ as $m \rightarrow \infty$. Since $\|Y^n - Y\|_{\mathbb{S}^2} + \|L^n - L\|_{\mathbb{S}^2} \rightarrow 0$, it follows that

$$-\mathbb{E} \left[\int_0^{\tau_m} (Y_{t-} - S_{t-}) dK_t \right] = \mathbb{E} \left[\int_0^{\tau_m} (Y_{t-} - S_{t-}) dL_t \right] = \lim_{n \rightarrow \infty} \mathbb{E} \left[\int_0^{\tau_m} (Y_{t-}^n - S_{t-}) dL_t^n \right] = 0.$$

Since K is non-decreasing and $Y \geq S$ on $[0, T]$, we thus obtain

$$\int_0^{\tau_m} (Y_{t-} - S_{t-}) dK_t = 0, \quad \mathbb{P} - \text{a.s.}$$

Letting $m \rightarrow \infty$, we see that the Skorokhod condition (3.3) holds true for K .

2. Finally, let us consider the general case when $p \in (1, \infty)$. In this case, we can define for $n \geq 1$

$$\xi^n := (-n) \vee \xi \wedge n, \quad S^n := (-n) \vee S \wedge n, \quad \text{and} \quad g^n(\cdot) := (-n) \vee g(\cdot) \wedge n,$$

so that $(\xi^n, S^n, g^n(\cdot, 0, 0))_{n \geq 1} \in \mathbb{L}^2 \times \mathbb{S}_r^2 \times \mathbb{H}_1^{2,\alpha}$, for any $\alpha \geq 0$. Thus by Step 1, we know that there is a unique solution $(Y^n, Z^n, M^n, K^n) \in \mathbb{S}_r^2 \times \mathbb{H}^2 \times \mathbb{M}^2 \times \mathbb{I}_{+,r}^2$ to (3.1). Since $(\xi^n, S^n, g^n(\cdot, 0, 0))_{n \geq 1}$ is a Cauchy sequence in $\mathbb{L}^p \times \mathbb{S}_r^p \times \mathbb{H}_1^{p,\alpha}$, the estimates of Proposition 3.2 and Theorem 2.2 show that the sequence $(Y^n, Z^n)_{n \geq 1}$ is also a Cauchy sequence in $\mathbb{S}_r^p \times \mathbb{H}^{p,\alpha}$, for any $\alpha \geq 0$. Moreover, by a similar equality as in (3.5), it follows that $(L^n := M^n - K^n)_{n \geq 1}$ is also a Cauchy sequence in \mathbb{S}_r^p . Using the same arguments as in (iii) of Step 1, it is easy to check that its limit is a solution to (3.1). \square

4 Side remark

Note that the existence and uniqueness of a solution $(Y, Z, M) \in \mathbb{S}_r^p \times \mathbb{H}^p \times \mathbb{M}^p$ to the BSDE

$$Y_t = \xi - \int_t^T g_s(Y_s, Z_s) ds - \int_t^T Z_s \cdot dW_s - \int_t^T dM_s \quad (4.1)$$

follows from the same arguments as the one used in Section 3 whenever $\xi \in \mathbb{L}^p$ and $g^0 \in \mathbb{H}_1^p$. Indeed, we can bound the component Y of a solution as in the proof of Proposition 3.2 by

$$X_t Y_t = \mathbb{E}_t^{\mathbb{Q}} \left[- \int_t^T X_s g_s(0, 0) ds + X_T \xi \right], \quad \mathbb{P} - \text{a.s.},$$

in which X and \mathbb{Q} are defined as in (3.2). The difference of the Y -components of two solutions can be bounded similarly. Then, it suffices to apply the same fixed point argument as in the proof of Theorem 3.1.

Theorem 4.1. *Let Assumption 2.1 hold true. Then, (4.1) admits a unique solution in $\mathbb{S}_T^p \times \mathbb{H}^p \times \mathbb{M}^p$. Moreover, for all $\alpha \geq 0$, there exists a constant $C_{4.1}^\alpha$ that depends only on L_y, L_z and α , such that*

$$\left\| e^{\frac{\alpha}{2} \cdot Y} \right\|_{\mathbb{S}^p}^p \leq C_{4.1}^\alpha \left(\|\xi\|_{\mathbb{L}^p}^p + \mathbb{E} \left[\left(\int_0^T e^{L_y s} |g_s^0| ds \right)^p \right] \right).$$

Remark 4.1. *Recalling that the difference of two solutions, with different terminal conditions and drivers, is still a solution to a BSDE, the bound of Theorem 4.1 applies. Using similar notations as above, we have:*

$$\left\| e^{\frac{\alpha}{2} \cdot \delta Y} \right\|_{\mathbb{S}^p}^p \leq \bar{C}_{4.1}^\alpha \left(\|\delta \xi\|_{\mathbb{L}^p}^p + \mathbb{E} \left[\left(\int_0^T e^{L_y s} |\delta g_s(Y_s^1, Z_s^1)| ds \right)^p \right] \right).$$

A Appendix

Let us consider a strong supermartingale $X \in \mathbb{S}^p$ on $[0, T]$. Then, its paths are almost surely l  d  g, and it admits the (unique) Doob-Meyer decomposition (see e.g. Mertens [18]) :

$$X_t = X_0 + M_t - A_t - I_t, \tag{A.1}$$

where M is a right-continuous martingale, A is a predictable non-decreasing right-continuous process with $A_0 = 0$, and I is a predictable non-decreasing left-continuous process with $I_0 = 0$.

The following extends [16, Thm 3.1], which is the key ingredient of our main results, Theorem 2.1 and Theorem 2.2.

Lemma A.1. *For every constant $p > 1$, there is some constant $C_{A.1}^p > 0$ such that, for all strong supermartingale $X \in \mathbb{S}^p$ (with the decomposition (A.1)), one has*

$$\|A\|_{\mathbb{I}^p} + \|I\|_{\mathbb{I}^p} \leq C_{A.1}^p \|X\|_{\mathbb{S}^p}.$$

Proof. (i) We suppose in addition that X is right-continuous so that $I \equiv 0$. Denote $X^* := \sup_{0 \leq t \leq T} |X_t|$ and define \tilde{X} as the right-continuous version of the martingale $\mathbb{E}[X^* | \mathcal{F}_t]$. Then $\hat{X} := X + \tilde{X}$ is a positive right-continuous supermartingale on $[0, T]$, with the Doob-Meyer decomposition

$$\hat{X}_t = (\tilde{X}_t + M_t) - A_t.$$

Setting $\hat{X}_t := \hat{X}_T$ for $t \in [T, T+1)$ and $\hat{X}_t = 0$ for $t \in [T+1, \infty)$, then \hat{X} is in fact a right-continuous potential (recall that a potential is a non-negative right-continuous supermartingale \hat{X} on $[0, \infty)$ such that $\lim_{t \rightarrow \infty} \mathbb{E}[\hat{X}_t] = 0$). Using Meyer [19, Thm 1] (see also [16, Thm 3.1]), there is some constant C'_p such that

$$\|A\|_{\mathbb{I}^p} \leq C'_p \left\| \hat{X} \right\|_{\mathbb{S}^p}.$$

By the definition of \hat{X} and Doob's martingale inequality, we get

$$\|A\|_{\mathbb{I}^p} \leq C'_p \left\| \hat{X} \right\|_{\mathbb{S}^p} \leq C'_p \left(1 + \frac{p}{p-1} \right) \|X\|_{\mathbb{S}^p}.$$

(ii) We now consider the case when X is not necessary right-continuous. By Mertens [18], we know that the process $X + I$ is a right-continuous strong supermartingale, and that the left-continuous process I is obtained as the limit of an increasing sequence $I := \lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} I^{\varepsilon, n}$, where $I^{\varepsilon, n}$ is defined by

$$I_t^{\varepsilon, n} := \sum_{k=1}^n (X_{\sigma_k^{\varepsilon, n}} - X_{\sigma_k^{\varepsilon, n}+}) \mathbf{1}_{\sigma_k^{\varepsilon, n} < t},$$

in which $(\sigma_k^{\varepsilon, n})_{1 \leq k \leq n}$ is the non-decreasing sequence of stopping times which exhausts the first n jumps of X such that $X_t - X_{t+} \geq \varepsilon$. In Step (iii) we will show that $\|I^{\varepsilon, n}\|_{\mathbb{I}^p} \leq C_p'' \|\hat{X}\|_{\mathbb{S}^p}$ for some constant $C_p'' > 0$ independent of ε and n , then it follows from the monotone convergence theorem that

$$\|I\|_{\mathbb{I}^p} \leq C_p'' \|\hat{X}\|_{\mathbb{S}^p} \leq C_p'' \left(1 + \frac{p}{p-1}\right) \|X\|_{\mathbb{S}^p},$$

and hence

$$\|A\|_{\mathbb{I}^p} \leq C_p' \left(1 + \frac{p}{p-1}\right) \|X + I\|_{\mathbb{S}^p} \leq C_p' \left(1 + \frac{p}{p-1}\right) (1 + C_p'') \|X\|_{\mathbb{S}^p}.$$

(iii) It is now enough to prove that $\|I^{\varepsilon, n}\|_{\mathbb{I}^p} \leq C_p'' \|X\|_{\mathbb{S}^p}$ for some constant C_p'' independent of (n, ε) . Notice that the discrete process $\bar{X}^n := (X_0, X_{\sigma_1^{\varepsilon, n}}, X_{\sigma_1^{\varepsilon, n}+}, \dots, X_{\sigma_n^{\varepsilon, n}}, X_{\sigma_n^{\varepsilon, n}+})$ is a discrete time supermartingale. By interpolation, we can turn it into a right-continuous strong supermartingale on $[0, T]$. Then, using the results in Step (i) we obtain that

$$\|I^{\varepsilon, n}\|_{\mathbb{I}^p} \leq C_p'' \|X\|_{\mathbb{S}^p}, \quad \text{with } C_p'' := C_p' \left(1 + \frac{p}{p-1}\right).$$

□

Remark A.1. A careful reading of the proof in [19] shows that the constant can be computed explicitly and is given by

$$C_p' := \min_{2 \leq k < p} \left(p \prod_{j=2}^k \frac{pj}{p-j} \right)^{\frac{k}{p-1}} \mathbf{1}_{p>2} + \left(\frac{p^2}{p-1} \right)^{\frac{1}{p-1}} \mathbf{1}_{p \in (1, 2]}.$$

Meyer's result show that for a supermartingale X with Doob-Meyer's decomposition $X = M - A$, we can control A by X ; the following example shows that given two supermartingales, we cannot control the difference of the A parts by the difference of the supermartingales.

Example A.1. Let W be a one dimensional Brownian motion. Fix $\varepsilon > 0$ and let V be defined by

$$V_t := \sum_{k \geq 0} W_{\tau_k} \mathbf{1}_{[\tau_k, \tau_{k+1})}(t), \quad \text{where } \tau_0 := 0, \quad \tau_{k+1} := \inf\{t \geq \tau_k : |W_t - W_{\tau_k}| \geq \varepsilon\}.$$

Notice that V is of finite variation with decomposition $V = V^+ - V^-$ such that V^+ and V^- are two non-negative non-decreasing and predictable process. Let $X^1 := W - V^+$ and $X^2 := -V^-$, then $\sup_t |X_t^1 - X_t^2| = \sup_t |W_t - V_t| \leq \varepsilon$, but $V^+ - V^- = V$ cannot be controlled by ε . □

We finally provide a technical lemma used in the paper. Recall the definition of ϕ_p in (2.7) and observe that $|\phi_{p-1}(y)| = |y|^{p-2} \mathbf{1}_{y \neq 0}$.

Lemma A.2. *Let X be a l  dl  g semimartingale. Then for all $p \in (1, 2)$ and $\alpha > 0$, we have \mathbb{P} – a.s. for any $t \in [0, T]$*

$$\begin{aligned} e^{p\frac{\alpha}{2}t} |X_t|^p &\leq e^{p\frac{\alpha}{2}T} |X_T|^p - p \int_t^T e^{p\frac{\alpha}{2}s} \frac{\alpha}{2} |X_s|^p ds - p \int_t^T e^{p\frac{\alpha}{2}s} \phi_p(X_{s-}) dX_s \\ &\quad - \frac{p(p-1)}{2} \int_t^T e^{p\frac{\alpha}{2}s} |\phi_{p-1}(X_s)| d[X]_s^c \\ &\quad - \frac{p(p-1)}{2} \sum_{t < s \leq T} e^{p\frac{\alpha}{2}s} |X_{s+} - X_{s-}|^2 \left(|X_{s-}|^2 \vee |X_{s+}|^2 \right)^{\frac{p}{2}-1} \mathbf{1}_{|X_{s-}| \vee |X_{s+}| \neq 0}, \end{aligned}$$

where we denote $X_{T+} := X_T$.

Proof. This is an immediate consequence of a straightforward adaptation of [13, Lemmas 7, 8 and 9], together with the It  ’s formula for l  dl  g processes in [15, p.538]. \square

References

- [1] Bouchard, B.,   lie, R., R  veillac, A. (2015). BSDEs with weak terminal condition, *Ann. Probab.*, 43(2):572–604.
- [2] Bouchard, B., Possama  , D., Tan, X. (2015). A general Doob-Meyer-Mertens decomposition for g-supermartingale system, *arXiv preprint, arXiv:1505.00597*
- [3] Briand, P., Delyon, B., Hu, Y., Pardoux,   ., Stoica, L. (2003). L^p solutions of backward stochastic differential equations, *Stoch. Proc. Appl.*, 108:109–129.
- [4] Cvitani  , J., Karatzas, I., Soner, H.M. (1998). Backward stochastic differential equations with constraints on the gain-process, *Ann. Probab.*, 26:1522–1551.
- [5] El Karoui, N. (1981). Les aspects probabilistes du contr  le stochastique, *Ecole d’Et   de Probabilit  s de Saint-Flour IX-1979, Lecture Notes in Mathematics*, 876:73–238.
- [6] El Karoui, N., Huang, S.-J. (1997). A general result of existence and uniqueness of backward stochastic differential equations, *In Backward stochastic differential equations (Paris, 1995-1996), volume 364 of Pitman Res. Notes Math. Ser.*, pp. 27–36. Longman, Harlow.
- [7] El Karoui, N., Peng, S., Quenez, M.-C. (1997). Backward stochastic differential equations in finance, *Mathematical Finance*, 7(1):1–71.
- [8] El Karoui, N., Kapoudjian, C., Pardoux, E., Peng, S., Quenez, M.-C. (1997). Reflected solutions of backward SDE’s, and related obstacle problems for PDE’s, *Annals of Applied Probability*, 25(2):702–737.
- [9] Grigorova, M., Imkeller, P., Offen, E., Ouknine, Y., Quenez, M.-C. (2015). Reflected BSDEs when the obstacle is not right-continuous and optimal stopping, *arXiv preprint arXiv:1504.06094*.
- [10] Klimsiak, T. (2013). BSDEs with monotone generator and two irregular reflecting barriers, *Bulletin des Sciences Math  matiques*, 137(3):268–321.

- [11] Klimsiak, T. (2014). Reflected BSDEs with monotone generator, *Electron. J. Probab.*, 17(107):1–25.
- [12] Klimsiak, T. (2014). Reflected BSDEs on filtered probability spaces, *arXiv preprint, arXiv:1408.5598*.
- [13] Kruse, T., Popier, A. (2015). BSDEs with monotone generator driven by Brownian and Poisson noises in a general filtration, *arXiv preprint, arXiv:1412.4622*.
- [14] Lepeltier, J. P. and Xu, M. (2005). Penalization method for reflected backward stochastic differential equations with one r.c.l.l. barrier, *Stat. and Prob. Lett.*, 75(1):58–66.
- [15] Lenglart, É. (1980). Tribus de Meyer et théorie des processus. *Séminaire de probabilité*, 14:500–546.
- [16] Lenglart, É., Lépine, D., Pratelli, M. (1980). Présentation unifiée de certaines inégalités de la théorie des martingales, *Séminaire de probabilités (Strasbourg)*, 14:26–48.
- [17] Lin, Q., Peng, S. (2000). Smallest g -supersolutions for BSDE with continuous drift coefficients, *Chin. Ann. of Math.*, 21B(3):359–366.
- [18] Mertens, J.-F. (1972). Théorie des processus stochastiques généraux applications, aux surmartingales. *Probability Theory and Related Fields*, 22(1):45–68.
- [19] Meyer, P.-A. (1968). Une majoration du processus croissant naturel associé à une surmartingale, *Séminaire de probabilités (Strasbourg)*, 2:166–170.
- [20] Osękowski, A. (2012). Sharp martingale and semimartingale inequalities, *Birkhäuser*.
- [21] Peng, S. (1999). Monotonic limit theorem of BSDE and nonlinear decomposition theorem of Doob-Meyer’s type, *Probability Theory and Related Fields*, 113(4):473–499.
- [22] Peng, S., Xu, M. (2005). The smallest g -supermartingale and reflected BSDE with single and double L^2 obstacles, *Ann. I. H. Poincaré - PR*, 41:605–630.
- [23] Peng, S., Xu, M. (2007). Constrained BSDE and viscosity solutions of variation inequalities, *arXiv preprint, arXiv:0712.0396*.
- [24] Peng, S., Xu, M. (2010). Reflected BSDE with a constraint and its applications in an incomplete market, *Bernoulli*, 16(3):614–640.
- [25] Possamai, D., Tan, X., Zhou, C. (2015). Stochastic control for a class of nonlinear stochastic kernels and applications, *in preparation*.
- [26] Soner, M., Touzi, N., Zhang, J. (2012). Wellposedness of second order backward SDEs, *Probability Theory and Related Fields*. 153(1-2):149–190.